

B e r i c h t e  
aus dem  
Institut für Meereskunde  
an der  
Christian-Albrechts-Universität Kiel

No 55

A PRELIMINARY<sup>+</sup> COMPILATION OF  
FISH LENGTH GROWTH PARAMETERS

by

Daniel Pauly

DOI 10.3289/IFM-BER-55

Copies of this compilation  
may be obtained from:  
Institut für Meereskunde  
Abt. Fischereibiologie  
Düsternbrooker Weg 20  
2300 Kiel, BRD

---

<sup>+</sup>This compilation contains a part of the data gathered for  
use in my dissertation, presently in preparation, and in which  
the quality of the data presented here will be assessed.

T A B L E   O F   C O N T E N T S

<u>Part I</u>	Page
1.1.,      Zusammenfassung .....	3
1.2.      Abstract .....	4
1.3.      Introduction .....	5
1.4.      Scope of compilation .....	5
1.5.      Source of data .....	8
1.6.      Methods used for the determination of growth parameters .....	10
1.6.1.,   Preliminary remarks .....	10
1.6.2.,   Methods (1 to 5) .....	13
1.6.2.1   von Bertalanffy Plot .....	13
1.6.2.2   Ford Walford Plot .....	13
1.6.2.3   Gulland and Holt Plot .....	13
1.6.2.4   Non-Linear Regression .....	14
1.6.2.5   "Ageing" Length Frequencies .....	14
1.7.      Presentation of the data .....	16
1.7.1.    General .....	16
1.7.2.    Taxonomic questions pertaining to data presentation .....	17
1.7.3.    Conversion to uniform length measurements	18
1.8.      Reliability of data .....	18
1.9.      Acknowledgement .....	19
1.10.     Bibliography I (References of Part I)	20
<u>Part II</u> .....	23
2.1.      Alphabetic Index to families .....	23
2.2.      Data by families: Acanthuridae .....	27
to Zoarcidae .....	117

	Page
2.3. Controversial data, Order Selachoides ...	118
2.4. Bibliography II .....	119
2.5. Addenda to bibliography II .....	198

### List of Tables

I	Summary of data included in previous compilations as compared with present compilation .....	6
II	Relationship of the number of taxa listed to number of recent fish taxa .....	8
III	Source or method used for the determination of the growth parameters included in this compilation .....	10
IV	Relative contribution of the various methods	12

### List of Figures

Figure 1	Geographical distribution of units included in this compilation .....	7
Figure 2	Example for method 5: length frequency data for <u>Sardinella sirm</u> from the Java Sea ..	15

## Part I

### 1.1. Zusammenfassung

Die Längenwachstumsparameter von 515 verschiedenen Fischarten, verteilt in 1 501 verschiedenen Beständen werden vorgestellt. Diese Parameter sind  $L_{\infty}$  und  $K$  (sowie  $t_0$  in zahlreichen Fällen) der von Bertalanffy Wachstumsfunktion. Die Parameterwerte wurden mit Hilfe verschiedener Methoden in 966 Fällen aus "Literaturrohdaten" wie Längen-Alter Wertepaare und verwandte Daten neu berechnet, der Rest stammt aus früheren, ähnlichen Zusammenstellungen und von der Primärliteratur. Die Methoden, die für die Berechnung der Parameter eingesetzt wurden, werden kurz charakterisiert. Sie werden in einer späteren Veröffentlichung diskutiert werden.

Die Zusammenstellung umfaßt vor allem Seefische aus temperierten Gebieten, aber es wurde darauf geachtet, so viele tropische Seefische einzubeziehen, wie nur möglich. Nur 12 % der aufgeführten Arten sind rein limnisch.

Die Qualität der aufgeführten Werte ist ziemlich schwankend, und sie sollten möglichst im Zusammenhang mit den Originalwerten betrachtet werden, sowie mit Daten aus verwandten Beständen.

Eine Bibliographie mit mehr als 650 Referenzen, die Fragen des Fischwachstums behandeln, schließt diese Zusammenstellung ab.

## 1.2. Abstract

The length growth parameters of 515 different species of fish, distributed in 1 501 different stocks are presented. The parameters are  $L_{\infty}$  and  $K$  (as well as  $t_0$  in many cases) of the von Bertalanffy growth function. They have been determined by various methods from "raw" size-at-age and related literature data in 966 cases, the rest originating from previous similar compilation and from the primary literature. The methods used for parameters estimation are briefly discussed, pending a detailed presentation elsewhere.

The compilation covers mainly temperate marine fishes, but care was taken to include as many tropical marine fishes as possible. Only 12 % of the species listed are purely limnic.

The parameter values listed are of varying quality and should be used preferably in the light of the original growth data and of data from related stocks.

A bibliography of more than 650 papers with data pertaining to fish growth is given.

### 1.3.

Previous compilations of fish growth parameters, especially those of Beverton & Holt (1959), Holt (1959, 1960), Beverton (1963), Taylor (1958) and Ursin (1967) have provided a great deal of insight into some of the mechanisms of growth in fishes.

In January 1977, I began a compilation of data on fish growth with the aim of identifying a general pattern to which the growth of all fish would conform and which could be used to predict the growth parameters of little-known fish stocks.

Such a general pattern does indeed exist and it will be exposed in my dissertation, presently in preparation,

The growth parameters calculated and/or used for the identification of this mechanism, however, do in themselves represent a fair amount of work and could be quite valuable to fishery biologists, especially those doing field work in areas where library and other facilities are not optimal.

### 1.4. Scope of the compilation

Table I should provide an idea of the scope of the present compilation, as compared with previous compilations.

Table II relates the number of taxa included here to the number of recent fish taxa. It will be noticed that, while 24 % of all families and 10 % of all recent genera are included, the bibliography covers only 2.4 % of all existing species. It should be realized, however, that literally all commercially relevant species are included here: it is those fish populations supporting relevant fisheries which get investigated first, in all countries.

Thus any good scanning of the literature will result in a very good coverage of those species which are commercially

Table I: Summary of data included in previous compilations of fish growth parameters, as compared with present compilation. ("Units" are pairs of value for  $L_{\infty}$  & K, "sexed units" are units relating to a given sex.)

No.	Group(s) of fish investigated	No of Genera	No. of Spp.	No. of Units	No of sexed Units	Author(s)
1	misc. marine & freshwater	44	58	88	32	Beverton & Holt 1959
2	Scombridae	2	4	14	0	Holt 1959
3	Sardines	3	10	31	6	Holt 1960
4	Clupeidae & Engraulidae	7	13	51	0	Beverton 1963
5	Sciaenidae	11	13	13	0	Rao 1966
6	Engraulidae	3	7	75	0	Bayliff 1957
7	Cod, <u>Gadus morhua</u>	1	1	11	0	Taylor 1958
8	misc. marine & freshwater	63	81	81	13	Ursin 1967 <sup>+</sup>
9	Trout, <u>Salmo trutta</u>	1	1	11	0	Gelhaar 1972
10	misc. marine, India	13	18	18	0	Banerji & Krishnan 1973
11	Cod, sprat and herring, Baltic	3	3	31	0	Thurøw 1974
12	misc. marine & freshwater	300	515	1501	332	Pauly 1978

<sup>+</sup>gives estimates of asymptotic weight

important. Fig. 1 shows the overall geographic distribution of the stocks whose growth parameters are included here.

The distribution pattern that is apparent in Fig. 1 requires little comment. In the case of the marine fishes, Fig. 1 is a quite truthful indicator of the general lack of knowledge on the growth of tropical stocks. And I believe that the situation would have been similar had I focused on fresh water stocks. On the other hand, the total number of stocks included here could have been doubled in a relatively short period of time, by simply recording more data on stocks from the Northern Atlantic, where practically all commercially relevant stocks have been repeatedly investigated.




Fig. 1 Geographical distribution of units.

1 small dot = 1 unit, 1 large dot = 10 units.



Table II: Relationship of the numbers of taxa listed in this compilation with numbers of recent fish taxa.

Recent taxa		listed	extant	listed in % of ext.	Source of estimates
Families	lower fishes	14	34	41.2	Berg 1958
	Teleosteans	93	417	22.3	Greenwood <u>et al.</u> 1966
	total	107	447	23.9	
Genera	lower fishes	36	160	22.5	after Jordan, 1963 <sup>+</sup>
	Teleosteans	264	2 840	9.3	" " "
	total	300	3 000	10.0	
Species	lower fishes	50	650	7.7	after Cohen, 1970
	Teleosteans	455	21 000	2.2	" " "
	total	515	21 650	2.4	

<sup>+</sup>discounting as probable synonyms 50 % of his generic names and rounding off the resulting figures.

### 1.5. Source of data

- a) A first source of data for this compilation were obviously the previous compilations listed in Table I. The data originating from this source generally have the code number "00" under the "method" heading. However, in most cases, the name(s) of the author(s) of the original data is (are) given under the "source" (of data) heading, this in order to enable users of this compilation to properly identify the stock in question.
- b) A second source of data were review papers and books in which "raw" growth data, mainly length-at-age data had been compiled for descriptive or comparative purposes. Such sources were for example the papers by D'Ancona (1937) or Bougis (1952) and especially

Nikolsky's "Special Ichthyology" (1957). In the case of this type of source it is, as in case a) the original author who was quoted - even if he was not consulted in the original - rather than the compiler (with the exception of the soviet authors cited by Nikolsky and whose papers are generally not accessible here). Again, the aim was to enable the user to track down the original growth data,

This feature seems very important as it appeared very early in the course of this work that many sets of growth data cited in different papers went back to one same author and/or fish stock,

- c) The third and most important source of growth data, however, was the primary literature, which was scanned quite thoroughly - at least for certain taxa and/or regions (see Fig. 1).

In those cases where the authors had calculated growth parameters themselves, this is indicated by a "O" under the "method" heading. In those cases where the parameters had not been calculated - that is in the majority of cases - this was done by using one of the methods listed in Table III, and the code number given correspond to the method used. In many cases, it was necessary to even reinterpret growth data, i.e. where authors had fitted their length-at-age data by straight lines (see for example Hureau 1970, Hureau & Ozouf 1977).

Thus, we summarize: All data with "O" and "OO" under the "method" heading are parameters taken from the literature, all other ones (64.4 %) are original values, determined by means of the method whose code number is given.

Table III

Source or method used for the determination of the growth parameters presented in this compilation.

	Code Nr.	Nr. of units
1) Growth parameter estimated by author given as reference	0	346
2) Growth parameter estimated by author different from the one cited (i.e. originating from one of the previous compilations in Table I)	00	190
3) "Von Bertalanffy Plot"	1	23
4) "Ford-Walford Plot"	2	158
5) "Gulland & Holt Plot"	3	12
6) Gulland & Holt Plot, with set value for asymptotic size	31	13
7) Non-Linear-Regression	4	604
8) Non Linear Regression, with set value for asymptotic size	41	104
9) "Ageing" length frequency data	5	42
10) Ageing length frequency data, with set value for asymptotic size	51	9
		<hr/> 1 501

#### 1.6. Methods used for the determination of growth parameters

##### 1.6.1. Preliminary remarks:

The work on this compilation gave ample opportunity to critically assess the various methods used in the determination of growth parameters. Some of them could be expanded and/or made more flexible. Also a new technique for the analysis of size-frequency samples was devised which allows for better reproducibility of results.

All of these methods, critiques and new techniques will be presented in a next issue of the "Berichte des Instituts für Meereskunde", together with programmes for their quick computation.

It seems therefore superfluous to do here more than listing and briefly outlining the characters of these methods.

The Von Bertalanffy Growth Formula (VBGF) in the notation of Beverton & Holt (1959) has for length growth the form

$$L_t = L_{\infty} (1 - e^{-K(t-t_0)})$$

The VBGF has been widely used in fisheries science, although it has been demonstrated that some of the assumptions made by von Bertalanffy (1934) do not apply (see for example Ursin 1967, Taylor 1962).

The VBGF, on the other hand, always produces an excellent fit to that part of a growth curve which corresponds to the exploited phase of fish stocks.

For this reason, and for the sake of comparability with previously published data, I have found it preferable not to use the four-parameters, generalized version of the VBGF, although it produces an even better fit and allows for a physiological interpretation of the parameters  $L_{\infty}$  &  $K$  (Richard 1959, Taylor 1962, Ursin 1967, Pauly, diss. in prep.)

The parameters of the VBGF may be defined as follow:

- 1) The asymptotic length, or  $L_{\infty}$  is about equal to the mean length the fish of a given stock would reach if they were to grow to a very old age. This definition differs from that of Ricker (1958) in that I insist that  $L_{\infty}$  should be a biologically "possible" length, not a mere mathematical entity. As will be shown elsewhere, this definition is of greater heuristic value than that of Ricker (1958) when the growth parameters of different fish stocks are to be compared.

In small and middle-sized fishes, however, the values of  $L_{\infty}$  are only slightly higher than the maximum size actually recorded ( $L_{\max}$ ). This feature was noted by Holt (1960), Beverton (1963) and especially Taylor (1958). Latter author suggested that, as a rule

$$L_{\max} \approx L_{\infty} \times 0.95$$

Table IV: Relative contribution of the various methods used for this compilation.

Code No of method	published growth parameters		original growth parameters							
	0	00	1	2	3	31	4	41	5	51
No of units	346	190	23	159	12	13	604	103	42	9
% of orig. values	--	--	2.4	16.5	1.2	1.4	62.5	10.7	4.4	0.9
% of all values	35.6 ( $\Sigma = 535$ )		64.4				( $\Sigma = 966$ )			

This property of  $L_{\max}$  has been used in this compilation to obtain preliminary estimates of asymptotic length in certain stocks. Such values I suggest to code " $L_{(\infty)}$ " in order to prevent confusion with calculated values of  $L_{\infty}$ .

Values of  $L_{(\infty)}$  must always be used, obviously, with those methods requiring independent estimates of asymptotic size (that is methods 1, 31, 41, and 51). Most of the values of  $L_{\max}$  used here come from the author who also produced the growth data.

- 2) The constant K has the dimension 1/time, and is here always expressed as 1/year. An exhaustive discussion of its character and properties will be given elsewhere.
- 3) The parameter  $t_0$  is "the age the fish would have had at length zero had they always grown in the manner described by the equation" (Ricker, 1958).  
Note that this definition implies that  $t_0$  cannot be interpreted biologically unless it is demonstrated that the fishes do always grow according to the equation.

This applies to the length at age zero which should not be interpreted as length at hatching or birth.

#### 1.6.2. Methods 1 to 5

1.6.2.1 Method 1, called here the "von Bertalanffy Plot" has been proposed by von Bertalanffy (1934, p. 627) for the estimation of  $K$  and by Beverton & Holt (1957, p. 283, fig. 16.4) and Ricker (1958) for the estimation of  $K$  and  $t_0$  on the base of a preliminary estimation of  $L(\infty)$ . The method has the advantage that the length-at-age samples can be weighed according to their size, and that data at unequal time interval can be used.

1.6.2.2 Method 2 is the well-known "Ford-Walford Plot" (Ford, 1933, Walford, 1946) which can be used for data at equal time interval only. For statistical reasons to be discussed in the planned manual of methods, a "functional" rather than a "predictive" regression has been fitted to the plot of  $L_t + 1$  against  $L_t$  data (see Ricker (1973), for a definition of the regression types). An exhaustive discussion of the FW Plot and its variants will be given elsewhere.

1.6.2.3 Method 3 is the "Gulland & Holt Plot" proposed for the treatment of data at unequal time interval (Gulland and Holt, 1959). The method has the code number 3. In certain cases a set value of  $L(\infty)$  used in conjunction with the "Gulland and Holt Plot" has proven necessary and helpful. This method, called here "Simplified Gulland & Holt Plot" has the code number 31.

1.6.2.4 Method 4 is the direct estimation of the growth parameters by means of a non-linear regression. Methods and programmes for the direct estimations of growth parameters have been introduced to fishery biology by Tomlinson & Abramson (1961), Fabens (1965) and Allen (1966). The programme used here was kindly written (in Algol) for the author by Dr. G. Gaschütz and was run on the Digital PDP 10 of the computer center of Kiel University. This programme, coded 4, estimates growth parameters from length or weight-at-age data, both at equal or unequal time intervals and the various data can be weighted by sample size. This latter feature allows for the use of a value of  $L_{(\infty)}$  or  $W_{(\infty)}$  which is input as a size at a very large age (say 500 years) and given a very high weight in term of sample size. This latter method is coded 41.

The programme and instructions for its use will be presented in the planned manual of methods.

#### 1.6.2.5 "Ageing" length frequencies

The two techniques generally used for the determination of values of  $L_{\infty}$  and  $K$  from length frequency data are:

- 1) the Petersen method
- 2) the "modal class progression analysis".

With the first method, assumptions are to be made on the time interval which separates various peaks assumed to represent certain age groups.

With the second method, assumptions are to be made as to which of the peaks can be interconnected that belong to two or several samples separated by (a) known time interval(s).

Thus, for example, questions to ask would be for the samples in figure 2A:

- method 1) - which time interval is represented by the length difference between the two peaks of the May sample?





Fig. 2 A. Length frequency data on *Sardinella sima* as presented by Burhanuddin *et al.* (1974)(copy of their Fig.2 p.20)  
 B. Same data as in Fig. 2A, replotted according to the integrated method.

Note special features:

- 1) The intervals on the time axis between the various samples are proportional to the time elapsed between the various sampling dates (c.f. Fig. 2A).
- 2) The original data are plotted twice along the time axis ("doubling up"). This allows for longer growth curves to be drawn. Note also that the two February samples coincide well.
- 3) All growth curves are drawn with a curved ruler and have approximately the same shape, varying only as to their origin.
- 4) The scale of the ordinate (length) starts at zero, thus allowing for the approximate birth date to be traced back (c.f. Fig. 2A).
- 5) Each growth curve interconnects several peaks; the more peaks a curve interconnects, the more likely it is to depict the actual growth of the population.
- 6) The length-at-age can be read off the curves at equal time intervals (say months), and these data plotted according to Ford-Walford (method 2). The various curves may yield slightly different estimates of  $L_{\infty}$  and  $K$ . In such case, geometric means should be taken.





method 2) - which of the three peaks of the November sample may be connected with the single January peak?

In most cases, the two questions of our example cannot be answered separately. Indeed, it appears that the two techniques may be combined into a new integrated method for "ageing" length frequency, where the question to ask may be formulated as follow:

- which combination of values of  $L_{\infty}$  and  $K$  produces a growth curve which "explains" - that is, interconnects - most of the peaks of a series of sequentially arranged length frequency samples?

The actual technique to use for the determination of the appropriate values of  $L_{\infty}$  and  $K$  is described in details on page 15, Fig. 2B.

An interesting feature of the integrated method is that it can be readily programmed. A computer programme for the processing of length and weight frequencies, including data where the growth rate oscillates seasonally is presently in preparation.

## 1.7. Presentation of data

### 1.7.1. General

The growth parameters are presented in "units", one unit containing at least a species name, a value of  $L_{\infty}$  (or  $L_{(\infty)}$ ) and of  $K$ , with a reference and a code number under the "source", resp. the "method" heading. In addition, a sampling location, more or less precise, is given, and/or additional comments under the "note" heading. A frequent note is a value of  $M$  (exponential coefficient of natural mortality), which in most cases originate from the papers by Beverton and Holt (1959) and Beverton (1963), but may also come from papers not cited but referring to the stock in question.

### 1.7.2. Taxonomic questions pertaining to data presentation

Quite often, the fishes whose growth parameters were used or determined belong to taxa whose valid names have not yet been agreed upon. I have assumed that the law of priority does not affect fish growth and stuck to the following rules :

- 1) Subspecific taxa were not considered and their names dropped from the specific names. Hence, for example, Notothenia rossi marmorata (fam. Notothenidae) is listed here as Notothenia rossi.
- 2) No serious attempts were made to check the validity of the specific and generic names used by the various authors. The taxa I put in synonymy are those I happened to know about; as a whole I have probably been more a "lumper" than a "splitter".
- 3) The genera recognized were ordered into families by means of Jordan's "Classification of Fishes" (1963) but the only families considered valid are those of Berg (1958) for the lower fishes and of Greenwood et al. (1966) for the Teleosteans, with the exception of the Thunnidae and Scombridae which were kept separate. Thus, for example, the data of Eisele (1943) on Coregonus wartmanni, to be found here under Coregonus lava-retus, are listed in the salmonid family, since the coregonid family, listed in Jordan (1963) is considered a subfamily of the Salmonidae by Greenwood et al. (1966).
- 4) The families, finally are listed alphabetically (see index p. 23). The only exceptions pertain to data by Holden (1974) on the order Selachioidei which are all grouped on one page (118), because of their controversial nature. (Holden (1974) interpreted  $t_0$  biologically, as the duration of gestation period, and the length-at-age zero as birth length).

### 1.7.3. Conversion to uniform length measurements

A considerable amount of problems have been caused by the need to use, throughout this compilation the same type of length measurements. Most authors do state the type of length they measured (standard, total, fork, eye-to-fork etc.). Some however, do not. Still there was a need to convert the data to uniform length measurements. The convention used here is the total length, except for the large scombroids such as tuna and billfishes, where fork length was used. Obviously, caudal filaments and the like have not been considered. Conversions from other lengths were made by using conversion factors given by the authors themselves, or from figures depicting the fishes in question. I have no doubts that, regrettably, these conversion problems are one of the main source for the errors which may occur in this compilation.

A few of the length growth parameters were derived from estimates of weight growth parameters, especially in the case of the clupeid Brevoortia tyrannus. Such conversions were done on the assumption that growth is isometric. A discussion of the value of this assumption will be given elsewhere.

### 1.8. Reliability of data

The growth parameters listed here, although in many cases calculated with the most sophisticated method presently available (Non-Linear Regression) should in any case be used with great circumspection.

The value of  $L_{\infty}$  (and consequently of  $K$ ) derived from a given set of length-at age data are strongly affected by biased sampling, by sex specific growth, by seasonal growth oscillation (especially in short-lived species), to mention only a few of the problems occurring.

The methods used provide for parameter estimates of varying quality. In term of reliability and goodness of fit they may be ordered as follow: 4, 2, 3, 5 and 41, 1, 51, 31, the second group being as a whole less reliable than the first.

The data of this compilation, it should be kept in mind do not represent a list of confirmed growth parameters, and are at the present stage nothing but raw data to be used in future comparative studies of fish growth. The parameters values presented here are for general orientation, and need to be checked in the light of the original growth data, and of data from related stocks everytime they appear controversial.

#### 1.9. Acknowledgement

My sincere thanks go to my supervisor Prof. Dr. G. Hempel for his support and guidance during the course of this work. I would also like to thank my friend Götz Gaschütz for providing me with the powerful, yet versatile tool that his programme for non-linear regression represents and which he offered me before I even knew that such things existed.

My gratitude also goes to Sandra Wade who turned my handwritten bibliography and notes into a readable whole, and to Mrs. Rockel for critically typing the manuscript. I would like to thank Uli Damm for discussing with me various aspects of, and contributing to this compilation, and, finally Prof. Dr. Daget of the Musée National d'Histoire Naturelle, Paris, for permission to work in the library of that Institution.

Bibliography I (References of Part I)

- ALLEN, K. R. 1966: A method of fitting growth curves of the Bertalanffy type to observed data. J. Fish. Res. Bd. Can. 23 (2): 163-179.
- BANERJI, S. K. and T. S. KRISHNAN 1973: Acceleration of assessment of fish populations and comparative studies of similar taxonomic groups, pp. 158-175 in: Proceedings of the Symposium on Living Resources of the Seas Around India, Spec. Publ., Cen. Mar. Fish. Res. Inst., Cochin, 748 pp.
- BAYLIFF, W. 1967: Growth, mortality, and exploitation of the Engraulidae, with special reference to the Ancho-veta, Cetengraulis mysticetus, and the colorado, Anchoa naso, in the Eastern Pacific Ocean. Int. Am. Trop. Tuna Comm. Bull. 12 (5): 365-432.
- BERG, L. 1958: System der rezenten und fossilen Fischartigen und Fische. VEB Verlag der Wissenschaften, Berlin (GDR) 310 pp.
- BERTALANFFY, L. von 1934: Untersuchungen über die Gesetzmäßigkeit des Wachstums. I Allgemeine Grundlagen der Theorie, mathematische und physiologische Gesetzmäßigkeiten des Wachstums bei Wassertieren. Roux' Archiv für Entwicklungsmechanik 131: 613-652.
- BEVERTON, R. J. H. 1963: Naturation, growth and mortality of Clupeid and Engraulid stocks in relation to fishing. Cons. Perm. Expl. Mer, Rapp. et Proc.-Verb., 154: 44-67.
- BEVERTON, R. J. H. & S. J. HOLT 1957: On the Dynamics of Exploited Fish Populations. Fish. Invest. Ser. II 19: 533 pp.
- BEVERTON, R. J. H. and S. J. HOLT 1959: A review of the lifespans and mortality rates of fish in nature, and their relation to growth and other physiological characteristics. CIBA Foundation Colloquia on Ageing 5: 142-180.
- BOUGIS, P. 1952: La croissance des poissons Méditerranéens. Vie et Milieu, Suppl. 2: 118-146.
- BURHANUDDIN, M. HUTOMO, S. MARTOSEWOJO & A. DJAMALI 1974: Some biological aspects of lemuru, Sardinella sirm (Waldbaum), around Panggang Islands In Indoneasian, with English abstract. Oceanologi di Indonesia 1974 (2): 17-25.
- COHEN, D. M. 1970: How many recent fishes are there? Proceedings of the California Academy of Sciences, 38 (17): 341-346.

- D'ANCONA, U. 1937: La croissance chez les animaux méditerranée Rapp. et P.-V. Comm. Int. Expl. Médit., X, pp. 162-224.
- EISELE, W. 1943: Über das Wachstum der Coregonen im Voralpengebiet, insbesondere über das Verhältnis von Schuppen- und Längenwachstum. Z. Fisch. und d. Hilfswiss. 61 (1): 23-45.
- FABENS, A. J. 1965: Properties and fitting of the von Bertalanffy growth curve. Growth 29: 265-289.
- FORD, E. 1933: An account of the herring investigations conducted at Plymouth during the years from 1924-1933. J. Mar. Biol. Ass. U.K. 19: 305-384.
- GEHLHAAR, C. E. 1972: Beiträge zur Biologie der Meeresforelle (Salmo trutta f. trutta L. in Schleswig-Holstein unter besonderer Berücksichtigung der Farver Au und der Rantzau. Thesis, Kiel University, 109 pp.
- GREENWOOD, P. H., D. E. ROSEN, S. H. WEITZMAN and G. S. MYERS 1966: Phyletic Studies of Teleostean Fishes, with a Provisional Classification of Living Forms. Bull. Am. Mus. Nat. History 131 (4): 339-456.
- GULLAND, J. A. & S. J. HOLT 1959: Estimation of growth parameters for data at unequal time intervals. J. du Conseil 25 (1): 47-49.
- HOLT, S. J. 1959: Report of the International Training Center on the Methodology of Research on Mackerel (Rastrelliger) FAO Expanded Technical Assistance Program, Rep. No. 1095, 129 pp.
- HOLT, S. J. 1960: A preliminary comparative study of the growth, maturity and mortality of sardine. Proc. World Scient. Meet. Biology Sardines and Related Species, FAO Rome, Vol. II, Subject Syn. (2): 553-561.
- HOLDEN, M. J. 1974: Problems in the rational exploitation of elasmobranch populations and some suggested solutions, p. 117-137 in: Harden Jones, F.R. (ed.) 1974, Sea Fisheries Research. Elek Scientific Books, Ltd., London, 510 pp.
- HUREAU, J. C. 1970: Biologie comparée de quelques Poissons antarctiques (Notothenidae). Bull. Inst. Oceanogr., Monaco, 68 (1391): 244 pp.
- HUREAU, J. C. and C. OZOUF 1977: Détermination de l'âge et croissance du coelacanth Latimera chalumnae Smith, 1939 (Poisson, Crossopterygien, Coelacanthidé). Cybium 3ieme Série 2: 129-137.

- JORDAN, D. S. 1963: The Genera of Fishes and a Classification of Fishes. Stanford University Press (Reprint Edition) 800 pp.
- NIKOLSKY, G. W. 1957: Spezielle Fischkunde, V.E.B. Deutscher Verlag Der Wissenschaften: Berlin, 632 pp.
- RAO, V. S. 1966: Age and growth of "Ghol", Pseudosciaena diacanthus (Lacépède), in Bombay and Saurashtra Waters. Ind. J. Fish. 13 (1-2): 251-291.
- RICHARD, F. C. 1959: A flexible growth function for empirical use. J. Exp. Bot. 10 (29):290-300.
- RICKER, W. E. 1958: Handbook of computation for biological statistics of fish population. Fish.Res.Bd Can.Bull. (119) 300 pp.
- RICKER, W. E. 1973: Linear Regressions in Fishery Research. J. Fish. Res. Bd. Can. 30: 409-434.
- TAYLOR, C. C. 1958: Cod growth and temperature. J. du Conseil, 23: 366-370.
- TAYLOR, C. C. 1962: Growth equation with metabolic parameters. J. du Conseil 27, 270-286.
- THUROW, W. 1974: Fischereiressourcen der Ostsee und ihre Bedeutung für die Kutterfischerei. Arbeiten des dt. Fischereiverbandes (17): 20-51.
- TOMLINSON, P. K. and ABRAMSON 1961: Fitting a von Bertalanffy growth curve by least squares, including table of polynomial. Fish. Bull. Calif. 116: 1-69.
- URSIN, E. 1967: A mathematical model of some aspects of fish growth, respiration, and mortality. J. Fish. Res. Bd. Canada 24 (11): 2355-2453.
- WALFORD, L. A. 1946: A new graphic method of describing the growth of animals. Biol. Bull. 90 (2): 141-147.

Part II

2.1. Alphabetic Index to families

	Family	Genera	Species	Units	Sexed Units	Page
1.	ACANTHURIDAE	1	4	4	0	27
2.	ACIFENSERIDAE	3	9	10	0	27
3.	AGONIDAE	1	1	1	0	27
4.	AMMODYTIDAE	1	3	12	0	28
5.	ANARHICHIDAE	1	3	8	4	28
6.	ANOPLOMATIDAE	1	1	2	2	29
7.	APOGONIDAE	1	2	2	0	29
8.	ARGENTINIDAE	1	3	10	2	29
9.	ARRIPIDAE	1	1	1	0	29
10.	ATHERINIDAE	1	1	2	2	30
11.	BALISTIDAE	2	2	2	0	30
12.	BATRACHOIDAE	2	2	3	2	30
13.	BERYCIDAE	1	1	1	0	30
14.	BLENNIDAE	2	3	3	0	30
15.	BOTHIDAE	4	4	11	7	31
16.	BRANCHIOSTEGIDAE	1	3	11	2	32
17.	CALLIONYMIDAE	1	2	4	4	32
18.	CALLORHYNCHIDAE	1	1	1	0	32
19.	CARANGIDAE	4	10	16	0	33
20.	CARCHARCHINIDAE	3	3	4	2	33
21.	CENTROPOMIDAE	1	2	3	0	34
22.	CHAENICHTHYIDAE	2	2	2	0	34
23.	CHAETODONTIDAE	1	1	1	0	34
24.	CHANIDAE	1	1	1	0	34
25.	CHARACINIDAE	1	1	1	0	34
26.	CHEILODACTYLIDAE	1	1	1	0	34
27.	CHIMAERIDAE	1	1	2	2	35
28.	CHIROCENTRIDAE	1	1	1	0	35
29.	CICHLIDAE	1	10 (incl. 2 hybrids)	39	10	36



Alphabetic Index (cont'd)

Family	Genera	Species	Units	Sexed Units	Page
30. GLUPEIDAE	14	25	185	10	38
31. CORYPHAENIDAE	1	1	2	2	48
32. COTTIDAE	4	6	10	7	48
33. CYCLOPTERIDAE	2	3	3	0	48
34. CYPRINIDAE	18	19	58	2	49
35. CYPRINIDONTIDAE	2	2	18	17	52
36. ELOPIDAE	1	1	1	0	53
37. EMBIOTOCIDAE	2	2	3	2	53
38. ENGRAULIDAE	7	18	90	4	54
39. ESOCIDAE	1	3	11	6	59
40. GADIDAE	13	17	190	19	60
41. GASTEROSTEIDAE	4	4	8	4	70
42. GERRIDAE	2	2	4	0	70
43. GOBIIDAE	4	6	9	4	71
44. HARPADONTIDAE	1	1	2	0	71
45. HOLOCENTRIDAE	1	1	1	0	71
46. ICTALURIDAE	1	1	1	0	72
47. ISTIOPHORIDAE	3	4	10	4	72
48. LABRIDAE	7	18	39	32	73
49. LACTARIIDAE	1	1	1	0	75
50. LAMNIDAE	1	1	1	0	75
51. LATIMERIDAE	1	1	2	2	75
52. LEIOGNATIDAE	1	6	19	0	76
53. LETHRINIDAE	1	2	3	0	77
54. LOPHIIDAE	1	1	2	0	77
55. LUTJANIDAE	1	3	5	0	77
56. MACRORHAMPHOSIDAE	1	1	1	0	78
57. MACROURIDAE	3	4	6	4	78
58. MAENIDAE	1	1	1	0	78
59. MERLUCCIDAE	1	8	32	12	79
60. MUGILIDAE	2	8	24	12	81

Alphabetic Index (contd)

Family	Genera	Species	Units	Sexed Units	Page
61. MYCTOPHIDAE	5	6	7	0	82
62. NEMIPTERIDAE	1	13	23	7	83
63. NOTOTHENIDAE	2	6	17	14	85
64. OSMERIDAE	2	2	16	4	86
65. PANGASIDAE	1	1	1	1	87
66. PERCIDAE	3	7	20	4	88
67. PLATYCEPHALIDAE	1	1	6	2	87
68. PLEURONECTIDAE	14	16	53	28	89
69. PLOTOSIDAE	2	2	3	0	92
70. POLYNEMIDAE	2	3	5	0	92
71. POLYDONTIDAE	1	1	1	0	92
72. POMACENTRIDAE	1	4	4	0	92
73. POMADASYDAE	5	5	10	2	93
74. POMATOMIDAE	1	1	2	0	93
75. PSETTODIDAE	1	1	2	2	93
76. RACHYCENTRIDAE	1	1	2	2	93
77. RAJIDAE	1	6	13	10	94
78. SALMONIDAE	6	18	57	8	95-97
79. SCARIDAE	2	5	5	0	94
80. SCIAENIDAE	17	31	66	17	98
81. SCOMBERESOCIDAE	1	1	3	0	101
82. SCOMBRIDAE	4	6	38	6	102
83. SCORPAENIDAE	9	14	22	7	104
84. SCYLIORHINIDAE	1	1	1	0	105
85. SERRANIDAE	11	14	25	10	105
86. SIGANIDAE	1	6	9	0	107
87. SILLAGINIDAE	1	3	3	0	107
88. SILURIDAE	1	1	1	0	107
89. SPARIDAE	11	20	35	2	108
90. SPHYRAENIDAE	1	2	3	2	110
91. SQUALIDAE	1	1	9	8	110
92. STICHAEIDAE	2	2	3	2	110

Alphabetic Index(cont'd)

Family	Genera	Species	Units	Sexed Units	Page
93. SYNGNATHIDAE	3	4	4	0	111
94. SYNODONTIDAE	1	2	5	0	111
95. TETRAODONTIDAE	1	1	1	0	111
96. THUNNIDAE	6	13	66	4	112
97. TORPEDINIDAE	1	1	1	0	115
98. TRIACANTHIDAE	1	1	1	0	115
99. TRICHIURIDAE	1	1	6	0	116
100. TRYGLIDAE	2	3	5	2	116
101. TRYGONIDAE	2	3	5	4	116
102. XIPHIDAE	1	1	1	0	116
103. ZEIDAE	1	1	3	1	117
104. ZOARCIDAE	1	1	2	0	117

2.2. Data by families: Acanthuridae to Zoarcidae

SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family ACANTHURIDAE							
Acanthurus bahianus	Virgin Islands, Caribbean	23.5	0.314	-		Randall 1962	31
" chirurgus	" "	33.2	0.254	-		" "	31
" coeruleus	" "	31.6	0.247	-		" "	31
" triostegus	Oahu, Hawaii	21.7	0.350	-		" 1961	31
Family ACIPENSERIDAE							
Acipenser dauricus	-	275.	0.073	-		d'Arcona 1937	4
" guldenstaedti	Azov Sea	253.	0.045	-3.5		in: Mikolsky 1957	4
" medirostris	-	190.	0.087	-		d'Arcona 1937	4
" ruthenus	-	57.8	0.130	-		d'Arcona 1937	4
" nudiventris	Southern Caspian & Ural River	205.	0.119	-		in: Mikolsky 1957	4
" "	Aral Lake	205.	0.053	-4.35		" "	4
Huso huso	Azov Sea	249.	0.097	-		" "	4
Pseudoscaphirhynchus kaufmanni	Amu-Dar'ja River	65.4	0.164	-0.95	16° spawn.	" "	4
Acipenser stellatus	--	201.	0.06	-		v. Bertalanffy 1951	0
" fulvescens	Nottaway River, Canada	156.	0.033	-5.59		Magnin 1965	4
Family AGONIDAE							
Agonus cataphractus	Seine Bight, Normandy Coast	15.	0.475	-0.15		Le Gall 1969	4

SPECIES NAME	SAMPLING LOCATION	Lo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family AMMODYTIDAE							
<i>Ammodytes dubius</i>	Nova Scotia Bank, Canada	41.7	0.228	-		Scott 1968	4
" <i>marinus</i>	Haddock Bank, Southern North Sea	21.8	0.89	-	M = 1.24	Macer 1966	00
"	Isle of Man, Irish Sea	12.2	0.98	-		Cameron 1958	4
"	North East Scotland	20.	0.393	-		Im: Reay 1970	41
"	Murman Coast, USSR	24.9	0.163	-		Kirillov 1936	4
"	Western Irish Sea	15.	0.334	-		Molloy 1967	41
" <i>tobianus</i>	Langstone Harbour, Gr. Britain	16.8	0.77	-	M = 1.29	Reay 1973	0
"	Kiel Bight, Western Baltic	21.9	0.449	-	fall spawners	Kändler 1941	4
"	Isle of Man, Irish Sea	22.2	0.633	-	spring "	Cameron 1958	4
"	Isle of Man, Irish Sea	23.4	0.552	-	fall "	Cameron 1958	4
"	Porthmouth, Southern England	16.1	0.720	-	spring "	Reay 1970	4
"	Porthmouth, Southern England	19.7	0.704	-	fall "	Reay 1970	4
Family ANARHICHIDAE							
<i>Anarhichas denticulatus</i>	North Atlantic "1-2° C"	150.	0.098	-		Beese & Kändler 1969	0
"	"	162.	0.081	-2.94		in: Nikolsky 1957	4
" <i>lupus</i> ♂	North Atlantic "2-4° C"	167.	0.044	-		Beese & Kändler 1969	0
" " ♀	North Atlantic "2-4° C"	158.	0.043	-		Beese & Kändler 1969	0
"	"	104.	0.121	-2.69		in: Nikolsky 1957	4
" <i>minor</i>	North Atlantic "1° C"	181.	0.061	-		Beese & Kändler 1969	0
" " ♂	Bear Island, North Norway	140.	0.124	-		Østvedt 1964	4
"	Bear Island, North Norway	153.	0.094	-		Østvedt 1964	4

SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family ANOPLONATIDAE							
Anoploma fimbria ♂	Oregon, USA	78.	0.23	-1.58		Pruter 1954	4
" " ♀	Oregon, USA	96.	0.19	-1.54		Pruter 1954	4
Family APOGONIDAE							
Apogon cyanosoma	Eilat, Red Sea	9.4	2.904	-0.07		Gundermann & Popper 1975	4
" imberbis	Nice to Menton, France, 20 m	15.	0.90	-		Garnaud 1962	41
Family ARGENTINIDAE							
Argentina semifasciata	Japan	19.	1.2	-		Kanayu 1956	00
" silus	44°N 52°W	45.4	0.199	-		Keisler 1966	4
" "	57°N 8°E	50.7	0.120	-		" "	4
" "	63°N 24°W	52.5	0.121	-		" "	41
" "	65°N 30°W	55.6	0.143	-		" "	41
" " ♂	43°N 64°W	42.5	0.166	-		Soviet data in: Emery & McCracken 1966	4
" " ♀	43°N 64°W La Have Bank	39.3	0.179	-		Emery & McCracken 1966	4
" "	43°N 60°W La Have Bank	32.0	0.248	-		Polish data in: Emery & McCracken 1966	4
" "	43°N 62°W max cpue. at 7.5° C	43.8	0.139	-1.74		Emery & McCracken 1966	4
" sphyraena	Mediterranean, off Sète, France	22.3	0.278	-		Lee 1963	4
Family ARRIPIDAE							
Arripis trutta	New South Wales, Australia	62.	0.3	-0.14		Nicolls 1973	0

SPECIES NAME	SAMPLING LOCATION	Loo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family ATHERINIDAE							
Menidia extensa ♂	Lake Waccamaw, North Carolina	9.6	1.36	-	} spawning at 70° F	Davis & Londer 1969	41
" " ♀	Lake Waccamaw, North Carolina	10.4	1.04	-		Davis & Londer 1969	41
Family BALISTIDAE							
Rudarius ercodes	Tomiooka Bay, Japan	6.0	1.20	-		Kikuchi 1966	4
Balistes vetula	Virgin Islands, Caribbean	60.	0.23	-		Randall 1962	31
Family BATRACHOIDAE							
Parichthys porosissimus	Port Aransas, Texas	13.0	1.752	-		Lane 1967	41
Opsanus tau ♂	Patuxent River (Chesapeake Bay)	38.1	0.106	-		Schwarz & Dutcher 1963	4
" " ♀	Patuxent River (Chesapeake Bay)	24.1	0.410	-		Schwarz & Dutcher 1963	4
Family BERYCIDAE							
Beryx splendens	Sagami Bay, Japan	40.8	0.457	-		Ikenouye & Masuzawa 1968	0
Family BLENNIDAE							
Acanthoclinus quadridactylus	Auckland Harbour, N. Zealand	21.5	0.231	-0.48		Jillett 1968	4
Blennius pavo	Castiglione, Algeria	15.	0.5	-		reference lost	4
" pholis	Southern England	17.	0.30	-		Qasim 1957	00

SPECIES NAME	SAMPLING LOCATION	Lo	K	t <sub>o</sub>	NOTES	SOURCE	METHOD
Family BOTHIDAE							
<i>Scophthalmus maximus</i> ♂	North Sea	49.2	0.37	-0.51		Jones 1974	0
" ♀	North Sea	64.8	0.26	-0.05		Jones 1974	0
"	North Sea (7)	106.	0.067	-		Kyle 1926	4
" ♂	Southern North Sea	52.5	0.320	-0.10		Mengl 1963	4
" ♀	Southern North Sea	70.	0.148	-1.79		Mengl 1963	4
<i>Lepidorhombus megastoma</i> ♂	North of Gulf of Biscay 200 m	59.4	0.130	-		FurnestIn 1935	4
" ♀	North of Gulf of Biscay 200 m	97.5	0.076	-		FurnestIn 1935	4
"	East of Shetlands, North Sea 200m 141.		0.047	-0.01		FurnestIn 1935	4
"	North Scottish Waters, N:Coast	61.1	0.13	-0.94		Robb 1963	5
<i>Pseudorhombus cinnamomeus</i>	-	38.4	0.311	-		Matsuura 1961	0
<i>Citharichthys sordidus</i> ♂	Californian waters	30.0	0.3	-	M<0.3	Arora 1951	00



log k = 8' - 2 log L<sub>∞</sub>

SPECIES NAME	SAMPLING LOCATION	L <sub>∞</sub>	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family BRANCHIOSTEGIDAE							
Branchiostegus japonicus	Pacific Coast, Central Japan	42.6	0.286	-0.72		Yasuda & Kosaka 1951	4
"	East China Sea 30°N 126°E 100 m	38.4	0.304	-0.38		Hayashi 1976	0
"	East China Sea, "	34.2	0.300	-		Hayashi 1976	0
"	-	57.4	0.176	0.75		Yasuda & Kosaka 1951	4
"	Japan Sea (Awajima)	41.6	0.358	-		"	4
"	East China Sea	43.5	0.331	-0.48		"	41
"	East China Sea	39.4	0.312	-0.58		"	4
"	argentatus	48.	0.164	-		"	4
"	auratus	42.6	0.286	-0.72		"	4
"	Pacific Coast of Central Japan	39.3	0.382	-		"	41
"	Japan Sea (Awajima)	43.5	0.331	-0.48		"	41
"	East China Sea	>40.0					
Family CALLIONYMIDAE							
Callionymus belcheri ♂	Moreton Bay, Australia 20 m	11.	0.477	-		Johnson 1972	4
"	Moreton Bay, Australia 20 m	10.5	0.411	-		Johnson 1972	4
"	lyra ♂	25.	0.43	-		Hsiao-Wei, Chang 1951	00
"	"	17.5	0.55	-		Hsiao-Wei, Chang 1951	00
Family CALLORHYNCHIDAE							
Callorhynchus milli	New Zealand 44°S 173°E 50 m	150.	0.26	-		Gorman 1963	5

1  
3  
2  
1

SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family CARANGIDAE							
Decapterus macrosoma	Palawan Waters, Philippines	24.4	0.949	-		Tiews <u>et al.</u> 1970	5
" punctatus	Al Ghardaqa, Red Sea	32.3	0.318	-0.56		Rafail 1972 a	0
" russelli	Corregidor Islands, Philippines	23.3	1.132	-		Tiews <u>et al.</u> 1970	5
Trachurus japonicus	Amakusa, Japan	64.7	0.138	-		Mukarani & Shindo 1949	4
" "	East China Sea 33°N 127°E	42.5	0.732	-0.07		Mitani & Ida 1964	0
" "	-	65.6	0.242	-0.65		Yamada & Kazihara 1955	4
" symmetricus	San Diego, California	88.7	0.207	-		Fitch 1956	41
" trachurus	Western South Africa 33°S 18°E	54.3	0.127	-0.31		Geldenhuys 1973	0
" "	North Sea , Coastal Waters	43.4	0.18	-0.48		Sarhage 1970	4
" "	Portuguese Coast	40.	0.158	-		Barraca 1954	41
" "	Black Sea	22.4	0.299	-		Mikolsky 1957	4
" murphyi	Valparaiso, Chile 33°S	75.7	0.62	-		Kaiser 1973	0
Seriola dorsalis	Southern California 33°N 118°W	148.	0.136	-1.9		Baxter <u>et al.</u> 1960	0
" quinqueradiata	Wakasa Bay, Japan	116.	0.327	-0.05		Mitani & Sato 1959	4
" "	Rensen, Korea	110.	0.556	-0.11	LT?	Honda 1950	4
Selaroides leptolepis	Inner Gulf of Thailand	20.	1.155	-		Morsuwan 1970	51
Family CARCHARHINIDAE							
Mustelus canis ♂	Buzzard Bay, 41°30'N 70°46'W	110.	0.07	-4.8		Moss 1972	0
" " ♀	Buzzard Bay "	150.	0.041	-5.4		Moss 1972	0
Prionace glauca	Southwestern England	423.	0.11	-1.04		Stevens 1975	0
Galeorhinus australis	Bass Strait (Australia-Tasmania)	168.	0.130	-1.24		Olsen 1954	4

SPECIES NAME	SAMPLING LOCATION	Loo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family CENTROPOMIDAE							
Lates calcarifer	culture ponds, Songkhla, Thailand	37.8	1.82	-		Wongsomnuk et al. 1971	3
" niloticus	Nozha Hydrodrome, Alexandria	180.	0.177	-		Hashem & Hussein 1973	1
" calcarifer	Daru, Papua New Guinea	113.	0.187	-		Reynolds & Moore 1973	2
Family CHAENICHTHYIDAE							
Pseudochaenichthys georgianus	South Georgia 20 m	54.4	0.316	-		Olsen 1955	4
Champsocoephalus gunnari	" " 20 m	41.4	0.394	-		Olsen 1955	2
Family CHAETOPONTIDAE							
Pomacanthus aureus	Virgin Islands, Caribbean	47.4	0.214	-	K too low?	Randall 1962	31
Family CHANIDAE							
Chanos chanos	culture pond, India	31.5	2.31	-		Tampi 1960	00
Family CHARACINIDAE							
Hydrocyon forskalii	inland waters, Sudan	55.4	0.452	-		Rafail et al. 1973	2
Family CHEILODACTYLIDAE							
Cheilodactylus macropterus	New Zealand	56.3	0.10	-4.3	M = 0.15 (but t <sub>0</sub> too high and K too low)	Vooren 1972	0

35

SPECIES NAME	SAMPLING LOCATION	Lo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family CHIMAERIDAE							
Hydrolagus collei ♂	Cape Blanco - Newport, Oregon USA	96	0.221	-		Johnson & Horton 1972	41
" "	" "	79	0.196	-		" "	41
Family CHIROCENTRIDAE							
Chirocentrus nudus	India 9°N 79°20'E	87.2	0.242	-		Luther 1973	41

SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>o</sub>	NOTES	SOURCE	METHOD
Family CICHLIDAE							
Tilapia esculenta	Lake Victoria, Kavirondo Gulf	32.0	0.50	-		Garrod 1959	0
"	" , outside Kav.Gulf	33.5	0.31	-		"	0
"	" , Mwanza Area	32.4	0.31	-		"	0
"	" , Jinja Area	33.8	0.32	-		"	0
	all: temp. 22,5° C						
	the following data stem from aquaria experiments, food <u>ad libitum</u>						
Tilapia esculenta	oligochaete worms	12.5	2.22	-	Table I	Cridland 1960	2
"	beef and mutton liver	10.5	2.03	-	" II	"	2
"	beef muscle	9.5	1.02	-	" III	"	2
"	Mormyrus stomach contents	25.1	0.427	-	" V	"	2
"	Tilapia muscle	12.2	0.908	-	" VI	"	2
"	Mormyrus muscle	9.0	1.89	-	" VII	"	2
"	Tilapia stomach contents	30.8	0.255	-	" VIII	"	2
"	prawns	16.3	0.794	-	" X	"	2
"	Euglena	8.0	1.02	-	" XII	"	2
"	maize meal	6.4	0.958	-	" XIV	"	2
"	small aquarium, Bemax, prawns & worms	12.8	3.29	-	" XVII	"	2
"	large aquar. Bemax, prawns & worms	10.1	3.80	-	" XVIII	"	2
"	small aquarium, Bemax, prawns & worms	12.5	2.22	-	" XIX	"	2
"	Bemax, prawns, worms & Daphnia	12.4	2.87	-	" XXI	"	2
Tilapia variabilis	Bemax, prawns, worms & Daphnia	8.0	4.26	-	" XXII	"	2

SPECIES NAME	SAMPLING LOCATION	L <sub>oo</sub>	K	t <sub>o</sub>	NOTES	SOURCE	METHOD
Family CICHLIDAE							
Tilapia mossambica ♂	Lake Sibaya, 27°25' S 32°40' E	27.5	0.350	-0.38		Bruton & Allanson 1974	4
" "	" "	21.5	0.425	-0.54		"	4
" "	Njelele Dam 22°44' S 30°7' E	30.0	0.521	-		Le Roux 1961	2
" "	Fundudzi Lake 22°50' S 30°20' E	33.0	0.148	-		"	2
" "	Loskop Dam 25°25' S 29°21' E	47.0	0.212	-		"	2
" "	Sheya-Io-ngubu Dam 25°43' S 31°21' E	34.8	0.373	-		"	2
" "	Rust-der-Winter Dam 25°14' S 28°28' E	41.7	0.226	-		"	2
" "	Harteespoort Dam 25°44' S 27°50' E	34.2	0.363	-		"	2
" "	experimental ponds, Thailand	12.2	7.03	-		Chotiarnwong 1971	2
T. mossambica ♀ x T. nilotica ♂♂,	" "	15.4	8.14	-		"	2
T. mossambica ♂♂ x T. nilotica ♀♀	" "	11.8	7.54	-		"	2
Tilapia nilotica	" "	10.6	8.14	-		"	2
" "	Lake Tiberias, Israel	35.7	0.501	-		Ben-Tuvia 1960	4
" "	" "	33.3	0.538	-		"	4
T. nilotica ♀♀ x T. macrochir ♂♂	experimental ponds, Bonaké, Ivory coast		1.491	-		Lessent 1968	3
Tilapia nigra ♂	experim. ponds, Sagana, Kenya	27.5	0.203	-		van Someren & Whitehead 1960	2
" zillii	Lake Tiberias, Israel	26.3	0.234	-		Ben-Tuvia 1960	4
" melanotheron	Lagos lagoon, Nigeria	33.1	0.159	-		Fagade 1974	4
" galilaea ♂	Lake Tiberias, Israel	35.7	0.501	-		Ben-Tuvia 1960	4
" "	" "	33.3	0.538	-		"	4

SPECIES NAME	SAMPLING LOCATION		Lo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family CLUPEIDAE								
Clupea harengus	East Anglia, YC	1932	28.2	0.378	-		Burd 1960	4
"	"	1933	28.4	0.361	-		"	4
"	"	1934	29.4	0.234	-		"	4
"	"	1935	29.3	0.264	-		"	4
"	"	1936	29.1	0.307	-		"	4
"	"	1943	29.5	0.290	-		"	4
"	"	1944	30.3	0.222	-		"	4
"	"	1945	30.2	0.267	-		"	4
"	"	1946	31.0	0.239	-		"	4
"	"	1947	29.9	0.375	-		"	4
"	"	1948	30.4	0.313	-		"	4
"	"	1949	31.0	0.296	-		"	4
"	"	1950	31.6	0.264	-		"	4
"	Buchan Herring YC	1952	31.1	0.28	-2.87		Hubold 1975	0
"	"	1953	30.6	0.42	-0.28		"	0
"	"	1954	30.7	0.30	-1.84		"	0
"	"	1955	32.1	0.31	-2.24		"	0
"	"	1956	35.1	0.16	-4.87		"	0
"	"	1957	32.9	0.28	-1.60		"	0
"	"	1958	33.3	0.29	-1.60		"	0

SPECIES NAME	SAMPLING LOCATION		Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family CLUPEIDAE								
Clupea harengus	Buchan Herring	YC 1959	32.9	0.32	-1.30		Hubold 1975	0
"	"	YC 1960	32.7	0.42	-0.81		"	0
"	"	YC 1961	32.7	0.48	-0.22		"	0
"	"	YC 1962	31.5	0.43	-0.84		"	0
"	Baltic, Gulf of Riga		17.	0.666	-		Nikolsky 1957	4
"	Baltic, Gulf of Gdansk	1936-39	23.9	0.295	-1.67	fall spawners	Popiel 1958	0
"	"	1946-57	24.4	0.35	-2.3	"	"	0
"	"	1936-37	24.7	0.273	-1.01	spring	"	0
"	"	1946-57	24.5	0.244	-1.68	"	"	0
"	"	Bornholm Deep	22.1	0.493	-		"	2
"	"	1943-45	23.6	0.381	-		"	2
"	"	1952-57	26.0	0.434	-1.43		"	0
"	"	Kiel Bight	27.9	0.620	-	spring spawn.	Weber 1970	0
"	"	"	28.7	0.630	-	fall	"	0
"	"large White Sea herring"		25.3	0.381	-	spawn.t.5-15° C	Nikolsky 1957	4
"	"lesser White Sea herring"		21.	0.296	-	" 0-6° C	"	41
"	Murman Herring (72°N 35°W)		28.3	0.292	0.289		"	4
"	"North Sea"		30.	0.38	-	M = 0.25	Beverton & Holt 1959	0

1  
3  
1



SPECIES NAME		SAMPLING LOCATION		Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family CLUPEIDAE									
Sardina pilchardus		Sète (French Mediterranean Coast)		19.9	0.360	-1.09		Lee 1961	4
"	"	Morocco		21.6	0.49	-	M = 0.50	Belveze 1972	4
"	"	Alicante (Altea), Spain		20.1	0.405	-1.86		Rodriguez-Roda & Larraneta 1955	4
"	"	Alicante (Torrevic.), Spain		20.9	0.529	-1.11		"	4
"	"	Agean (Mytilini)		16.7	0.483	-1.31		Laskaridis 1948	4
"	"	Agean (Oreos),		16.6	0.399	-1.64		Laskaridis 1948	4
"	"	Upper Adriatic (off Venedice)		17.0	0.689	-1.02		Mozzi & Duo 1958	4
"	"	Castellon, Spain		19.6	0.31	-		Larrañeta & Lopez 1958	4
"	"	Formentera, Spain		22.5	0.33	-1.81		Andreu <u>et al.</u> 1950	4
"	"	Gulf of Lion		16.4	0.557	-0.66		Fage 1920	4
"	"	Castellon, Spain		16.6	0.565	-1.03		Andreu <u>et al.</u> 1950	4
"	"	Spanish Medit. Coast		17.	0.60	-		Bougis 1952	00
"	♂	Loire to Gironde Mouths, Biscay		20.	0.32	-		d'Ancona 1937	00
"	♀	Loire to Gironde Mouths, Biscay		25.	0.25	-		d'Ancona 1937	00
"	"	off Split, Yugoslavia		17.	0.40	-		Muzinič 1957	00
"	"	Castellon, Spain		20.3	0.306	-1.54		Larrañeta 1965	0
"	"	Barcelona, Spain		20.4	0.272	-2.21		Lopez 1963	4
"	"	Bay of Biscay		19.	0.40	-		Bougis 1952	00

SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family CLUPEIDAE							
<i>Sardinella albella</i>	Gulf of Manaar, India	13.0	1.654	-		Bennett 1961	5
"	Gulf of Manaar, India	13.3	1.298	-		Benett 1961	5
"	Gulf of Manaar (?)	17.	1.1	-		Nair 1960	00
"	Mandapam, India	13.3	1.44	-		Sekaran 1955	0
" anchovia	Venezuela, Gulf of Coriaco 64°W	24.2	0.725	-		Heald & Griffiths 1967	0
"	Venezuela	24.0	0.7	-		Simpson 1971	0
" eba	Senegal	35.	0.608	-		Postel 1955	4
"	Congo	39.6	0.277	-		Rossignol 1955	4
"	Congo	24.9	0.990	-		Chéno & Le Guen 1968	0
" fimbriata	Karwar, India	18.4	1.316	-		Radhakrishnan 1964	4
"	Philippines	14.	1.6	-		Ronquillo 1960	00
" gibbosa	Mandapam, India	17.1	1.08	-		Sekaran 1955	00
" longiceps	Bali Strait, Indonesia	23.8	0.505	-		Dwiponggo 1972	0
"	Indian Waters	21.0	0.60	-1.12		Anthony Raja 1970	0
"	India, West Coast	20.7	0.528	-1.33		Banerji 1973	0
"	India	27.	0.4	-		Nair 1960	00
"	Malabar Coast, India	20.7	0.50	-		Hornell & Naidu 1924	00
" sirm	Thousand Islands, Java Sea	24.3	0.586	-		Barhanuddin et al 1974	5
"	Al Ghardaqa, Red Sea	22.6	0.65	-		Rafail 1972 a	41

SPECIES NAME	SAMPLING LOCATION	LOO	K	t <sub>o</sub>	NOTES	SOURCE	METHOD
Family CLUPEIDAE							
<i>Sardinella aurita</i>	Pointe Noire, Congo, January Brood	26.0	1.023	-0.09		Ghéno 1975	0
"	Pointe Noire, Congo, May Brood	26.0	1.208	-0.03		Ghéno 1975	0
"	West Africa 19°N 17°W	40.7	0.326	-		Pham-Thuoc & Szypula 1975	0
"	Canary Islands	37.0	0.257	-0.82		Navarro 1932	4
"	Israel Coast	31.1	0.250	-1.81		Ben-Tuvia 1956	4
"	Senegal	51.9	0.170	-1.05		Postel 1955	4
"	Balearic Islands	27.7	0.447	-1.41		Navarro 1932	4
"	Pointe Noire, Congo	31.0	0.693	-		Rossignol 1955	4
"	Rio de Janeiro, Brazil	24.5	0.423	-1.23		Richardson et al 1960	4
"	Agean	29.0	0.50	-		Anamitades 1952	00
"	Balearic Islands	30.	0.35	-		d'Ancona 1937	00
"	Balearic Islands	35.	0.25	-		d'Ancona 1937	00
"	Castiglione Bight, Algeria	25.	0.534	-		Dieuzelde & Roland 1957	1
<i>Sardinops melanostica</i>	Japan	27.0	0.65	-		Tokai Lab. 1960	0
"	N.West Pacific (Japan, recent)	27.0	0.90	-		Nakai 1960	00
"	N.West Pacific (Japan, early years)	29.0	0.50	-		Holt 1960	0
"	Walvis Bay, Namibia	36.0	0.358	-0.235		Matthew 1960	4
"	Walvis Bay, Namibia 23° S 14° E	30.6	0.225	-		Newman 1970	0
"	33° S 17° E, South Africa	29.5	0.45	-		Davies 1958	00

SPECIES NAME	SAMPLING LOCATION	Loo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family CLUPEIDAE							
Sardinops caerulea	California	30.5	0.396	-0.59	M = 0.15	Phillips 1948	4
"	"	29.	0.60	-		"	00
"	"	31.	0.46	-		"	00
"	San Francisco, Calif.	29.5	0.398	-2.10	M = 0.45	in: Marr 1960	4
"	California	29.3	0.45	-	M = 0.4	Beverton 1963	0
"	Mexico (Baja California ?)	26.	0.50	-		Granados 1958	0
"	British Columbia, Year class 37	25.1	0.57	-		Felin 1954	00
"	"	29.1	0.55	-		"	00
"	"	29.3	0.52	-		"	00
"	"	30.0	0.52	-		"	00
"	"	30.2	0.53	-		"	4
"	Monterey, California	30.0	0.35	-		in: Marr 1960	4
"	San Pedro, Calif. YC 1937	26.0	0.54	-		Felin 1954	00
"	"	26.1	0.52	-		"	00
"	"	27.6	0.53	-		"	00
"	"	26.9	0.56	-		"	00
"	"	25.7	0.55	-		"	00
"	"	26.8	0.59	-		"	00
"	California	27.	0.70	-	M = 0.15	Silliman 1943	00

SPECIES NAME	SAMPLING LOCATION		LOO	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family CLUPEIDAE								
Brevoortia tyrannus	Chesapeake Bay	Year Class 1954	24.3	0.639	-		Henry 1971	4
"	"	" 1955	28.3	0.338	-		"	4
"	"	" 1957	34.1	0.218	-		"	4
"	"	" 1959	36.8	0.238	-		"	4
"	"	" 1960	33.8	0.432	-		"	4
"	"	" 1961	35.1	0.434	-		"	4
"	"	" 1962	30.9	0.511	-		"	4
"	"	" 1963	31.7	0.533	-		"	4
"	North Carolina Fall Fishery	YC 1952	41.8	0.218	-		"	4
"	"	" 1953	36.4	0.561	-		"	4
"	"	" 1954	36.8	0.576	-		"	4
"	"	" 1955	36.6	0.545	-		"	4
"	"	" 1956	36.9	0.442	-		"	4
"	"	" 1958	36.6	0.480	-		"	4
"	"	" 1961	38.8	0.409	-		"	4
"	"	" 1962	36.9	0.455	-		"	4
"	"	" 1963	40.2	0.307	-		"	4
"	"	" 1964	38.4	0.388	-		"	4
"	"	" 1965	37.5	0.441	-		"	4
"	"	" 1951	37.1	0.353	-		"	4

SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family CLUPEIDAE							
Brevoortia tyrannus	South Atlantic YC 1957	25.0	0.490	-		Henry 1971	4
"	" " 1959	22.7	0.463	-		"	4
"	" " 1960	20.6	0.741	-		"	4
"	" " 1962	22.7	0.463	-		"	4
"	" " 1963	21.9	0.770	-		"	4
"	" " 1964	20.9	1.091	-		"	4
"	Middle " " 1951	35.1	0.37	-		"	4
"	" " " 1956	40.5	0.26	-		"	4
"	" " " 1958	44.0	0.209	-		"	4
"	" " " 1959	35.8	0.370	-		"	4
"	" " " 1963	42.8	0.282	-		"	4
"	" " " 1964	39.6	0.343	-		"	4
"	North " " 1951	41.0	0.168	-		"	4
"	" " " 1952	45.8	0.119	-		"	4
"	" " " 1953	37.6	0.290	-		"	4
"	" " " 1954	37.7	0.242	-		"	4
"	" " " 1955	35.1	0.549	-		"	4
"	" " " 1956	39.9	0.231	-		"	4
"	" " " 1962	35.7	0.586	-		"	4
"	" Catch 1955-1956	39.	0.542	-		Reintjes 1969	4

SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family CLUPEIDAE							
Brevoortia tyrannus	Beaufort, North Carolina	47.6	0.289	-		Rush 1952	4
"	North Carolina 1959-1962	35.5	0.475	-		Reintjes 1969	0
"	North Carolina 1959-1962	36.9	0.478	-		Reintjes 1969	0
Sprattus sprattus	Norwegian Coast	16.	0.65	-		Sund 1911	00
"	Western England	16.4	0.53	-	early spawners	Iles & Johnson	0
"	Western England	14.0	0.63	-	late spawners	Iles & Johnson	0
"	Kiel Bight	14.9	1.02	-0.04		Hohendorf 1966	0
"	Black Sea, Soviet Coast	18.0	0.136	-2.5		Aslanova 1954	4
"	Black Sea, Soviet Coast	11.9	0.31	-0.83		Berg et al. 1949	4
"	Southern France	14.2	0.368	-2.30		Furneston 1948	4
"	"North Sea"	13.0	0.70	-	M < 1.2	Robertson 1938	4
"	Northern Adriatic	14.4	0.452	-		Zavodnik 1969	4
"	Gulf of Biscay 0-50 m	17.5	0.297	-		Porche 1976	2
Clupea pallasii	Southern Alaska 50°-60° N	32.	0.349	-		Rounsefell 1930	4
"	-	35.6	0.558	-		Nikolsky 1957	4
"	Canadian West Coast	23.	0.29	-	M = 0.56	Ricker 1958	00
Clupeonella delicatula	Azov Sea	10.9	0.246	-		Berg et al. 1949	2
"	Caspian, Volga-Caspian Rayon	16.1	0.236	-		"	2
"	Caspian, Dagestan Coast	11.0	1.134	-		"	2
Caspialosa saposhnikovi	Caspian, Azerbaydzhan	28.8	0.353	-		"	2

SPECIES NAME	SAMPLING LOCATION	Lo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family CLUPEIDAE							
Hilsa ilisha	Mandapam area, India	51.1	0.49	-		Sekaran 1955	00
" " ♂	Hoogly River, India	44.7	0.652	-		Pillay 1958	4
" " ♀	Hoogly River, India	46.1	1.030	-		Pillay 1958	4
Sardinops neopilchardus	Australia, New South Wales	22.4	0.467	-0.09		Blackburn 1949	4
" "	Australia, Victoria	20.5	0.22	-		Blackburn 1950	00
" "	New Zealand 41° S 173° E	21.9	0.201	-		Baker 1972	2
Anadontostoma chacunda	Godavary Estuary, India	20.8	0.869	-		Babu Rao 1965	5
Clupanodon pseudohispanicus	Venezuela, Gulf of Cariaco	28.5	0.864	-		Martin & Gonzales 1960	5
" "	Venezuela, Gulf of Cariaco	27.6	0.678	-		Martin & Gonzales 1960	5
Etrumeus microps	Amakusa, Japan	21.	1.379	-		Makami & Shindo 1949	4
Dorosoma cepedianum	ponds, Indiana, USA	43.6	0.179	-		Miller 1960	4
Raonda russelina	Indian waters	24.7	0.40	-		Verghese 1961a	00



SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family CORYPHAENIDAE							
Coryphaena hippurus ♂	Florida Strait	167.	0.53	-	} very tentative	Beardsley 1967	4
" " ♀	" "	135.	0.62	-		"	41
Family COTTIDAE							
Cottus beldingii	Lake Tahoe, California	9.	0.316	-		Ebert & Summerfeldt 1969	4
Taurulus bubalis ♂	Kiel Bight, Baltic	18.4	0.207	-		Lamp 1965	4
" " ♀	" "	18.7	0.251	-		"	4
Cottus gobio ♂	Windermere, Gr. Britain	7.2	0.7	-	M = 1.1	Smyly 1957	00
" " ♀	" "	7.3	0.4	-	M = 0.9	"	00
Oligocottus maculosus	Port Renfrew, Brit. Columbia	10.2	0.208	-1.35		Chadwick 1976	4
Myoxocephalus scorpius ♂	Kiel Bight, Baltic	19.6	0.758	-		Lamp 1965	4
" " ♀	" "	32.3	0.358	-		"	4
" " ♀	Heligoland, North Sea	26.7	0.539	-0.572		Hass 1937	4
" octodecimspinosus	New England	30.	0.72	-		Morrow 1951	2
Family CYCLOPTERIDAE							
Liparis inquilinus	39°N, near Hudson Canyon 40 m	6.6	2.78	-		Able & Musik 1976	4
" pulchellus	Northern California, Humboldt Cty. 20. -		0.3	-		Johnson 1970	41
Cyclopterus lumpus	Islandic Waters	55.	0.12	-0.12	very tentative	Saemundsson 1926	1

SPECIES NAME		SAMPLING LOCATION	L <sub>50</sub>	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family CYPRINIDAE								
Abramis brama								
"	"	Aral Lake, USSR	51.9	0.153	-		Nikolsky 1957	4
"	"	"	54.1	0.241	-		Berg <u>et al</u> 1949	2
"	"	Pskov Reservoir, USSR	72.9	0.119	-		"	2
"	"	Ural River Delta, USSR	52.4	0.429	-		in: Backiel & Zawisza 1968	2
"	"	Rybinsk Reservoir, USSR	86.1	0.068	-		"	2
"	"	Volgograd Reservoir, USSR	111.	0.077	-		"	2
"	"	Lake Ladoga, USSR	90.	0.076	-		"	2
"	"	Njemen River, USSR	109.	0.096	-		"	2
"	"	Danube Delta	74.4	0.165	-		"	2
"	"	Middle Course of Dnepr River	65.5	0.191	-		"	2
"	"	Dnepr River Delta, USSR	73.	0.191	-		"	2
"	"	Ilmen Lake, USSR	86.4	0.093	-		"	2
"	"	Volga, near Kuybyshev	93.1	0.081	-		"	2
"	"	Bodensee, FRG	84.2	0.110	-		"	2
"	"	Müggelsee (Berlin)	61.8	0.065	-		"	2
"	"	Lake Hjälmären, Sweden	77.2	0.058	-		"	2
"	"	Lake Onkamo, Finland	105.	0.064	-		"	2
"	"	Danube River, near Medvedovo	71.6	0.109	-		"	2
"	"	Lake Znin Duzy, Poland	91.4	0.085	-		"	2
"	"	North German Lakes	94.5	0.053	-		"	2

SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family CYPRINIDAE							
Abramis brama	Lake Havgardsjön, Sweden	68.6	0.131	-		in: Backiel & Zawisza 1968	2
"	Caspian, brackish waters	54.6	0.257	-		"	2
"	Hommanás-Pellige, Finland, brackish	70.1	0.058	-		"	2
"	Vistula Firth, Baltic, brackish	71.6	0.075	-		"	2
"	Arkona, Baltic, brackish	74.5	0.091	-		"	2
"	Azov Sea, brackish	57.7	0.287	-		"	2
"	-		0.235	-	page 281	v. Bertalanffy 1951	0
Catla catla	River Yamuna, India	128.	0.28	0.11		Natarajan & Jhingran 1963	0
Cirrhinus mrigla	River Godavary, India	140.	0.122	-0.46		Hanumantharo 1974	0
Labeo niloticus ♂	Nozha Hydrodrom, Alexandria	55.8	0.662	-		Hashem 1972	2
"	"	69.2	0.400	-		"	2
Rutilus rutilus	Petschora, USSR	42.5	0.08	-	stationary	Nikolsky 1957	2
"	Ilmen Lake	26.2	0.180	-	form	"	2
"	Aral Lake	51.3	0.101	-	migratory	"	2
"	Don River (Taran)	35.5	0.173	-	form	"	2
Tor tor	Narbadi River, India	109.	0.082	-		in: Jhingran 1975	2
Acanthobrama terra-sanctae	Lake Tiberias, Israel	21.5	0.277	-		Steinitz 1959	1
Brachydanio rerio	Laboratory	5.2	3.97	-		Eaton & Farley 1974	4

SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family CYPRINIDAE							
Parabramis pekinensis	Amur River	60.	0.096	-	All USSR	Nikolsky 1957	1
Erythroculter erythropterus	" "	80.	0.117	-		"	1
Hypophthalmichthys molitrix	" "	70.	0.242	-		"	2
Ctenopharyngodon idella	" "	167.	0.049	-		"	2
Mylopharyngodon piceus	" "	151.	0.085	-		"	2
Barbus brachycephalus	Syr-Darja River, at Tschardara	25.6	0.682	-		"	2
" "	" , at Arys	73.6	0.122	-		"	2
" "	" , at Kasalinsk	143.	0.080	-		"	2
" "	" , at Kostam	110.	0.118	-		"	2
Varicorhinus capoeta	Murgab River, "Taschkepu"	61.1	0.092	-		"	2
" "	" , "Sultanbent"	67.9	0.087	-		"	2
" "	Amur Darja River	44.9	0.170	-		"	2
Carassius auratus	Amur River	28.9	0.284	-		"	2
Carassius carassius	"culture ponds"	22.8	0.358	-		"	2
" "	Petschora River	29.8	0.093	-		"	2
Tinca tinca	Lower Wolga River	29.4	0.710	-		"	2
Cyprinus carpio	Aral Lake	74.8	0.157	-		"	2
" "	Asow Sea	57.3	0.335	-		"	2
" "	Ili River	32.0	0.286	-		"	2
" "	Southern Caspian	65.1	0.396	-		"	2

SPECIES NAME	SAMPLING LOCATION	Loo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family CYPRINIDONTIDAE							
Lebistes reticulatus ♀	Laboratory	5.0	1.74	-0.12	fish no 1	Ursin 1967	4
" ♂	"	2.4	6.90	-0.07	" 2	"	4
" unsexed	"	3.0	5.11	-0.06	" 3	"	4
" ♀	"	5.0	2.12	-0.11	" 4	"	4
" ♂	"	2.3	6.87	-0.07	" 5	"	4
" ♂	"	6.1	1.10	-0.14	" 6	"	4
" ♀	"	4.4	1.98	-0.12	" 7	"	4
" ♂	"	2.6	6.74	-0.06	" 8	"	4
" ♀	"	4.9	1.77	-0.10	" 9	"	4
" ♀	"	5.0	1.56	-0.13	" 10	"	4
" ♂	"	2.2	5.46	-0.08	" 11	"	4
" ♀	"	4.2	2.3	-0.10	" 12	"	4
" ♂	"	2.7	12.3	-		v. Bertalanffy 1951	0
" ♀	"	4.4	4.44	-		"	0
Gambusia affinis ♂	Portugal	3.6	1.20	-	M < 1.6	Da Franca 1953	00
" ♀	Portugal	6.2	0.8	-	N < 0.8	"	00
Lebistes reticulatus ♂	Aquarium 25 ° C	1.5	9.12	-		Svårdson 1943	2
" ♀	Aquarium 25 ° C	3.5	3.23	-		"	2

SPECIES NAME	SAMPLING LOCATION	L <sub>oo</sub>	K	t <sub>o</sub>	NOTES	SOURCE	METHOD
Family ELOPIDAE							
Elops saurus	Cuba, inshore waters	47.	0.24	-0.48		Carles 1967	4
Family EMBIOTOCIDAE							
Amphigonopterus aurora	California, tide pools	15.5	0.363	-0.89		Hubbs 1921.	4
Amphistichus argenteus ♂	California 33.5° N 10-20 m	42.	0.184	-		Carlisle et al. 1960	4
" "	" "	53.	0.147	-		"	4

SPECIES NAME		SAMPLING LOCATION	Loo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family ENGRAULIDAE								
Engraulis japonicus		Japan	15.	1.6	-		Beverton 1963	0
"	"	Sagami Bay, Japan	18.7	0.824	-0.17	spring spawn.	Kondo 1974	4
"	"	"	20.6	0.586	-0.27	fall	"	4
"	"	Wakasa Bay, Japan	18.2	1.76	-		Wanatabe 1958	00
"	"	Northern Kyushu	13.2	4.59	-		"	00
"	"	Suo Nada, Japan	28.1	0.60	-0.20		"	00
"	"	Sea of Japan	18.8	1.48	-		"	00
"	"	Hyuga Nada, Japan	19.8	1.04	-0.17	spring spawn.	Asami & Hanaoka 1957	00
"	"	"	22.6	1.05	-	fall	"	00
"	"	Japan, Pacific Coast	17.8	1.70	-	spring	Hayashi & Kondo 1957	00
"	"	Hyuga Nada, Japan	23.7	0.98	-0.12		Yokota & Asami 1956	00
"	"	"	20.6	0.94	-0.17		Yokota & Furukawa 1952	00
"	"	Harima Nada, Japan	14.4	2.02	-		Wanatabe 1958	00
"	"	Ise Bay, Japan	10.0	3.24	-		"	00
"	"	Japan, Pacific Coast	18.3	1.01	-	fall spawn.	Hayashi & Kondo 1957	00
Engraulis anchoita		Mar del Plata, Argentina	24.1	0.24	-2.2		Fuster de Plaza, no date	00
"	"	"	22.3	0.30	-1.2		"	00
"	ringens	Peru 5°50' - 37°04' S, 10-20 m	15.6	1.38	-		Sanchez 1966	4
"	"	Chile 20° S 75° W	16.9	1.6	-0.02		Simpson & Buzeta 1967	0
"	"	Peru 14 - 17° S	15.0	1.7	-	M ≈ 1.5	Boerema et al. 1965	0

SPECIES NAME		SAMPLING LOCATION	Loo	K	t <sub>o</sub>	NOTES	SOURCE	METHOD
Family ENGRAULIDAE								
Engraulis encrasicolus		Mediterranean, Algero-Maroccan Border	20.	0.396	-		Arrignon 1966	1
"	"	Black Sea, Western	15.	1.26	-0.23		Majarova & Chugunova 1954	00
"	"	Black Sea, Soviet Waters	13.9	1.73	-		Berg et al 1949	00
"	"	Black Sea, Eastern	14.2	0.81	-0.54		Majarova & Chugunova 1954	00
"	"	Black Sea, Eastern	14.1	1.01	-0.28		Svetovidov 1964	00
"	"	Azov Sea	11.7	1.16	-0.17		Berg et al 1949	00
"	"	Bay of Naples	17.6	1.28	-		Page 1920	00
"	"	Channel & Zuidersee	21.5	0.88	-		Hošk 1912	00
"	"	Black Sea, Bulgaria	16.6	0.581	-0.92		Stoyanov 1961	4
"	"	Greek Waters	16.5	1.39	-0.02		Vidalis 1949	00
"	"	Upper Adriatic	15.3	2.44	-		Padoan 1963	00
"	"	Gulf of Lion, France	20.0	0.42	-1.15		Lee & Juge 1965	00
"	"	Azov Sea	9.4	1.21	-0.37		Svetovidov 1964	00
"	"	Gulf of Biscay, France	23.2	0.21	-2.52		Furnestin 1943	00
"	"	North Sea	18.6	1.21	-0.01		Meek 1916	00
"	"	Azov Sea	10.8	7.66	-		Nikolsky 1957	00
Cetengraulis edentulus ♂		Venezuela 10.64° N 64° W	14.5	2.52	-0.34	} spawning	Simpson & Griffiths 1973	0
"	♀	Venezuela, off Taguapire	16.9	1.72	-0.40		"	0
"	"	"	14.0	2.72	-0.43	} spawning	"	0
"	♂	"	16.9	1.86	-0.38		"	0
"	♀	" , 10.64° N 64° W	16.9	1.86	-0.38	} July-Sept.	"	0



SPECIES NAME	SAMPLING LOCATION	LOO	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family ENGRAULIDAE							
Cetengraulis mysticetus	Central East, Pacific	17.5	1.70	-		Beverton 1963	0
"	Almejas Bay, Gulf of California	19.9	1.23	-		Bayliff 1969	0
"	Guaymas Bay,	17.0	2.58	-		Barett & Howard 1961	00
"	Ahome Point,	17.5	2.42	-		"	00
"	Colombia	17.2	2.09	-		Bayliff 1967	0
"	Gulf of Fonseca, Honduras, 1952	18.5	2.92	-		Barett & Howard 1961	00
"	" , 1951, 1954, 1955	20.5	0.90	-		"	00
"	Montijo Bay, Panama (Pac.)	19.1	2.42	-		Bayliff 1963	0
"	Gulf of Panama 1951-1960	17.9	2.36	-		Bayliff 1966	0
"	" 1961-1963	20.4	1.31	-		"	0
"	Gulf of Guayaquil 3° S 80° W	17.4	1.34	-		Barett & Howard 1961	00
Stolephorus bucaneri	Corregidor Islands, Philippines	11.6	1.226	-		Tiews et al 1970	5
" heterolobus	Corregidor Islands	10.3	1.286	-		"	5
" indicus	Singapore Strait	21.8	0.712	-0.13		Tham Ah Kow 1967	0
" insularis	"	11.9	2.08	-0.08		"	0
" indicus	Corregidor Islands, Philippines	17.2	1.085	-		Tiews et al 1970	5
" pseudoheterolobus	Singapore Strait	10.3	2.08	-0.04		Tham Ah Kow 1967	0
" purpureus	Kaneohe Bay, Hawaii	11.4	1.737	-0.04		Struhsaker & Uchiyama 1976	41
" , Species A	Corregidor Islands, Philippines	10.2	1.236	-		Tiews et al 1970	51

SPECIES NAME		SAMPLING LOCATION	Lo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family ENGRAULIDAE								
Engraulis mordax		San Diego, California	16.6	0.69	-1.8	1955-56	Miller & Wolf 1958	00
"	"	San Diego, California	18.2	0.48	-0.20	1956-57	"	00
"	"	Monterey, California	23.1	0.2	-4.3	1946-51	Clark & Phillips 1952	00
"	"	Southern California	17.6	0.349	-2.43	1967	Messersmith 1969	4
"	"	San Francisco	20.1	0.32	-3.1	1952-53	Miller <u>et al.</u> 1955	00
"	"	Los Angeles, California	17.2	0.83	-2.1	1956-57	Miller & Wolf 1958	00
"	"	Los Angeles, California	18.7	0.41	-3.6	1955-56	"	00
"	"	"	18.3	0.48	-2.6	1954-55	"	00
"	"	"	21.1	0.21	-4.4	1953-54	Miller <u>et al.</u> 1955	00
"	"	"	18.4	0.45	-2.7	1952-53	"	00
"	"	Santa Barbara,	18.6	0.44	-2.9	1954-55	Miller & Wolf 1958	00
"	"	"	20.0	0.26	-4.2	1953-54	Miller <u>et al.</u> 1955	00
"	"	Monterey, California	21.9	0.19	-5.3	1955-56	Miller & Wolf 1958	00
"	"	"	19.8	0.20	-6.4	1954-55	"	00
"	"	"	27.0	0.10	-6.3	1953-54	Miller <u>et al.</u> 1955	00
"	"	"	28.5	0.20	-8.1	1952-53	"	00
"	"	Oceanside,	24.1	0.17	-4.5	1956-57	Miller & Wolf 1958	00
"	"	"	31.0	0.09	-5.4	1955-56	"	00
"	"	San Clemente,	19.1	0.40	-2.6	1956-57	"	00
"	"	Newport,	19.3	0.34	-3.1	1956-57	"	00

SPECIES NAME	SAMPLING LOCATION	Loo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family ENGRAULIDAE							
Engraulis mordax	Newport, California	16.5	0.61	-0.5	1955-56	Miller & Wolf 1958	00
"	San Pedro, "	19.0	0.56	-1.4	1956-57	"	00
"	"	16.4	0.95	-1.4	1955-56	"	00
"	Santa Monica, California	18.9	0.47	-2.1	1956-57	"	00
"	Malibu, "	18.7	0.58	-1.5	1956-57	"	00
"	Santa Barbara, "	20.5	0.33	-2.6	1956-57	"	00
Anchoa naso	Manta, Ecuador	8.6	1.78	-		Bayliff 1967	0
"	"	8.5	1.82	-		Joseph 1963	0
Coila borneensis	-	18.5	0.70	-		Vergheze 1971	00
Setipinna taty	Lower Sunderbans, India	20.5	0.332	-1.17		D.K. De & Datta 1974	41
Thrissocles mystax	Malabar Coast, India	23.6	0.76	-		Venkatamaran 1956	0

SPECIES NAME	SAMPLING LOCATION	Loo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family ESOCIDAE							
Esox lucius ♂	Windermere, England 4 m	75.	0.238	-0.18		in: Johnson 1966	0
" " ♀	Windermere, England 4 m	100.	0.264	-		"	0
" "	Wisconsin, USA	93.3	0.310	-	p. 233	Mikolsky 1957	2
" "	Aral Lake, USSR	80.6	0.204	-	"	"	2
" " ♂	Peipus Lake, USSR	64.4	0.332	-	"	"	2
" " ♀	Peipus Lake, USSR	97.5	0.208	-	"	"	2
" " ♂	Chany Lake, USSR	106.	0.123	-	"	"	2
" " ♀	" "	141.	0.097	-	"	"	2
" masquinongy	Wisconsin, USA	143.	0.150	-	"	"	2
" richerti	Tumen River, Sachalin Isl.	153.	0.089	-	"	"	2
" "	Bolon Lake, Amur, USSR	77.4	0.218	-	"	"	2

SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family GADIDAE							
Gadus morhua	North Sea	132.	0.20	-	M < 0.20	Graham 1934	00
"	Northern Norway	154.	0.069	-		Jensen & Hansen 1931	00
"	Baltic, Bornholm Deep	120.	0.125	-0.33		Kosior 1976	0
"	Southern North Sea 54° N 3° E	111.	0.333	-		Daan 1974	0
"	Baltic, Bornholm Deep	99.5	0.2	-	M = 0.44	Kändler 1944	00
"	Baltic, Bornholm Deep	106.	0.177	-		Tiews 1971	00
"	South Western Iceland	102.	0.234	-0.35		Jonsson 1965	4
"	Baltic, off Gdansk	112.	0.154	-	M = 0.31	Kändler 1944	4
"	Baltic, Gdansk Deep	108.	0.153	-		Kosior 1976	0
"	Calais to Boulogne, France	123.	0.232	-		Lefranc 1970	4
"	Calais to Boulogne, France	122.	0.247	-		Lefranc 1970	4
"	Deep of Gdansk, Baltic	72.8	0.209	-		Rutkiewicz 1957	4
"	Baltic, Bornholm	116.	0.095	-		Stanek 1962	4
"	Labrador	73.3	0.154	-	(200 m)	May 1959	4
"	Barents Sea	129.	0.126	-		Trout 1954	4
"	Grand Bank, Newfoundland	176.	0.070	-		Ancellin 1954	4
"	South West Greenland	92.3	0.218	-		Hansen 1949	4
"	North West Greenland	106.	0.164	-		Hansen 1949	4
"	Iceland	99.9	0.209	-0.47		Jonsson 1959	4
"	Southern North Sea	126.	0.219	-		Lefranc 1966	4

SPECIES NAME		SAMPLING LOCATION	Lo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family GADIDAE								
Gadus morhua		North Eastern North Sea	105.	0.134		(200 m)	Lundbeck 1953	4
"	"	Northern North Sea	126.	0.209		(200 m)	"	4
"	"	Celtic Sea	126.	0.217			Letaconnoux 1954	4
"	"	ICNAF 3K 51° N 53° W	66.6	0.419			Figuera 1964	4
"	"	ICNAF 1B 66° N 54° W	90.6	0.236		(200 m)	"	4
"	"	ICNAF 3L 46° N 47° W	101.	0.125		(200 m)	"	4
"	"	Northern North Sea 60° N 3° E	119.	0.269		(200 m)	Daan 1974	4
"	"	Southern New England	98.8	0.281		(200 m)	Schoeder 1930	00
"	"	Lofoten	95.9	0.284		(200 m)	Rollefsen 1934	00
"	"	ICNAF Subarea 4	124.	0.118			Martin 1953	00
"	"	Southern Baltic	120.	0.13			Draganik & Netzel	0
"	"	ICNAF 3L 46° N 47° W	83.9	0.245			Figuera 1964	4
"	"	Faroe Bank	115.	0.19		(200 m)	Jones 1966	0
"	"	ICNAF Area Ps	101.	0.17			May et al 1965	0
"	"	ICNAF Area N-0	130.	0.12			"	0
"	"	ICNAF Area 3L	102.	0.16			"	0
"	"	ICNAF 2J	65.	0.31			"	0
"	"	ICNAF Pn	78.	0.25			"	0
"	"	ICNAF 3M	98.	0.15			"	0
"	"	ICNAF 3K	77.	0.26			"	0

SPECIES NAME	SAMPLING LOCATION	Lo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family GADIDAE							
Gadus morhua	ICNAF 2J 53° N 53° W	68.5	0.241	-1.57		Figueras 1964	4
"	ICNAF 4V 44° N 56° W	110.	0.080	-		"	4
"	Eastern Iceland	110.	0.147	-		Jonsson 1965	4
"	ICNAF Area 2H	64.	0.24	-		May et al 1965	0
"	ICNAF 1B 66° N 54° W	87.	0.314	-		Figueras 1964	4
"	ICNAF 3K 51° N 53° W	66.4	0.501	-		"	4
"	Faroe Plateau 62° N 7° E	115.	0.19	-0.42	M = 0.17	Jones 1966	0
"	East Iceland	147.	0.096	-		Saemundsson 1923	00
"	South Iceland	128.	0.147	-		"	00
Onos cimbrius	Eckernförde Bay, Baltic	43.	0.245	-		Hoffmeister 1957	4
Brosimius brosme ♂	62° N 5° W	77.6	0.162	-		Joennes 1961	4
"	"	84.3	0.135	-		"	4
Gadiculus thori	Bay of Biscay 200 m	15.5	0.693	-0.37		Letaconnoux 1947	
Boreogadus saida	Arctic Ocean	22.	0.67	-		in: Beverton & Holt 1959	0
"	Barents Sea	23.6	0.472	-		Nikolsky 1957	4
Gadus macrocephalus	Bering Sea (Virgin stock)	105.	0.157	-		Mosher 1953	4
"	Hecate Strait (Canada)	94.	0.27	-		Ketchen 1964	00
Pollachius pollachius	Southern Ireland	85.6	0.186			Moreau 1964	2

SPECIES NAME	SAMPLING LOCATION	Lo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family GADIDAE							
Melanogrammus aeglefinus	North Sea	53.	0.20	-		Beverton & Holt 1957	00
"	ICNAF Subarea 4 (N. Scotia Shelf)	67.5	0.195	-		Beverton 1965	0
"	ICNAF Subarea 3 (Grand Bank)	57.5	0.22	-		"	0
"	Faroe	77.9	0.22	-		Jones 1962	0
"		96.8	0.115	-0.69		Nikolsky 1957	4
"	Rockall Island	43.8	0.269	-0.56		Blacker 1971	4
"	North Sea, "fast growth area"	58.1	0.24	-		Jones 1962	0
"	" " "slow growth area"	48.3	0.28	-		Jones 1962	0
"	Norway 63° N	89.0	0.123	-1.72		Andersen 1938	4
"	Iceland	81.0	0.228	-0.25		Blacker 1971	4
"	Skagerrak	115.	0.084	-1.39		Andersen 1938	4
"	Barents Sea	75.5	0.234	-		Blacker 1971	4
"	ICNAF Subarea 5 (Georges Bank)	73.	0.28	-	M = 2.0	Beverton 1965	0
Trisopterus minutus ♂	English Channel	20.	0.42	-	M = 1.1	Menon 1950	00
" " ♀	" "	24.	0.40	-	M 0.9	"	00
" esmarkii	Scotland 59° N 0° longitude	19.0	0.44	-		Raitt 1968	0
"	" "	19.3	0.59	-	M = 1.6	"	0
"	Aberdeen Bay & Moray Firth	27.2	0.363	-		Fulton 1901	5



SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family GADIDAE							
Theragra chalcogramma	Bering Sea (Virgin stock?)	75.8	0.126	-		Mosher 1953	4
"	Bering Sea	127.	0.078	-		Nikolsky 1957	4
"	"	79.8	0.097	-		Hirschhorn 1974	4
"	"	94.4	0.092	-		"	4
Molva molva ♂	North Atlantic 62° N 5° W	132.	0.170	-		Joenoës 1961	41
" ♀	"	183.	0.118	-		"	41
Molva byrkelange ♂	"	113.	0.157	-		"	41
" ♀	"	155.	0.126	-		"	41
Micromestitius poutassou	Faroe Islands	33.4	0.23	-2.94		Raitt 1966	0
"	Costa Brava, Spain, West Medit.	27.9	0.60	-0.91		Bas 1963	00
"	Tuscan Archipelago, Italy	28.1	0.48	-1.60		Matta 1959	00
"	Iceland	34.7	0.36	-1.07		Saemundsson 1929	4
"	Scotland 57° N 11° W	39.9	0.15	-3.51		Raitt 1966	0
"	Eastern Spanish Coast, Medit.	31.8	0.365	-1.19		Bas 1965	4
Pollachius virens ♂	Norway 63° N 150 m	87.7	0.211	-0.04		Damas 1909	4
"	"	120.	0.15	-0.58	M = 0.3	Schmidt 1959	00
"	66° N 17° W	128.	0.13	-0.90	M = 0.3	Jones & Jonsson 1971	0
"	-	106.	0.14	-1.61		Nikolsky 1957	4
"	Norwegian Sea	107.	0.19	-	M = 0.15	Gottlieb 1957	00

SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family GADIDAE							
Lota lota	Petschora River USSR	84.4	0.135	-		Nikolsky 1957	2
"	Ilmen Lake USSR	66.1	0.210	-		"	2
"	Oderhaff, Northern GDR	88.	0.24	-	very	Müller 1960	2
"	Spree River (Neu Zittau) GDR	74.	0.11	-	tentative	"	2
the following data on M. merlangus have been compiled in cooperation with U. Damm, IFM							
Merlangius merlangus	N + W North Sea YCL 1952	37.7	0.426	-0.299		Damm 1975	4
"	" " " 1953	41.4	0.250	-1.71		"	4
"	" " " 1954	34.5	0.524	-0.467		"	4
"	" " " 1955	33.8	0.537	-0.704		"	4
"	" " " 1957	36.6	0.500	-0.559		"	4
"	" " " 1958	42.2	0.323	-0.731		"	4
"	" " " 1959	42.7	0.340	-0.698	YCL 1956 and	"	4
"	" " " 1960	42.5	0.357	-0.720	1964 data not	"	4
"	" " " 1961	37.3	0.440	-0.847	asymptotic	"	4
"	" " " 1962	38.2	0.405	-0.507		"	4
"	" " " 1963	36.4	0.506	-0.317		"	4
"	" " " 1965	40.3	0.471	-0.340		"	4
"	" " " 1966	41.3	0.222	-2.36		"	4
"	" " " 1967	32.9	0.271	0.032		"	4
"	SE North Sea " 1954	68.9	0.0905	-2.05		"	4

SPECIES NAME	SAMPLING LOCATION		Lo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family GADIDAE								
Merlangius merlangus	SE North Sea	YCL 1955	28.5	1.25	0.464		Dunn 1975	4
"	"	" 1957	94.7	0.0742	-1.48		"	4
"	"	" 1958	40.4	0.294	-0.874	YCL 1956 data	"	4
"	"	" 1959	39.9	0.324	-0.853		"	4
"	"	" 1960	57.4	0.204	-0.688	not	"	4
"	"	" 1961	45.3	0.240	-1.242	asymptotic	"	4
"	"	" 1962	63.7	0.150	-1.02		"	4
"	"	" 1963	37.4	0.492	-0.160		"	4
"	"	" 1964	36.0	0.497	-0.368		"	4
"	"	" 1965	40.1	0.324	-1.02		"	4
"	"	" 1966	37.7	0.301	-1.57		"	4
"	"	" 1967	33.2	0.633	-0.173		"	4
"	Moray Firth & Orkney area, North Sea	YCL 1958	66.3	0.146	-1.23		Jones & Hislop 1975	4
"	"	" 1959	62.1	0.197	-0.627		"	4
"	"	" 1960	31.1	0.323	0.154	YCL 1957 & 1963	"	4
"	"	" 1961	39.8	0.396	-0.613	data not	"	4
"	"	" 1962	37.7	0.415	-0.375	asymptotic	"	4
"	"	" 1964	47.2	0.292	-0.751		"	4
"	"	" 1965	36.3	0.619	-0.495		"	4
"	"	" 1966	37.0	0.496	-0.516		"	4

SPECIES NAME	SAMPLING LOCATION		Lo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family GADIDAE								
Merlangius merlangus	Moray Firth & Orkney Area, North Sea	YCL 1967	53.7	0.140	-2.28		Jones & Hislop 1975	4
"	"	" 1968	34.3	0.741	0.082		"	4
"	"	" 1969	36.8	0.738	0.119		"	1
"	Northern area, North Sea	" 1959	37.8	0.567	-0.167		"	4
"	"	" 1960	52.2	0.167	-2.23		"	4
"	"	" 1961	43.42	0.180	-3.37		"	4
"	"	" 1962	80.9	0.0704	-3.176		"	4
"	"	" 1963	33.9	1.791	1.18		"	4
"	"	" 1964	75.6	0.107	-2.08	YCL 1957, 1958, " 1965, 1971,	"	4
"	"	" 1966	50.	0.180	-1.953	data not	"	1
"	"	" 1967	62.5	0.116	-3.70	asymptotic	"	4
"	"	" 1969	58.0	0.230	-0.912		"	4
"	"	" 1970	92.4	0.080	-2.675		"	4
"	North central area, North Sea	" 1957	33.8	0.556	-0.217		"	4
"	"	" 1959	38.1	0.432	-0.500		"	4
"	"	" 1960	34.0	0.653	-0.165	YCL 1958 data	"	4
"	"	" 1961	38.0	0.370	0.785	not asymptotic	"	4
"	"	" 1962	55.2	0.158	-1.46			
"	"	" 1963	39.0	0.560	0.018		"	4
"	"	" 1964	50.	0.272	-0.657		"	1

SPECIES NAME	SAMPLING LOCATION	Lo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family GADIDAE							
Merlangius merlangus	North Central area, North Sea	49.2	0.256	-1.05		Jones & Hislop	4
"	"	35.8	0.656	-0.070	YCL 1967 data	"	4
"	"	50.8	0.198	-0.803	not asymptotic	"	4
"	"	50.	0.280	-0.995	"	"	1
"	"	43.6	0.278	-1.73	"	"	4
"	Central area, North Sea	32.9	0.649	-0.208	"	"	4
"	"	45.5	0.278	-0.852	"	"	4
"	"	57.3	0.229	-0.513	"	"	4
"	"	29.5	1.533	0.307	"	"	4
"	"	45.9	0.217	-1.09	"	"	4
"	"	31.9	0.928	-0.065	"	"	4
"	"	35.1	0.714	-0.096	YCL 1958 data	"	4
"	"	66.4	0.105	-2.47	not asymptotic	"	4
"	"	47.5	0.207	-1.246	"	"	4
"	"	36.5	0.632	-0.216	"	"	4
"	"	56.5	0.173	-1.64	"	"	4
"	"	35.4	0.548	-0.767	"	"	4
"	N + E North Sea & Morray Firth	44.2	0.383	-0.44		Ellis & Jones 1956	0
"	SE North Sea	40.0	0.29	-0.51		Knudsen 1964	0
"	W North Sea	44.44	0.23	-1.07		"	0

SPECIES NAME		SAMPLING LOCATION		Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family GADIDAE									
Merlangius merlangus									
"	"	Skagerak		39.53	0.33	-0.55		Knudsen 1964	0
"	"	Skagerak - Kattegat		39.9	0.69	-	tagging data	"	0
"	"	NW North Sea		37.4	0.47	-	"	"	0
"	"	Skagerak		43.9	0.266	-1.132		in: Desbrosses 1948	4
"	"	Skagerak - Kattegat	N	62.4	0.162	-		in: Messtorff 1959	4
"	"	Belt Sea	S	37.8	0.458	-		"	4
"	♀	Bay of Biscay		70.	0.238	-		in: Desbrosses 1948	1
"	"	Iceland		71.0	0.250	-		Saemundsson 1925	4
"	♂	Irish Sea		54.7	0.151	-		Bowers 1954	4
"	"	Clyde, Scotland		54.5	0.322	-0.48		Garrod & Gambell 1965	0
"	"	West Coast, Scotland		45.0	0.483	-0.42		"	0
"	"	Baltic Sea		44.0	0.247	-		Hoffmeister 1957	00
"	♂	North Sea	YCL 1958	32.1	0.729	-0.0797		Hannerz 1964	00
"	♀	"	"	38.9	0.492	-0.0567	back calculated	"	00
"	"	"	"	36.3	0.554	-0.0333	lengths from	"	00
"	♂	"	1959	39.9	0.399	-0.273	3-years old	"	00
"	♀	"	"	47.3	0.318	-0.199		"	00
"	"	"	"	44.2	0.341	-0.265		"	00

SPECIES NAME	SAMPLING LOCATION	Lo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family GASTEOSTEIDAE							
Gasteosteus aculeatus ♂	Brackish Waters, Roscoff, France	4.6	4.20	-0.01		Bertin 1923	4
" " ♀	"	6.0	2.40	-0.04		"	4
"	Kiel Bay, eelgrass belt	6.9	1.79	-0.285		Northmann 1975	4
"	Ooster Schelde, Holland	6.7	2.32	-		van Mullem & van der Vlugt 1964	4
Apeltes quadracus ♂	Chesapeake Bay, eelgrass	4.8	1.50	-		Schwarz 1965	41
" " ♀	"	7.0	0.84	-		"	41
Spinachia spinachia	Kiel Bight, eelgrass	17.	1.78	-		Northmann 1975	41
Pygosteus pungitius	Cheshire, England	4.3	1.60	-	M = 1.1	Jones & Hines 1950	00
Family GERREIDAE							
Gerres punctatus	Ambaro Bay, Madagascar	21.9	2.616	-		Chabanne & Pianté 1969	51
"	"	32.9	0.756	-		"	51
"	"	26.9	1.788	-		"	51
Diapterus rhombeus	Puerto Rico, Southwest	22.0	2.196	-		Austin 1971	51

SPECIES NAME	SAMPLING LOCATION		Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family GOBIIDAE								
Gobiosoma robustum ♂	Tampa Bay, Florida		3.6	2.006	-		Springer & McErlan 1961	5
" " ♀	" "		6.	0.813	-		"	5
Gobius paganellus	tidal pools, Isle of Man		11.6	0.408	-	M = 0.57	Miller 1961	4
Microgobius thalassinus ♂	Chesapeake Bay, 10-15 m		5.5	2.407	-		Schwarz 1971	1
" " ♀	" "		6.9	1.388	-		"	1
Pomatochistus microps	Isle of Man, River mouths		11.2	0.295	-		Miller 1975	4
" minutus	Dutch Wadden Sea		9.2	0.928	-0.01		Fonds 1973	0
" "	Gullmarsfjord, Sweden 58° N		7.1	1.38	-0.32		Swedmark 1957	4
Gobius cobitis	Roscoff, France		25.3	0.217	-		Gibson 1970	4
Family HARPADONTIDAE								
Harpadon neherus	Bombay, India		35.	0.52	-		Bapat 1970	41
" "	"		28.4	0.85	-		Bapat <u>et al.</u> 1952	41
Family HOLOCENTRIDAE								
Holocentrus diadema	Eilat, Red Sea		16.8	1.129	-0.12		Gundermann & Popper 1975	4



SPECIES NAME	SAMPLING LOCATION	LOO	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family ICTALURIDAE							
Ictalurus punctatus	St. Laurent River, Quebec	83.4	0.056	-1.92		Magnin & Fradette 1975	0
Family ISTIOPHORIDAE (LF and including bills)							
Istiophorus platypterus	Western North Atlantic	220.	1.10	-		De Sylva 1957	00
"	"	243.	1.20	-		Mather III et al. 1974	3
"	East China Sea	341.	0.442	-		Koto & Kodama 1962	3
Makaira nigricans ♂	Hawaii	371.	0.285	-		Skillman & Young 1976	0
" ♀	"	659.	0.116	-		"	0
Tetrapterus albidus	"	261.	0.58	-	tagging, best growth only	Mather et al. 1974	3
" audax	Western North Pacific	273.	0.262	-		Koto 1963	3
"	East Africa 5° S 43° E 400 m	312.	0.201	-		Nerett 1971	4
" ♂	Hawaii	277.	0.417	-0.52		Skillman & Young 1976	0
" ♀	"	240.	0.810	-		"	0

SPECIES NAME	SAMPLING LOCATION	Loo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family LABRIDAE							
<i>Ctenilabrus tinca</i> ♂	Black Sea, Soviet Coast	28.4	0.183	-		Kafinina 1963	41
" " ♀	" "	21.4	0.186	-		"	4
" <i>quinquemaculatus</i> ♂	" "	16.7	0.303	-		"	4
" " ♀	" "	15.0	0.200	-		"	4
" <i>rupestris</i>	Nyborg Fjord, Denmark	20.	0.412	-		In: Johansen 1925	41
" " ♂	Roscoff, France	16.1	0.307	-		Guinard 1966	4
" " ♀	" "	15.4	0.297	-		"	4
<i>Labrus berggylta</i> ♂	" "	57.8	0.113	-		"	4
" " ♀	" "	61.1	0.100	-		"	4
" "	St. Andrew Bay, Western Gr. Brit.	38.4	0.194	-	1 specimen (backcalculated)	Meek 1916	4
" <i>bimaculatus</i> ♂	Sète, Mediterranean, France	34.1	0.133	-		Guinard 1966	4
" " ♀	" "	42.9	0.093	-		"	4
<i>Symphodus cinereus</i> ♂	Arcachon, France	15.5	0.348	-		"	4
" " ♀	" "	14.3	0.291	-		"	4
" <i>exoletus</i> ♂	Sud Finistère (Brettony, France)	12.9	0.713	-		"	4
" " ♀	" "	12.3	0.618	-		"	4
" <i>mediterraneus</i> ♂	Sète, Mediterranean, France	17.0	0.628	-		"	4
" " ♀	" "	15.7	0.470	-		"	4
" <i>melops</i> ♂	Roscoff, France	24.2	0.409	-		"	4
" " ♀	" "	21.4	0.308	-		"	4

SPECIES NAME	SAMPLING LOCATION	Lo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family LABRIDAE							
<i>Symphodus ocellatus</i> ♂	Toulon, France	10.8	0.949	-		Guinard 1966	4
" " ♀	" "	10.3	0.645	-		"	4
" <i>roissali</i> ♂	Sète, Mediterranean	13.6	0.617	-		"	4
" " ♀	" "	13.0	0.593	-		"	4
" <i>rostratus</i> ♂	Toulon, France	11.6	0.898	-		"	4
" " ♀	" "	10.6	0.709	-		"	41
" "	Villefranche-sur-Mer, France	14.2	0.363	-0.69		Nival 1966	41
" <i>tinca</i> ♂	Sète, Mediterranean	31.3	0.248	-		Guinard 1966	4
" " ♀	" "	24.8	0.244	-		"	4
<i>Tautoga onitis</i> ♂	Rhodes Islands, USA	58.2	0.147	-		Cooper 1967	4
" " ♀	" "	49.5	0.182	-		"	4
<i>Tautogalabrus adspersus</i>	Weweeantic River Estuary, Mass. USA	28.5	0.198	-		Serchuk & Cole 1974	0
" " ♂	41°20' N 71°55' W 20 m	29.7	0.222	-		Dew 1976	4
" " ♀	" "	24.2	0.239	-		"	4
" "	Gulf of St. Lawrence	37.	0.152	-		Johansen 1925	41
<i>Thalassoma bifasciatum</i>	Florida (open air aquarium)	17.	0.75	-		Feddern 1965	4
<i>Xyrichtys novacula</i>	Balearic Islands	26.1	0.264	-		In: Guinard 1966	4
<i>Labrus merula</i> ♂	Sète, France	44.	0.215	-		Guinard 1966	1
" " ♀	" "	35.1	0.252	-		"	2

SPECIES NAME	SAMPLING LOCATION	Loo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family LACTARIIDAE							
Lactarius lactarius	Waltair, India	32.	0.854	-		Apparao 1966	5
Family LAMNIDAE							
Lamna nasus	Atlantic, 44° N 62° W 30 m	280.	0.111	-	M 0.113	Aasen 1963	00
Family LATIMERIDAE							
Latimeria chalumnae ♂	East Africa	150.	0.26	-0.94	very rough	Hureau & Ozouf 1977	1
" ♀	East Africa	200.	0.15	-1.16	estimates	"	1

SPECIES NAME	SAMPLING LOCATION		Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family LEOGNATHIDAE								
Leiognathus bindus	Calicut, India		12.7	2.04		-	Balan 1967	5
"	Manilla Bay, Philippines		11.3	2.06	Feb. brood		Tiews et al 1965 *	5
"	"	"	10.5	2.24	May	"	"	5
"	"	"	10.2	2.63	Sept.	"	"	5
"	"	"	10.3	2.58	Dec.	"	"	5
"	"	"	9.5	2.42	March	"	"	5
" daura	"	"	8.1	2.93	Jan.	"	"	5
"	"	"	8.5	2.52	Sept.	"	"	5
" equulus	Ambaro Bay, Madagascar		21.1	1.88	July	"	Chabanne & Plante 1969	5
"	"	"	21.2	1.73	Sept.	"	"	5
" lineolatus	Manilla Bay, Philippines		9.8	2.10	Febr.	"	Tiews et al 1965	5
"	"	"	8.8	2.60	Dec.	"	"	5
"	"	"	8.6	2.83	Jan.	"	"	5
" nuchalis	Kasaoka Bay, Japan		12.0	2.56	-		Kakuda, pers. comm.	5
" splendens	San Miguel Bay, Philippines		13.5	1.03	April	"	Tiews et al. 1965	5
"	"	"	11.7	1.12	March	"	"	5
"	"	"	11.7	1.20	March	"	"	5
"	"	"	13.0	1.60	Aug.	"	"	5
"	"	"	14.3	1.04	April	"	"	5

\*Tiews et al (1965) do not state which length was measured. Total length ?

SPECIES NAME	SAMPLING LOCATION	Lo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family LETHRINIDAE							
Lethrinus enigmaticus	Indian Ocean 10°30'N 62° E	62.6	0.149	-1.21	} M = 0.2	Lebeau & Cueff 1975	0
"	Indian Ocean 10°30'N 62° E	52.6	0.179	-1.07		"	0
" lentjan	Gulf of Manaar, India	64.0	0.27	-		Toor 1964	0
Family LOPHIIDAE							
Lophius piscatorius	Bay of Fundy, Canada	197.	0.060	-0.08		Connolly 1920	4
"	Aberdeen, Scotland 40 m	137.	0.164	-0.06		Fulton 1903	41
Family LUTJANIDAE							
Lutjanus griseus	Everglades, Florida	55.	0.167	- V		Crocker 1962	41
" purpureus	off Recife, Brazil	96.7	0.096	-1.20		Ponteles-Filho 1970	4
" synagris	Cuban Waters	42.2	0.268	-0.44 V		Rodriguez-Pino 1962	4
"	off Caera State, Brazil	50.5	0.231	-0.19		Alegria & Ferrairade Menezes 1970	0
" purpureus	North Eastern & Northern Brazil	98.9	0.090	-	M≈0.37 from other source	Ferreira de Menezes & Vasconcelof Gesteira 1974	0

SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>o</sub>	NOTES	SOURCE	METHOD
Family MACRORHAMPHOSIDAE							
Macrorhamphosus scolopax	Meteor Sea Mound, Cent. East Atl.	16.5	0.745	-0.24		Ehrich 1976	0
Family MACROURIDAE							
Macrurus rupestris ♂	N-W Atlantic 2000 m	105.	0.082	-		Sarratimskii 1973	41
" ♀	"	110.	0.099	-		"	41
Nezumia glerorhynchus	Alboran Sea 500-800 m	26.4	0.160	-		Rannou 1976	0
" steglipedis	Gulf of California 500 m	43.	0.20	-		Mathew 1974	4
Trachyrinchus trachyrinchus ♂	off Monaco, Mediterranean, 500 m	45.6	0.347	-0.95		Motais 1960	4
" ♀	"	49.3	0.222	-2.28		"	4
Family MAENIDAE							
Maena smarís	Adriatic	21.	0.218	-1.79		Zel 1951	41

SPECIES NAME	SAMPLING LOCATION	Loo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family MERLUCCIDAE							
Merluccius merluccius	Gulf of Biscay	105.	0.184	-		Belloc 1929	4
"	off Morocco	85.	0.210	-		Maurin 1954	41
"	off Morocco 30-40° N	100.	0.158	-		Belloc 1929	4
"	off Ireland	83.9	0.296	-		"	4
"	off Morocco, Agadir Bight	112.	0.121	-		Maurin 1954	41
"	Mediterranean, off Fano, Italy	42.1	0.405	-0.01		Ghirardelli 1959	41
"	Mediterranean, North Africa	38.4	0.220	-		Belloc 1929	4
"	Marmara Sea	44.	0.13	-	M = 0.6	Akyüz 1959	00
"	Marmara Sea	60.	0.10	-	M = 0.5	"	00
"	Central Adriatic 43°20' N 15°10'E	62.	0.344	-		Zupanovič 1968	4
"	Mediterranean	65.3	0.181	-		Letaconnoux 1955	4
"	off Tunisia	40.	0.2	-		Matta 1956	4
"	Can Farré Bank, Costa Brava	28.9	0.634	-	400 m	Figueras 1967	4
"	Malica Bank, Costa Brava, Spain	59.6	0.39	-	500 m	"	4
"	Barana Bank,	60.8	0.255	-	700 m	"	4
"	Capet Bank,	19.7	0.446	-	90 m	"	4
"	off Castellon, Spain	43.3	0.142	-		"	4



SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family MERLUCCIDAE							
Merluccius gayi ♂	off Chile	54.0	0.428	-		Aguayo 1971	4
" ♀	off Chile	81.9	0.203	-		"	4
" ♂	off Peru 7° S 80-81° W	67.9	0.190	-0.38		Misa & Hamasaki 1971	0
" ♀	off Peru 7° S 80-81° W	116.	0.092	-		"	0
"	off Chile, Valparaíso	106.	0.100	-0.07		Alberti & Zuniga 1966	00
" ♂	off Chile	49.	0.45	-		Saetersdal & Villegas 1968	0
" ♀	off Chile	62.	0.36	-		"	0
" cadenati	off Dakar, Senegal 200 m	64.4	0.394	-		Doutre 1960	41
" hubbsi ♂	46° S 63° W, Argentina, 90 m	51.5	0.308	-0.14		Rojo & Silvosa 1969	4
" ♀	"	85.9	0.128	-0.59		"	41
" productus ♂	off Portland, Oregon 100 m	56.3	0.39	-	} M = 0.56 Ehrlich, pers. comm.	Dark 1975	0
" ♀	"	61.2	0.30	-		"	0
" senegalensis	off Dakar, Senegal 200 m	85.2	0.171	-0.61		Doutre 1960	41
" augustimanus	Gulf of California, Southeast	38.	0.335	-		Mathews 1974	41
" capensis	South Africa	141.	0.091	-		Botha 1971	0

1  
8  
0  
1

SPECIES NAME		SAMPLING LOCATION		Loo	K	t <sub>o</sub>	NOTES	SOURCE	METHOD
Family MUGILIDAE									
Mugil cephalus	♂	Taiwan Waters	15 m	49.8	0.393	0.05	LT ?	Ih-Hsiu Tung 1970	0
"	♀	"	"	59.3	0.301	-0.12	M = 0.31	"	0
"	♂	Tunisian Coast		48.6	0.29	-		Farrugio 1975	0
"	♀	"	"	54.6	0.20	-	LT ?	"	0
"	"	Turkey (Bosphorus)		105.	0.11	-		Erman 1959	4
"	♂	Texas Coast		(40.)	0.367	-0.48		Cech & Wohlschlag 1975	4
"	♀	"	"	42.2	0.327	-0.70		"	4
"	"	Sea of Marmara, Turkey		71.5	0.254	-		Erman 1959	41
"	"	Australia		65.	0.271	-0.15	LT ?	Kesteven 1942	4
"	labrosus	Tunisian Lagoons		(37.8)	0.36	0.06		Farrugio 1975	0
"	♀	"	"	57.4	0.17	-0.51		"	0
"	labrosus	Caspian Sea		47.1	0.188	-1.33		Nikolsky 1957	4
"	ramada	Tunisian Lagoons		31.6	0.45	-0.40		Farrugio 1975	0
"	♀	"	"	(40.5)	0.31	-0.23		"	0
"	saliens	-		34.3	0.208	-0.48		Aleev 1956	4
"	♂	Lake Quarum (brackish)	Egypt	(36.5)	0.288	-1.23		El Zarka & El Zedfy 1970	4
"	♀	"	"	41.9	0.195	-1.90		"	4
"	cunnesius	Bengal, India		29.2	0.31	-		Sarofini 1958	00
"	parsia	"	"	30.7	0.32	-	LT ?	" 1957	00

SPECIES NAME	SAMPLING LOCATION	L <sub>50</sub>	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family MUGILIDAE							
<i>Liza provinsalis</i>	-	44.1	0.282	-0.86		Nikolsky 1957	4
" <i>aurata</i>	Brackish Waters, Italy	53.7	0.341	-		in: Bougis 1952	4
"	Black Sea	54.1	0.199	-		Nikolsky 1957	4
"	Gulf of Biscay	34.5	0.366			Arné 1938	4
"	Caspian	66.	0.169			Nikolsky 1957	4
Family MYCTOPHIDAE							
<i>Benthoosema glaciale</i>	Canadian Waters (East)	8.5	0.36	-0.23	M = 1.75	Halliday 1970	00
"	Norwegian Fjords, below 100 m	8.6	0.45	-	M = 0.74	Gjøsæter 1970	0
<i>Lobianchia dofleini</i>	off Bermuda	4.6	1.386	-		Karnella & Gibbs 19	1
<i>Myctophum</i> affine	29-31° N 149-154° E	10.0	0.42	-0.23		Odote 1966	4
" <i>punctatum</i>	North West African, upwelling Area	9.0	0.32	-0.91		Wörner 1975	41
<i>Notolychnus valdiviae</i>	24° S 110° E 200 m	2.8	1.411	-0.16		Legand 1967	4
<i>Stenobrachius leucopsaurus</i>	off Monterey Bay, California	14.3	0.244	-0.23		Bolin 1956	4

SPECIES NAME	SAMPLING LOCATION	Loo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family NEMIPTERIDAE							
Nemipterus bathybius ♂	Hongkong Shelf 50 m	30.2	0.324	-0.80		Eggleston 1972	41
" bleekeri	off Brunei 30-40 m	30.4	0.439	-	M = 1.4	Weber & Jothy , undated	41
" hexodon	" "	27.8	0.494	-	M = 1.0	"	41
" delagoae	" "	27.8	0.701	-	M = 2.2	"	41
" marginatus	" "	28.4	0.368	-	M = 1.6	"	41
" mesoprion	" "	19.5	0.629	-	M = 0.6	"	41
" nematophorus	" "	27.3	0.531	-	M = 1.9	"	41
" nemurus	" "	28.9	0.264	-	M = 1.3	"	41
" peronli	" "	28.9	0.455	-	M 1.4	"	41
" tolu	" "	26.8	0.523	-	M 0.4	"	41
" sp.	" "	31.5	0.440	-	M 1.3	"	41
" virgatus ♂	112,5° E 20,8° N 70 m	37.6	0.397	-0.42		Eggleston 1970	4
" ♀	" "	35.	0.237	-		" 1972	41
" ♂	} Hongkong, South West of Pearl River Mouth 50 m	44.7	0.408	-0.12		"	4
" ♀		36.2	0.352	-0.71		"	4
"	21.3° N 114° E	44.0	0.348	-		Li Kwang Ming 1960	4
"	" "	47.0	0.298	-		"	4

SPECIES NAME	SAMPLING LOCATION	Loo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family NEMIPTERIDAE							
Nemipterus japonicus	1964-65 Andhra-Orissa Coast, India	30.5	0.314	-1.11		Krisnamoorthi 1971	0
"	1965-66	"	0.648	-	" draught period "	"	0
"	1966-67	"	0.294	-		"	0
"	Hong Kong	38.	0.13	-		Jones 1976	0
"	"	34.	0.19	-		"	0
"	off Brunei 30-40 m	28.9	0.470	-	M = 2.3	Weber & Jothy MS	41

SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family NOTOTHENIDAE							
Nototothenia cyanobranchia	Kerguelen Islands, Antarctica	39.1	0.186	-		Hureau 1970	0
" macrocephala	"	46.4	0.26	-0.4		"	0
" neglecta ♂	Sydney Islands	45.3	0.129	-0.8	} M = 0.36	Everson 1970	0
" " ♀	"	51.6	0.091	-1.7		"	0
" " ♂	Terre Adélie	36.	0.22	-		Hureau 1970	41
" " ♀	"	48.	0.22	-		"	41
" rossii ♂	South Georgia	86.	0.128	-		Olsen 1954	4
" " ♀	"	92.	0.124	-		"	4
" " "	"	44.7	0.293	-		Crisp & Carrick 1975	4
Trematomus bernachii ♂	Terre Adélie	28.6	0.185			Hureau 1970	4
" " ♀	"	31.4	0.192			"	4
" " ♂	McMurdo Sound	20.0	0.557			Wohlschag 1961	4
" " ♀	"	25.5	0.370			"	4
" " ♂	"	23.	0.36			" 1962	0
" " ♀	"	30.7	0.42			"	0
" hansonii ♂	Terre Adélie	26.8	0.255	-		Hureau 1970	41
" " ♀	"	37.3	0.229	-		"	41

SPECIES NAME	SAMPLING LOCATION	L <sub>0</sub>	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family OSMERIDAE							
Osmerus eperlanus	Elbe Estuary, FRG	32.2	0.274	-		Lillelund 1961	4
"	Ilmen Lake, USSR	17.0	0.336	-		in: Belyanina 1958	2
"	Rybinsk Reservoir, USSR	9.6	1.386	-		"	2
"	Dadey Lake (formerly Eastern Prussia)	13.3	0.875	-		"	2
"	Onega Lake, USSR	12.6	0.370	-		"	2
"	Kurishes Haff, Baltic	32.4	0.257	-	Sea migrants	"	2
"	Neva River, USSR	25.1	0.210	-		"	2
"	Basin of Maine River, USA	31.8	0.313	-		"	2
"	White Sea, Dvina Bay	38.3	0.152	-		"	2
"	" , Kandalaksha Bay	30.9	0.379	-		"	2
"	Chesha Bay	31.8	0.181	-		"	2
"	Yenisey River, USSR	28.8	0.252	-		"	2
Mallotus villosus ♂	Labrador	20.	0.48	-	} M = 1.3	Templeman 1948	00
" ♀	"	19.	0.48	-		"	00
" ♂	Barents Sea & Finmark Coast	27.8	0.259	-0.82		Dragesund et al. 1973	4
" ♀	"	19.8	0.454	-0.53		"	4

SPECIES NAME	SAMPLING LOCATION	L <sub>∞</sub>	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family PANGASIDAE							
Pangasius pangasius	Ganga & Hoogly Rivers, India	227.	0.043	0.56	bad fit, L <sub>∞</sub> too high	Pantulu 1962	0
Family PLATYCEPHALIDAE							
Neoplatycephalus macrodon	Marion Bay, Tasmania 43° S	65.2	0.197	-	≈ 50 m	Fairbridge 1951a	4
"	Eastern Australia 35°30' S	79.1	0.175	-	"	"	4
"	" 34° S	65.6	0.246	-	"	"	4
"	" 37° S	56.5	0.302	-	"	"	4
"	♂ New South Wales, Australia	52.0	0.376	-	"	Fairbridge 1951b	4
"	♀ " "	62.2	0.292	-	"	"	4



SPECIES NAME		SAMPLING LOCATION		Lco	K	t <sub>o</sub>	NOTES	SOURCE	METHOD
Family PERCIDAE									
Perca	schenkli	lower Illi River, USSR		60.	0.069	-	p. 404	Nikolsky 1957	2
"	flavescens ♂	Saginaw Bay, Lake Huron		42.9	0.193	-		Hile & Jobe 1941	2
"	" ♀	"		49.3	0.165	-		"	2
"	fluviatilis	Aral Lake, USSR		23.3	0.405	-		Nikolsky 1957	2
"	"	Pestchora, USSR		38.0	0.124	-		"	2
"	"	Sweden		30.	0.20	-	M = 0.29	Alm 1952	00
"	"	"		34.	0.13	-	M = 0.16	"	00
"	"	Orava Reservoir, Slovakia		41.9	0.123	-		Hnatevič 1960	2
"	"	-		36.9	0.119	-	"reservoir"	in: Hnatevic 1960	2
"	"	-		29.3	0.354	-	"25 lakes"	"	2
Stizostedion glaucum									
"	canadensis	Lake Erie, USA		54.4	0.237	-		Deason 1933	2
"	"	Lake Nipigon, Canada		40.	0.14	-	M = 0.44	Beverton & Holt 1959	0
"	"	Lake Erie, USA		54.4	0.237	-		Deason 1933	2
"	vitreum ♂	Lake La Ronge, Canada 10 m		110.	0.055	-	} max. c.p.u.e. at 16.5° C	Rawson 1956	2
"	" ♀	"		137.	0.037	-		"	2
Lucioperca lucioperca									
"	"	Kuban River, USSR		85.6	0.238	-		Nikolsky 1957	2
"	"	Don River, USSR		86.	0.168	-		"	2
"	"	Aral Lake, USSR		79.5	0.204	-		"	2
"	"	Ilmen Lake, USSR		104.	0.129	-		"	2
"	"	Southern Caspian, USSR		40.	0.333	-		"	2

SPECIES NAME	SAMPLING LOCATION	LOO	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family PLEURONECTIDAE							
Eopsetta jordani ♂	British Columbia	49.	0.160	-2.60		Ketchen & Forrester 1966	0
" " ♀	" "	58.6	0.167	-		"	0
Isopsetta isolepis ♂	Canada, West Coast	38.	0.36	-		Hart 1948	0
" " ♀	" "	42.	0.26	-		"	0
Lepidopsetta bilineata	Bering Sea (Virgin Stock)	73.5	0.098	-		Mosher 1953	4
Glyptocephalus cynoglossus ♂	Isle of Man 70 m	30.9	0.312	-		Bower 1959	0
" " ♀	" "	36.1	0.38	-		"	0
Limanda aspera ♂	Peter the Great Bay 1928 Virgin	74.4	0.085	-		Nikolsky 1957	4
" " ♀	" " 1933 exploit.	52.2	0.148			"	4
Lophopsetta aquosa	Hell Hole, Rhodes Island	41.	0.242			Moore 1947	41
Parophrys vetulus ♂	Puget Sound, Washington	30.7	0.347			Holland 1969	4
" " ♀	" "	41.6	0.243			"	4
Rheinhardtius hippoglossoides	West Greenland 400 m				M = 0.48	Smidt 1969	4
" " "	Okhotsk Sea	109.	0.074			Nikolsky 1957	4
" " "	Barents Sea	138.	0.056			"	4
Pleuronectes platessa ♂	North Sea	45.	0.15		M = 0.22	Beverton & Holt 1959	0
" " ♀	" "	70.	0.08		M = 0.12	"	0
" " "	Barents Sea	59.	0.115			Nikolsky 1957	4
" quadrituberculatus	Bering Sea, Virgin Stock	57.	0.099			Mosher 1953	41

SPECIES NAME	SAMPLING LOCATION	L <sub>50</sub>	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family PLEURONECTIDAE							
<i>Pseudopleuronectes americanus</i>	Canada, East Coast	44.	0.4	-	M = 0.3	Dickie & McCracken 1955	00
"	Peronic & Gardiners Bay, Long Island	50.6	0.175	-		McCracken 1954	4
"	Anapolis Bay, USA	43.6	0.323	-		"	4
"	Passamaquoddy Bay	41.6	0.195	-		"	4
"	♂ Narraganset Bay, Rhode Island	43.6	0.199	-		Berryet al. 1965	4
"	♀ " "	45.5	0.283	-		Berry et al. 1965	4
"	" Northumberland Strait	39.5	0.259	-		McCracken 1954	4
"	" Pubnico Bay	57.7	0.117	-		"	4
"	♂ Charlestown "Pond", Rhode Isl.	32.6	0.428	-		Berry et al. 1965	4
"	♀ " "	39.8	0.384	-		"	4
<i>Plathychthys stellatus</i>	California 41° N 125° W	44.8	0.229	-		Orcutt 1950	4
"	" "	51.0	0.192	-		"	4
"	Bering Sea (Virgin Stock)	44.3	0.303	-		Mosher 1959	4
"	flesus ♂	39.1	0.377	-0.179	} M = 0.18	Saeger 1974	0
"	♀	46.5	0.302	-0.291		"	0
"	" "	47.3	0.229	-		Hohendorf 1966	0
<i>Drepanopsetta platessoides</i>	♂ " , Western Baltic	25.3	0.337	-0.28		Kröger 1942	4
"	♀ " "	36.1	0.428	-		"	4

SPECIES NAME	SAMPLING LOCATION	Loo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family PLEURONECTIDAE							
Hippoglossoides platessoides ♂	46° N 56° W 50-200 m	65.6	0.08	-1.21		Minet 1974	0
"	"	80.7	0.06	-1.45		"	0
"	ICNAF Res.Div. 3L 1957-60	64.2	0.08	-		Pitt 1975	0
"	"	80.2	0.076	-		"	0
"	"	72.9	0.099	-		"	0
"	"	65.9	0.084	-		"	0
"	"	64.2	0.108	-		"	0
"	"	72.5	0.125	-		"	0
"	"	72.5	0.067	-		"	0
"	"	70.4	0.093	-		"	0
Hippoglossus hippoglossus ♂	North Atlantic	170.	0.04	-		Devold 1938	00
"	"	250.	0.02	-		"	00
"	Alaska, Portlock Albatros Ground	215.	0.054	-		Southward & Chapman 1965	4
"	Bering Sea (Virgin Stock)	132.	0.105	-		Mosher 1959	41
"	Northern Norway 71° N 27° E 500 m	131.	0.2	-	} 0.20	Mathisen & Olsen 1968	0
"	"	303.	0.03	-		"	0

SPECIES NAME	SAMPLING LOCATION	Lo	K	t <sub>o</sub>	NOTES	SOURCE	METHOD
Family PLOTOSIDAE							
Plotosus anguillaris	Ago Bay, Japan	27.	0.449	-		Kajikawa 1973	4
"	Tamiooka Bay, Japan, eelgrass	28.4	0.445	-		Kikuchi 1966	4
Tandanus tandanus	North West of Sydney, Australia	59.6	0.341	-	fresh water	Davis 1977	0
Family POLYNEMIDAE							
Galedoides decadactylus	off Sierra Leone 50 m	75.2	0.139	-0.08		Longhurst 1963	4
"	off Lagos, Nigeria 50 m	54.5	0.41	-		" 1965	0
Pentanemus quinarius	off Sierra Leone 50 m	36.8	0.382	-		" 1963	41
"	off Lagos	34.	1.081	-0.14		" 1965	41
Polynemus indicus	Indian Coast (E) 22° N 50 m	150.	0.189	-0.441		Kagwade 1970	1
Family POLYDONTIDAE							
Polyodon spathula	Mississippi River, USA	113.	0.581	-		Adams 1942	4
Family POMACENTRIDAE							
Amphiprion chrysopterus	Enivetok Atoll, Pacific	11.3	0.812	-		Allen 1975	3
" melanopus	"	9.1	0.528	-		"	3
" perideraion	"	9.7	0.480	-		"	3
" tricluctus	"	16.5	0.360	-		"	3

1  
9  
2  
1

SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>o</sub>	NOTES	SOURCE	METHOD
Family POMADASYDAE							
Brachydeuterus auritus	Lagos, Nigeria 50 m	20.6	0.363	-		Raitht & Sagua 1969	4
"	"						
"	Togo 30-40 m	25.3	0.292	-		Beck 1974	0
"	Togo 30-40 m	24.7	0.322	-		"	0
"	Ivory Coast 30-50 m	25.3	0.4	-		Barro 1968	0
Pomadasys hasta	India 20° N 70° E	74.1	0.274	-		Deshmukh 1973	4
"	Bombay, India	62.2	0.496	-		Nammalwar 1974	4
Pristipoma jubelini	Sierra Leone 50 m	45.	0.3	-		Longhurst 1963	41
Parapristipoma trilineatum	Japan, Miara Peninsula	32.8	0.342	-0.43		Masuzawa 1967	4
Parapristipoma trilineatum	" " "	52.6	0.352	-		Yasuda et al. 1962	4
Rhonciscus striatus	Gulf of Suez	18.	0.300	-		Latif & Shenouda 1972	4
Family POMATOMIDAE							
Pomatomus saltatrix	Natal, South Africa	84.	0.197	-		van der Elst 1976	0
"	Cape Cod, USA	115.	0.135	-		Bachus 1962	4
Family PSETTODIDAE							
Psettodes erumei	Gulf of Thailand	33.1	0.330	-0.63		Kühlmoorgen-Hille 1977	4
"	"	50.9	0.265	-0.18		"	4
Family RACHYCENTRIDAE							
Rachycentron canadus	Chesapeake Bay Area	134.	0.21	-		Richard 1967	0
"	"	160.	0.20	-		"	0

SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>o</sub>	NOTES	SOURCE	METHOD
Family RAJIDAE							
Raja brachyura ♂	Irish Sea	139.	0.12	-1.52		Holden 1972	0
" " ♀	"	118.	0.19	-0.8		"	0
" clavata ♂	"	85.6	0.21	-0.60		"	0
" " ♀	"	128.	0.09	-1.32		"	0
" eglanteria	lower Delaware Bay, USA	78.5	0.20	-7.78		Fitz & Daiber 1963	4
" erinacea	Long Island Sound, USA	52.1	0.290	-0.83		Richard et al. 1963	4
" "	"	53.5	0.287	-0.91		"	4
" hollandi ♂	East China Sea	57.0	0.521	-		Ishiyama 1951	4
" " ♀	"	62.9	0.408	-		"	4
" " ♂	"	45.5	1.29	-		"	4
" " ♀	"	50.	1.34	-		"	41
" montagui ♂	Irish Sea	68.7	0.19	-0.56		Holden 1972	0
" " ♀	"	72.8	0.18	-0.37		"	0
Family SCARIDAE							
Scarus coeruleus	Virgin Islands, Caribbean	61.5	0.325	-		Randall 1962	31
" guacamaia	"	80.	0.293	-		"	31
" vetula	"	53.4	0.599	-		"	31
Sparisoma chrysopterum	"	41.8	0.782	-		"	31
" rubripinne	"	46.5	0.584	-		"	31

SPECIES NAME	SAMPLING LOCATION	Lo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family SALMONIDAE							
Salmo trutta	Loch Leven 56°12' N 3°22'	40.6	0.276	-		Thorpe 1974	0
"	River Gryfe, Scotland	63.5	0.178	-		Mackay 1970	2
"	Douglas Waters, Scotland	34.7	0.275	-		"	2
"	Northern Sweden, Baltic	98.8	0.219	-1.26		Rosen 1918	00
"	Farver Au (FRG), Baltic	119.	0.108	-2.14		Galhaar 1972	0
"	Rantzau (FRG), North Sea	84.	0.262	-0.92		"	0
"	Vistula, Baltic	142.	0.173	-1.69		Chrzan 1959	00
"	"Stolpe" (?), Baltic	89.1	0.603	-0.34		"	00
"	"Wipper" (?), Baltic	94.9	0.469	-0.56		"	00
"	"Persante" (?), Baltic	101.	0.334	-1.06		"	00
"	Gudenaa River, Denmark, Baltic	88.6	0.388	-0.504		Poulsen 1935	00
"	Dunajec River, Poland, Baltic	94.9	0.493	-0.54		Dixon 1931	00
"	Reda River, Poland, Baltic	95.5	0.344	-0.34		"	00
"	Afva River, Sweden, Baltic	98.7	0.157	-1.90		Alm 1936	00
"	Lake Windermere, England	30.	0.36	-		Frost & Smyly 1952	00
"	Scotland	125.	0.75	-	M = 1.1 (♀)	Nall 1927	00
"	Baltic	156.	0.288	-0.46		Hohendorf 1966	0
"	"	120.	0.439	-0.34		Thurrow 1973	0



SPECIES NAME	SAMPLING LOCATION	Loo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family SALMONIDAE							
Coregonus clupeaformis	Lake Nipigon, Canada	50.	0.13	-	M = 0.17	Hart 1931	00
"	Lake Opeongo, Canada	70.	0.06	-	M < 0.5	Kennedy 1943	00
"	Trout Lake, Wisconsin	44.	0.09	-		Hile & Deason 1934	00
"	Lake Huron	77.0	0.188	-		Van Oosten 1939	2
"	"	79.8	0.182	-		"	2
"	Lake Opeongo, Canada	14.	0.43	-	Dwarf whitefish	Beverton & Holt 1959	0
Coregonus sardinella	Lake Ikroavik, Alaska	38.	0.40	-	M = 0.6	"	0
"	Pereslawski Lake, USSR	35.8	0.318	-		Nikolsky 1957	2
"	Vesijärvi Lake, S. Finland	26.7	0.531	-		Järvi 1947	2
"	Attersee (FRG)	35.8	0.378	-		Eisele 1943	2
"	Lower Lena River, USSR	96.8	0.032	-		Awerizew & Ssytsch-Awerinzewa 1937	4
"	Peene-Achterwasser, Baltic	42.	0.64	-		Thienemann 1935	1
"	Haweswater & Hullswaters, Gr.Br.	41.	0.741	-		Bagenal 1970	0
"	Attersee (FRG)	43.8	0.447	-		Eisele 1943	2
"	Bodensee (Lake Constance)	44.0	0.298	-	YC 29-30	Nüman 1959	2
"	"	39.8	0.417	-	YC 30-31	"	2
"	"	52.8	0.267	-	YC 52-53	"	2
"	"	51.	0.232	-		Surbeck 1916	1
"	Attersee (& other FRG lakes)	48.8	0.383	-		Eisele 1943	2
"	Lena River, upper course	45.4	0.241	-		Nikolsky 1957	2

SPECIES NAME	SAMPLING LOCATION	Lo	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family SALMONIDAE							
Thymallus arcticus	Western Siberia	49.8	0.139	-		Nikolsky 1957	2
"	Baikal Lake (White)	46.8	0.235	-		"	2
"	" (Black)	55.6	0.187	-		"	2
" thymallus	"European waters"	43.2	0.170	-		"	2
"	River Gryfe, Scotland	45.0	0.262	-		Mackay 1970	2
"	Douglas Waters, Scotland	33.9	0.412	-		"	2
" brevirostris	North Western Mongolia	40.	0.288	-		Nikolsky 1957	1
Oncorhynchus nerka	Cultus Lake, Canada	69.	0.58	-		Foerster 1929	00
" keta ♂	Columbia River, Canada	120.	0.27	-	} 4 years } spawners	Marr 1943	00
" " ♀	" "	105.	0.30	-		"	00
" " ♂	" "	106.	0.45	-	} 3 years } spawners	"	00
" " ♀	" "	102.	0.39	-		"	00
Leucichthys artedii	Trout Lake, Wisconsin, USA	29.	0.162	-		Hile 1936	2
"	Silver Lake "	23.	0.488	-		"	2
"	Muskellunge Lake "	21.	0.775	-		"	2
" " ♂	Clear Lake "	41.3	0.417	-		"	2
" " ♀	" "	43.0	0.394	-		"	2
" kiyi	"USA"	28.	0.51	-	M = 0.8	Deason & Hile 1947	00
Cristivomer namaycush	Great Slave Lake, Canada	56.	0.07	-	M = 0.6	Kennedy 1954	00

SPECIES NAME	SAMPLING LOCATION	L <sub>∞</sub>	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family SCIAENIDAE							
<i>Pseudotolithus senegalensis</i> ♂	mouth of Congo River 20 m	42.6	0.947	0.79	Z = 1.1	Troadec 1969	0
" "	" "	50.7	0.607	0.78		"	0
" "	off Lagos, Nigeria 20 m	47.8	0.66	-	Z = 1.25	Longhurst 1964	0
" "	" "	81.	0.33	0.0		Bayagbona 1969	0
" "	off Ghana 20 m	54.	0.44	-		Poinsard & Troadec 1966	0
" typus	off Lagos, 20 m	105.	0.217	-0.26		Anon. 1963	41
" " ♂	mouth of Congo River 10 m	61.1	0.346	-0.36		Poinsard 1973	0
" " ♀	" "	77.4	0.255	-0.41		"	0
" "	off Lagos 20 m	103.	0.29	-0.15		Bayagbona 1969	0
" "	Nigeria	61.2	0.37	-	Z = 1.25	Longhurst 1963	0
" "	Congo River mouth 20 m	61.3	0.42	-0.24		Poinsard & Troadec 1966	4
Nibea nibe							
" "	Formosa Strait	58.5	0.116	-		Lin 1956	00
" "	27° N 122° E	58.7	0.145	-1.04		Matsui & Amio 1951	4
" "	27° N 123° E 60 m	37.1	0.354	-2.0		Matsui & Amio 1951	41
" "	" "	37.1	0.523	-0.01		Matsui & Takai 1951	4
" "	27° N 122° E 60 m	57.3	0.177	-1.09		"	4
" "	32° N 127° E 60 m	50.9	0.227	-1.08		"	4
" ♂	-	40.1	0.238	-0.89		"	4
" ♀	-	45.0	0.225	-1.05		"	4
" "	32° N 127° E 60 m	46.9	0.297	-1.09		Matsui & Amio 1951	4

SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family SCIAENIDAE							
Cheilotrema saturnum	Baja California	47.	0.183	-2.64		Limbaugh 1961	41
Lagodon rhomboides	30°25' N 87°07' W	19.5	1.019	-0.537		Hansen 1970	4
Leiostomus xanthurus	Texas	34.	0.43	-		In: Rao 1966	0
"	Lake Pontchartrain, Louisiana	23.9	0.89	-0.04		Sundaraj 1960	4
"	Chesapeake Bay (York River Mouth)	24.1	0.725	-1.13		Pacheco 1962	4
Macrodon ancylodon ♂	31° S, Brazil	33.0	0.44	-		Yamaguti & dos Santos 1966	0
" ♀	31° S, "	43.6	0.27	-		"	0
Menticirrhus undulatus	Southern California	44.4	0.385	-		Joseph 1962	00
Micropogon furnieri	28° S 49° W	63.3	0.12	0.37		Wootton Valozzer 1962	0
" undulatus	Texas	39.	0.35	-		In: Rao 1966	0
"	Florida, Pensacola Estuary	18.2	1.452	0.18		Hansen 1970	41
Nibea argentea	East China Sea, "Urride" 30 m	69.6	0.138	-0.10		Yasuda & Kosaka 1950	4
Pseudosciaena polyactis	37.5°N 125.5° E, South Korea	46.0	0.163	-2.82		Woo II Choo 1970	0
Pseudotolithus elongatus	Congo River Estuary	46.7	0.274	-0.67	M = 0.34	Le Guen 1971	0
"	off Sierra Leone 20 m	51.8	0.257	-0.44	Z = 1.49	Longhurst 1963	00
Roncador stearnsi	Southern California	47.6	0.287	-		Joseph 1962	00
Sciaena aquila	Port Said, Egypt	70.	0.195	-		Rafail 1971	41
" ruselli	13°10' N 100°45' E, Thailand	18.5	1.012	-		Phetthongkam & Thasananul- kit 1972	4

SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family SCIAENIDAE							
<i>Cynoscion leiarchus</i>	Off Caera State, Brazil	45.4	0.35	-0.57		Namura 1966	0
" <i>macdonaldi</i>	mouth of Colorado River, lower California	128.	0.3	-	M = 0.3	Berdegue 1955	00
"	Gulf of California	208.	0.255	-		Nakashima 1916	41
" <i>maracaiboensis</i>	Lake Macaraibo, Venezuela (a lagoon)	106.	0.27	0.21		de Espinosa 1972	0
" <i>nobilis</i>	33°30' N 119°30' W	146.	0.128	-0.28	M = 0.3	Thomas 1968	0
" <i>petranus</i> ♂	24°30' S, Southern Brazil	30.3	0.342	-	} M = 0.58	dos Santos 1963	0
" " ♀	" "	32.1	0.345	-		"	0
" <i>nebulosus</i> ♂	North West Florida	57.0	0.222	-		Klima & Tabb 1969	4
" " ♀	" "	70.6	0.174	-		"	4
"	Punta Gorda, Florida 10 m	55.8	0.361	-		Welsh & Broder 1964	4
"	Corpus Christi, Texas	85.5	0.134	-0.74		Pearson 1929	4
"	Fort Myers, Florida	70.1	0.169	-0.44		Moffett 1961	4
"	Cedar Keys, "	60.6	0.274	-0.02		"	1
"	" "	60.4	0.223	-0.289		"	4
"	Indian River, Florida	61.8	0.190	-0.14		Tabb 1961	41
"	" "	79.4	0.147	-0.47		"	41

SPECIES NAME	SAMPLING LOCATION	L <sub>∞</sub>	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family SCIAENIDAE							
<i>Michthys imbricatus</i>	East China Sea	80.6	0.32	-		Hanabuchi 1967	0
<i>Otolithoides brunneus</i>	off Bombay, India	150.	0.212	-		Narayanan Kutty 1961	1
<i>Pogonias chromis</i>	Texas	105.	0.17	-		Richard 1972	00
<i>Pseudosciaena coibor</i>	Chilka Lake, India	75.	0.386	-0.08		Rajan 1964	41
" <i>diacanthus</i>	Bombay-Saurashtra, India	122.	0.315	-0.31	M = 0.8	Rao 1966	0
"	off Bombay, India	122.	0.294	-		"	4
<i>Sciaenops ocellatus</i>	Texas	90.	0.416	-0.08		Miles 1950	41
"	"	105.	0.30	-		Pearson 1929	00
<i>Umbrina canariensis</i>	off Morocco 100 m	57.2	0.259	-		Dardignac 1961	4
" <i>cirrrosa</i>	" 50 m	86.5	0.279	-		"	4
<i>Umbrina fusca</i>	" 50 m	77.1	0.393	-		"	4
" <i>cirrrosa</i>	Port Said, Egypt	30.	1.452	-		Rafail 1971	4
Family SCOMBERESOCIDAE							
<i>Cololabis saira</i>	Southern California to Vancouver	37.6	0.34	-1.19	spring spawn. M ≈ 1.6	Hughes 1974	0
"	"	37.8	0.38	-0.72	fall spawn.	"	0
"	Northern California	39.4	0.595	-1.56		Sunada 1974	4

SPECIES NAME		SAMPLING LOCATION		L <sub>∞</sub>	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family SCOMBRIDAE									
Rastrelliger kanagurta		off Cochin, India		21.8	5.16	-		George & Banerji 1964	0
"	"	off Calicut, India		23.3	3.12	-		"	0
"	"	off Karwar, India		22.4	4.32	-		"	0
"	"	Java Sea		23.9	2.76	0.08	M = 4.44	Sudjastani 1973	0
"	"	Kerala to Maharashtra, India		23.9	4.92	0.17	M = 7.8	Banerji 1973	0
"	"	Al Ghardaqa, Red Sea, Egypt		42.0	0.288	0.94		Rafail 1972a	0
"	"	Indian waters		31.3	0.64	0.06		Seshappa 1958	4
"	"	"		31.6	0.60	-		Sekaran <u>et al</u> 1940	00
"	neglectus	10° N 100° E Gulf of Thailand		20.	3.528	-	January brood	Sacondharman <u>et al</u> , 1970	0
"	"	"		19.6	4.14	-	July brood	"	0
"	"	Tanjung Satai (Western Borneo)		22.9	2.28	0.01	M = 4.56	Sudjastani 1973	0
"	"	Inner Gulf of Thailand		20.9	3.38	0.002	M = 7.22	Mongskul 1974	0
"	"	"		20.9	4.2	-	M = 4.0	Somjaiwong <u>et al</u> , 1972	0
Scomberomorus cavalla ♂		off Caera State, Brazil		116.	0.18	-0.22	LF	Momura 1967	0
"	♀	"		137.	0.15	-0.13	LF	"	0
"	maculatus ♂	Florida, Coastal waters		61.2	0.373	-	LF	Klima 1959	4
"	♀	"		76.7	0.356	-	LF	"	4
"	♂	off Caera State, Brazil		83.	0.20	-0.40	LF	Momura 1967	0
"	"	"		94.4	0.20	-0.05	LF	"	0

SPECIES NAME	SAMPLING LOCATION	Loo	K	t <sub>o</sub>	NOTES	SOURCE	METHOD
Family SCOMBRIDAE							
Scomber scombrus	English Channel & Irish Sea	41.8	0.43	-		Steven 1952	00
"	Mediterranean (Western)	35.	0.5	-		Andreu & Rodriguez-Roda 1953	00
"	North East Atlantic	41.	0.44	-		Le Gall 1939	00
"	New England	46.1	0.234	-2.11		ICNAF Redbook 1973	4
"	ICNAF Subarea 4	44.5	0.269	-1.57		Stobo & Hunt 1974	4
"	Central North Sea	33.6	0.837	-0.18		Nedelec 1958	4
"	Southern North Sea	42.	0.238	-2.07		"	4
"	Celtic Sea	37.4	0.47	-0.97		"	4
"	English Channel	37.	0.56	-		Steven & Corbin 1939	00
"	New England	47.3	0.262	-1.34		Isakov 1973	4
"	off Newfoundland	42.9	0.36	-1.14		Moore et al 1975	0
"	North Sea	39.	0.43	-		Ehrenbaum 1923	00
Pneumatophorus japonicus	Sea of Marmara & Bosphorus	33.	0.473	-		Taggac 1957	41
"	Japanese Waters	46.4	0.40	-		Kondo 1966	00
"	Soviet Pacific Waters	60.	0.306	-		Berg et al 1949	41
"	Red Sea, Egyptian Waters	27.9	0.485	-0.03		Rafail 1972 a	0
"	North West Pacific	45.5	0.34	-		Japanese Fish.Agency 1952	00
"	Central Japan Sea	44.0	0.441	-0.42		in: Nikolsky 1957	4
"	California	40.	0.4	-	M = 0.9	Fitch 1951, 1956	00



SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family SCORPAENIDAE							
<i>Sebastes inermis</i>	Senday Bay, Japan, Zostera Belt	29.6	0.250	-0.76		Hatanaka & Sekino 1962	4
" <i>pachycephalus</i>	Amakusa Island, Japan	25.	0.372	-		Shiokawa 1962	0
" <i>marinus</i>	Barents Sea	73.2	0.064	-0.11		Travin 1951	4
" " ♂	Gulf of Maine 42.7° N 70.2° W	35.2	0.145	-		Kelly & Wolf 1959	4
" " ♀	" "	46.2	0.082	-		"	4
" "	Browns Bank	40.7	0.067	-0.82		Pearlmutter & Clarke 1949	4
" "	Barents Sea	75.6	0.059	-		Veschezerov 1944	4
" <i>mentella</i>	" >300 m	49.2	0.100	-0.22		Travin 1951	4
" " ♂	Gulf of Maine	32.7	0.151	-		Elvertowsky 1964	4
" " ♀	"	52.5	0.058	-		"	4
<i>Scorpaenichthys marmoratus</i> ♂	California	57.7	0.342	-		O'Connel 1953	4
<i>Scorpaena guttata</i>	Santa Monica, California 30 m	34.6	0.389	-		Turner et al 1969	3
<i>Sebasticus marmoratus</i>	Amakusa Islands	29.2	0.335	-0.06		Shiokawa 1962	0
<i>Sebastes alutus</i>	Bering Sea, 4° C spawn-temp.	45.3	0.124	-		In: Major & Shippen 1970	4
" " ♂	off Washington State	40.3	0.20	-0.71	} M = 0.10	"	0
" " ♀	" "	43.2	0.18	-		"	0
" <i>mystinus</i>	Monterey Bay, California	38.7	0.168	-0.60		Miller & Geiber 1973	4
<i>Helicolenus dactylopterus</i>	Morocco 33° S 8° W 200 m	42.1	0.204	-0.38		Boutière 1958	4
<i>Gymnapistes marmoratus</i>	South Tasmania, Zostera Belt 43° S 21.4	21.4	0.117	-2.3		Grant 1972	0
<i>Hypodytes rubribinnis</i>	Tamioka Bay, Japan, Zostera Belt	9.6	0.548	-0.65		Kikuchi 1966	4
<i>Helicolenus dactylopterus</i>	off Algier, Mediterranean	21.2	0.210	-0.85		Siblot-Boutefika 1976	0
<i>Scorpaena porcus</i>	" "	29.4	0.162	-0.97		Siblot-Boutefika 1976	0

SPECIES NAME	SAMPLING LOCATION	Loo	K	t <sub>o</sub>	NOTES	SOURCE	METHOD
Family SCYLLIORHINIDAE							
Scyllorhinus canicula	Yugoslavia 100-200 m	56.8	0.530	-		Zupanovič 1961	5
Family SERRANIDAE							
Epinephelus aeneus ♂	off Morocco, 50 m	85.	0.247	-		Cadenat 1935	41
" " ♀	"	100.	0.226	-		"	1
" striatus	Virgin Islands, Caribbean	110.	0.224	-		Randall 1962	31
Lateolabrax japonicus	off Sanriku, Japan (Pacific)	78.	0.225	-0.76		Yasuda & Koike 1950	4
" "	Japan Inland Sea	125.	0.142	-0.27		"	4
" "	Matsushima Bay, Japan	85.	0.193	-0.66		Hanataka & Sekino 1962	4
Dicentrarchus labrax	Casablanca to Agadir, Morocco	105.	0.089	-0.56		Gravier 1961	41
" " ♂	off Sète, France, Mediterranean	62.5	0.282	-0.16		Barnabé 1973	4
" " ♀	"	78.5	0.228	-0.05		"	4
" " ♂	off Southern Ireland	70.4	0.14	-0.03	M = 0.10	Holden & William 1974	0
" " ♀	"	72.3	0.14	-0.20		"	0
Mycteroperca microlepis	Tampa Bay, Florida	124.	0.124	-		McErlan 1963	4
Serranus alexandrinus	West of Alexandria, Egypt	72.5	0.154	-0.73		Rafail 1972 b	0

SPECIES NAME	SAMPLING LOCATION	Lo	K	t <sub>o</sub>	NOTES	SOURCE	METHOD
Family SERRANIDAE							
<i>Anthias squamipinnis</i>	Eilat, Red Sea	16.8	0.602	-		Gundermann & Popper 1975	4
<i>Diplecctum formosum</i>	Florida	31.	0.287	-		Bortone 1971	4
<i>Döderlinia berycoides</i>	35°10' N 132° E 100 m	30.8	0.351	-		Kibesaki 1950	41
<i>Morone americana</i> ♂	Chesapeake Bay, York River Mouth	35.	0.131	-0.98		St. Pierre & Davis 1972	4
" " ♀	"	52.	0.073	-1.52		"	4
" punctata	Port Said, Egypt	40.	0.156	-		Rafail 1971	4
"	Casablanca to Agadir, Morocco	103.	0.066	-1.316		Gravier 1961	4
<i>Paralabrax clathrus</i>	Southern California	64.0	0.103	-0.77		Young 1963	4
<i>Roccus lineatus</i> ♂	San Francisco Bay	76.3	0.246	-		Scofield 1931	4
" " ♀	"	123.	0.125	-		"	4
" saxatilis	Coos Bay, Oregon	95.8	0.188	-		Morgan & Gerlach 1950	4
"	Chesapeake Bay	98.7	0.158	-2.96		Vladykov & Wallace 1952	41

SPECIES NAME	SAMPLING LOCATION	Lo	K	t <sub>o</sub>	NOTES	SOURCE	METHOD
Family SIGANIDAE							
Siganus canaliculatus	Southern Negros, Philippines	25.2	1.872	0.02		Lam 1974	41
"	Aquaria	16.1	3.3	-0.09		von Westernhagen & Rosenthal 1976	4
"	Sea Cages, Mactan Islands, Phil.	31.6	1.978	-0.15		Horstmann 1975	4
"	Aquaria	30.8	1.548	-0.08		Tsuda et al. 1974	4
"	Sea Cages, Mactan Islands, Phil.	32.7	1.659	-0.22		Horstmann 1975	4
"	Aquaria	16.3	3.3	-0.07		v. Westernhagen & Rosenthal 1975	4
"	Israel Coast, Mediterranean	42.	1.016	-		Ben Tuvia et al. 1973	41
"	Sea Cages, Mactan Islands, Phil.	24.4	2.321	-0.21		Horstmann 1975	4
"	Ponds, Fidji Islands	31.	1.90	-		Lichatowich & Popper 1975	4
Family SILLAGINIDAE							
Sillago panius	Hoogly Estuary, Bay of Bengal	153.	0.054	+0.27	bad fit, L <sub>∞</sub> too high	Krishnaya 1963	0
"	New South Wales, Australia 10 m	36.5	0.677	-		Cleland 1947	4
"	Gulf of Manaar, India	28.1	0.75	-		Radhakrishnan 1957	00
Family SILURIDAE							
Silurus glanis	Aral Lake	296.	0.042	-	p. 334	Nikolsky 1957	2

SPECIES NAME	SAMPLING LOCATION	LOO	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family SPARIDAE							
<i>Acanthopagrus bifasciatus</i>	Gulf of Aden 50 m	48.1	0.181	-0.93		Druzhinin 1975	4
<i>Argyrops spinifer</i>	" 50 m	96.4	0.077	-1.43		"	4
" <i>filamentosus</i>	" 50 m	45.6	0.227	-0.291		"	4
<i>Cheimerus nufar</i>	" 50 m	100.	0.067	-2.96		"	4
<i>Sparus swinhonis</i>	Tokyo Bay ? Japan	33.4	0.559	-0.06		Oshima 1942	4
<i>Taius tumifrons</i>	Hong Kong	41.5	0.249	-0.30		Li Kwang Ming 1958	4
" "	East China & Yellow Sea	93.2	0.069	-1.18		Tanaka 1956	4
" "	East China Sea, "A"	47.4	0.249	-0.30		Yasuda 1950	4
" "	East China Sea, "B"	34.3	0.373	-0.37		"	4
<i>Dentex congoensis</i>	Cape Verde, West Africa 100 m	30.3	0.16	-0.81		"	0
" <i>macrophthalmus</i>	Cap Blanc, West Africa	37.5	0.162	-1.32		"	0
" "	Cape Verde, West Africa	39.0	0.18	-0.88		"	0
" <i>maroccanus</i>	Cap Blanc, West Africa	34.3	0.18	-0.49		"	0
" "	Cape Verde, West Africa	32.5	0.18	-0.62		"	0
<i>Dentex polli</i>	Cap Blanc, West Africa	46.4	0.22	-0.28		"	0
<i>Pagrus ehrenbergi</i>	off Ghana	44.8	0.49	-		Rivajec 1973	0
" "	off Togo	41.2	0.501	-0.18		Beck 1974	4
<i>Polysteganus undulosus</i>	Cape to Delagoa Bay, South Africa	98.4	0.181	-0.74		Ahrens 1966	4
<i>Dentex canariensis</i>	off Ghana, 50 m	70.	0.173	-0.136		Rivajec 1973	41
" <i>filosus</i>	Cap Blanc, West Africa	107.	0.124	-0.60			0

SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family SPARIDAE							
Chrysophrys aurata	43°25' N 3°40' E, near Sète	59.1	0.275	-0.26		Audoin 1962	4
"	off Sète (France), Mediterr.	53.9	0.26	-0.74		Lassère & Labour 1974	0
"	"	76.	0.13	-1.22	Z = 1.17	Lassère 1976	0
"	Hauraki Gulf, New Zealand	40.1	0.288	-0.05		Cassie 1956	4
" major	Wakasa Bay, Japan	70.8	0.117	-		Akazaki 1960	4
"	Seto Inland Sea, Japan	55.3	0.18	-0.47		Kuniyuki et al. 1975	0
"	near Amakusa, Japan	71.	0.178	-		Mukarami & Shindo 1950	4
"	32° N 125° E	84.3	0.100	-		Mukarami & Okado 1967	0
Pagellus acarne	20° N West Africa 120 m	36.	0.23	0.97	Z = 1.15	Le Trong Phan & Kompowski 1972	0
" centrodonatus ♂	Bay of Biscay	48.1	0.162	-0.76		Guéguin 1969	4
" ♀	"	49.9	0.146	-0.93		"	4
"	French Mediterranean Coast	61.2	0.091	-		Olivier 1928	4
" coupei	off Dakar, Senegal	39.4	0.180	-0.81	Z = 1.58	Le Trong Phan & Kompowski 1972	0
"	off Rio de Oro, West Africa	40.2	0.19	-0.63		"	0
" natalensis	Gulf of Aden 50 m	45.5	0.20	-1.01		Druzhinin 1975	4

SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family SPHYRAENIDAE							
Sphyraena argentea	off Los Angeles	140.	0.139	-1.58		Walford 1932	41
" barracuda ♂	Florida	178.	0.087	-		De Sylva 1963	4
" " ♀	"	156.	0.113	-		"	4
Family SQUALIDAE							
Squalus acanthias	Sea of Japan	132.	0.074	-		Kaganovskaya 1933	41
" " ♂	Washington State, USA	102.	0.071	-5.2		Bonham et al. 1949	00
" " ♀	"	153.	0.036	-6.7		"	00
" " ♂	Hecate Strait, British Columbia	84.7	0.092	-3.7		Ketchen 1975	0
" " ♀	"	125.	0.031	-10.6		"	0
" " ♂	Georgia Strait,	96.1	0.067	-5.0		"	0
" " ♀	"	129.	0.034	-7.3		"	0
" " ♂	Northern Scotland	79.7	0.21	-2.0		Holden & Meadow 1962	0
" " ♀	"	101.	0.11	-3.6		"	0
Family STICHAEIDAE							
Lumpenus lampretiformis	Firth of Forth Area, Scotland	47.6	0.205	-		Fulton 1904	4
Ulvaria subbifurcata ♂	Trinity Bay, Newfoundland	17.0	0.186	-0.45		LeDrew & Green 1975	4
" " ♀	"	14.9	0.215	-		"	4

SPECIES NAME	SAMPLING LOCATION	Loo	K	t <sub>o</sub>	NOTES	SOURCE	METHOD
Family SYNGNATHIDAE							
Hippocampus zosterae	Cedar Key, Florida	2.0	12.	-	M = 2.5	Strawn 1958	00
" hudsonius	Florida	14.0	2.5	-	M = 1.0	Herald & Rakowicz 1951	00
Syngnathus typhle	Kiel Bight, Baltic	26.2	0.558	-		Worthmann 1975	51
Syngnathus rostellatus	"	20.0	0.747	-		"	51
Family SYNODONTIDAE							
Saurida tumbil	Mouth of Manilla Bay	43.6	0.431	-		Tiews et al. 1972	5
" "	28° N 124° E East China Sea	63.2	0.286	-0.289	M = 0.46	Shindo 1972	41
" undosquamis	Inner Gulf of Thailand, off Trat	37.9	0.889	-		Kühlmorgan-Hille 1970	5
" "	Gulf of Thailand, off Prachuap Khirikhan	41.3	1.132	-		"	5
" "	Mediterranean, Israel Coast	31.5	0.698	-		Ben Yami & Glaser 1974	41
Family TETRAODONTIDAE							
Uranostoma richiei	Lyttelton Harbour, New Zealand	21.9	0.246	-		Habib 1976	4



SPECIES NAME	SAMPLING LOCATION	Lco	K	t <sub>0</sub>	NOTES	SOURCE	METHOD	
Family THUNNIDAE (LF)								
Thunnus albacares	off Mexico 19,5° N 111° E	YC 1953	172.	0.56	-	Hennemuth 1961	0	
"	"	YC 1954	162.	0.66	-	"	0	
"	"	YC 1955	173.	0.66	-	"	0	
"	Eastern Atlantic		195.	0.42	-	M = 0.8	Le Guen & Sagakawa 1973	0
"	Japanese Waters		169.	0.564	-		Yabuta & Yakinawa 1957	4
"	Japanese Waters		191.	0.327	-1.02		Yabuta et al. 1969	4
"	Hawaii		190.	0.454	-0.75		Moore 1951	4
"	Eastern Atlantic		159.	0.160	-0.47		LeGall 1951	4
"	off Dakar, trop. Atlantic		182.	0.431	-		LeGuen et al. 1969	0
"	"		207.	0.315	-		LeGuen & Champagnac 1968	0
"	off Pointe Noire, Congo		166.	0.557	-		"	0
"	"		182.	0.443	-		LeGuen et al. 1969	0
♂	off Mauritania		167.	0.183	-		Vincent Guaz 1969	4
♀	"		228.	0.116	-		"	4
"	-		152.	0.661	-0.62		Yakinawa 1959	4
"	off Senegal		187.	0.152	-0.14		Postel 1955a	4
"	- (data: Zharov)		229.	0.277	-		int. Naudin Laperouse 1968	4
"	- (data: Postel)		205.	0.112	-4.12		"	4

SPECIES NAME	SAMPLING LOCATION	Lat.	K	t <sub>0</sub>	ROUTS	SOURCE	METHOD
Family THUNNIDAE (LP)							
Thunnus alalunga	Gulf of Biscay 45°30' N 10° W	135.	0.138	-		Alonge & Delaporte 1974	4
"	" 50° N 10° W	150.	0.119	-		"	4
"	South African Waters	151.	0.118	-		Talbot & Penrith 1963	4
"	Tropical Atlantic	135.	0.190	-		Yang 1970	00
"	Bay of Biscay	134.	0.183	-	M = 0.2	Bard 1971	4
"	off Northern Spain	104.	0.274	-		Anon. 1973	0
"	Eastern Atlantic (off Brittany?)	118.	0.262	-1.31		Friol 1944	4
"	off California	109.	0.233	-2.14		Bell 1962	4
"	Central North Pacific	189.	0.093	-		Otsu 1960	4
"	North Pacific 40° N 150° W	136.	0.169	-1.87		Clemens 1961	4
"	North Pac. Current 28-40° N	145.	0.159	-3.06		Nose <u>et al.</u> 1957	4
"	Central Atlantic	140.	0.141	-	M = 0.23	Beardsley 1971	0
Sarda sarda	off Morocco (Atlantic)	64.0	0.693	-1.42		Dardignac 1962	4
"	Black Sea & Eastern Mediterran.	81.5	0.525	-		Nikolsky 1957	4
"	"	67.8	0.795	-		Mümann 1955	4
"	"	103.	0.132	-1.80		Zusser 1954	4
"	"	95.6	0.237	-1.24		Nikolov 1960	4
"	"	64.0	0.860	-		Türgan 1958	4

SPECIES NAME	SAMPLING LOCATION	Loc	K	t <sub>0</sub>	NOTES	SOURCE	METHOD
Family THUNNIDAE (LF)							
Auxis thazard	South West of Sri Lanka	61.	0.83	-		Sivasubramaniam 1973	5
Euthynnus affinis	Seychelles	90.	0.444	-		Wheeler & Ommanney 1953	41
" alletteratus	Tunisia	136.	0.164	-		Postel 1956	4
" "	Senegal	99.5	0.315	-		Postel 1955	4
Thunnus atlanticus	Cuba	78.	0.33	-	M = 0.67	Charles Martin 1975	0
" "	Florida	90.	0.216	-1.50		Idyll & Sylva 1963	41
" maccoyi	38° S 155° E, Australia	222.	0.141	-		Robins 1963	0
" "	Australia	220.	0.14	-		Yukinawa 1970	0
" "	Australia	220.	0.151	-	M = 0.2	Hynd & Lucas 1974	0
" orientalis	Suruga Bay, Japan	266.	0.075	-1.01		Kimura 1932	4
" tonggol	Northern Australia	110.	0.324	-		Serventy 1949	4
Katsuwonus pelamis	Hawaiian waters	84.6	1.16	-		Uchiyama & Struhsaker 1975	4
" "	" "	85.1	0.95	-		Brock 1954	4
" "	Eastern Pacific	85.1	0.44	-0.01		Schaefer 1961	4
" "	Hawaiian waters	82.3	0.77	-		Rothschild 1967	0
" "	10° N 100° W (Pacific)	80.5	0.63	-	M = 1.68	Joseph & Calkins 1969	0
" "	North of Madagascar 12° S 48° E	62.3	0.98	-		Marcille & Stequert 1976	0

SPECIES NAME	SAMPLING LOCATION	LOO	K	t <sub>O</sub>	NOTES	SOURCE	METHOD
Family THUNNIDAE (LF)							
Thynnus thynnus	European Waters	389.	0.067	-1.60		Sella 1930	4
"	Bay of Biscay	419.	0.056	-1.92		Anon. 1954	4
"	Long Island, New York	318.	0.091	-0.308		Westmann & Neville 1942	4
"	Spain, off Barbate	384.	0.064	-2.52		Rodriguez-Roda 1972	4
"	Long Island, New York	218.	0.162	-1.12		Westmann & Gilbert 1941	4
"	Cape Cod - Long Island	468.	0.050	-	M ≈ 0.15	Mather & Schuck 1960	4
" obesus	10° S 115° W	187.	0.38	-		Kume & Joseph 1966	0
"	Pacific	236.	0.125	-	tagging data in table III	Alverson & Peterson 1963	31
"	North Pacific Current 28-40° N	195.	0.106	-1.13		Nose et al, 1957	4
"	Central & Western Pacific	222.	0.183	-		Iversen 1955	3
"	Eastern Pacific	169.	0.508	-		Hennemuth 1961	4
" ♂	Central Pac. (Hawaiian Longlines)	222	0.114	-		Shomura & Keala 1963	0
" ♀	"	196.	0.167	-		"	0
Family TORPEDINIDAE							
Torpedo torpedo	Gulf of Tunis	43.1	0.110	-2.58		Quignard 1973	4
Family TRIACANTHIDAE							
Triacanthus brevirostris	Ise & Mikawa Bay, Japan	41.4	0.216	-		Ohsima & Nakamura 1941	4

SPECIES NAME	SAMPLING LOCATION	Loo	K	t <sub>o</sub>	NOTES	SOURCE	METHOD
Family TRICHIURIDAE							
<i>Trichiurus lepturus</i>	Madras to Mangalore, India	98.	0.200	-		Prabhu 1955	00
"	Cape Blanc - Cape Verde, W.Africa	147.	0.296	-0.46		Wojciechowski 1971	4
"	Yellow Sea	50.5	0.286	-		Hamada 1971	4
"	-	45.4	0.411	-		in: Hamada 1971	4
"	Suruga Bay, Japan	43.4	0.298	-		Kosaka <u>et al</u> 1963	2
"	Gulf of Manaar Area, India			-		James 1967	4
Family TRIGLIDAE							
<i>Trigla hirundo</i>	off Morocco 50-120 m	65.	0.148	-1.11		Collignon 1968	4
<i>Chelidonichthys spinosus</i>	Japan	54.8	0.494	-		Yukonawa 1951	00
" kumu	New Zealand, Pegasus Bay 45° S 173° E	54.9	0.406	0.29		Staples 1972	0
" ♂	Hauraki Gulf, N.Zealand 36,50° S 175° E	28.8	0.549	-0.55		Elder 1976	0
" ♀	" "	36.4	0.641	0.19		"	0
Family TRYGONIDAE							
<i>Dasyatis satina</i> ♂	Texas Coast, USA	109.	0.309	-0.90		Sage <u>et al</u> 1972	51
" ♀	"	144.	0.259	-0.72		"	5
" akajei ♂	Miyazaki Prefecture, Japan	105.	0.1	-	M = 1.8 (?)	Yokota 1951	00
" ♀	" "	150.	0.1	-	M = 0.45	"	00
<i>Urolophus halleri</i>	Belmont-Newport, Calif. 10 m	46.0	0.152	-2.03		Babel 1967	4
Family XIPHIDAE							
<i>Xiphias gladius</i>	Canada Gulf stream waters (W∞ = 630 kg → LF, with bill)	630.	0.153	-	catch temp. 18° C	Beckett 1974	41

SPECIES NAME	SAMPLING LOCATION	Loo	K	t <sub>o</sub>	NOTES	SOURCE	METHOD
Family ZEIDAE							
Zeus faber	Mauritania 20° N	57.9	0.298	-		Debrosse 1937	41
"	Bay of Biscay to Channel	56.7	0.267	-	♂♂ & ♀♀	"	41
"	"	69.3	0.147	-		"	41
Family ZOARCIDAE							
Macrozoarces americanus	Southern New England	122.	0.107	-		Olsen & Merriman 1946	4
"	Bay of Fundy	91.	0.076	-0.53		Clemens & Clemens 1921	4

### 2.3. Controversial data, Order Selachoidae

Order SELACHOIDEI (due to the method utilized (by Holden, 1974), the data given here seem particularly unreliable and need confirmation).

Family	Hexanchidae	L (co)	"K"	Family	Carcharhinidae** (cont'd)	L (co)	"K"
	Heptanchia perlo	214	0.121		Carcharias taurus	318	0.193
	Hexanchus griseus	482	0.125*		Carcharias maculipinnis	244	0.137
Family	Orectolobidae				Eulemia milberti	239	0.133
	Ginglymostoma cirratum	427	0.141		Galeocerdo cuvier	457	0.106
Family	Lamnidae**				Galeorhinus zyopterus**	192	0.104
	Alopias superciliosus	549	0.168		Mustelus canis**	152	0.138*
	Alopias vulpinus	610	0.127*		Negaprion brevirostris	284	0.132
	Aprionodon isodon	160	0.168		Prionace glauca**	383	0.110
	Carcharodon carcharias	1 113	0.113		Scoliodon tetrarhynchus	93	0.226*
	Cetorhinus maximus	1 372	0.131*		Tiakia barbouri	38	0.210
Family	Scyliorhinidae**			Family	Sphyrnidae		
	Isurus oxyrinchus	610	0.143		Sphyrna diplana	305	0.150
	Lamna nasus**	305	0.112		Sphyrna tiburo	152	0.110
	Pseudotriakis microdon	295	0.170		Sphyrna tudes	457	0.166
Family	Carcharhinidae**				Sphyrna zygaena	656	0.158
	Carcharhinus acronotus	168*	0.159*	Family	Squalidae**		
	Carcharhinus leucas	305	0.125		Centroscyllium coelestis	112	0.115
	Carcharhinus limbatus	228*	0.139		Dalatias licha	182	0.180
	Carcharhinus longimanus	396	0.186		Etmopterus hillianus	32	0.170
	Carcharhinus obscurus	335	0.171		Etmopterus spinax	52	0.144
	Carcharhinus oxyrinchus	152	0.148				

all data from Holden (1974); \* mean of two values; Lmax  $\pm$  L (co);

\*\* families, genera or species listed in previous tables.

#### 2.4. Bibliography II

As explained in Part I, I have not seen all of the papers listed here, and their reference had to be taken verbatim from various author's reference list. Several of these references appear to be at least incomplete. I felt, however, that nothing would have been gained by not including them, together with the corresponding growth parameters. It is hoped that an improved version of this compilation can be produced soon in which the references have been revised.



- AASEN, O. 1963: Length and Growth of the Porbeagle (Lamna nasus, Bonnaterre) in the North West Atlantic. Reports on Norwegian Fishery and Marine Investigations 13 (6): 20-37.
- ABLE, K. and J. MUSICK 1976: Life history, ecology and behavior of Liparis inquilinus (Pisces, Cyclopteridae) associated with the sea scallop, Placopecten magellanicus. Fish. Bull., U.S., 74 (2): 409-421.
- ADAMS, L. A. 1942: Age determination and rate of growth of Polyodon spathula, by means of growth rings of the otolith and dentary bone. Am. Midland Naturalist, 28 (3): 617-630.
- AGUAYO, M. 1971: Determinacion de la edad en la Merluza (Merluccius gayi Guichenot 1848) mediante la lectura de otolitos. Inst. Fom. Pesq. Bol. Científico, (16): 1-49.
- AHRENS, R. 1966: Age determination of the Seventy-four. South African Assoc. for Marine Biological Research, Bull., (6): 44-47.
- AKAZAKI, M. 1960: Growth and age of the sea bream, Chrisophrys major, caught in Wakasa Bay. Bull. Jap. Soc. sci. Fish., 26 (3): 216-222.
- AKYÜZ, E. 1959: Unpublished data used by Beverton & Holt (1959).
- ALEEV, Y. 1956: [Vop. Ikhtiol.] Journal of Ichthyology 6: 75-95.
- ALEGRIA, J. and M. FERREIRA DE MENEZES, 1970: Edad y crecimiento del ariacó Lutjanus synagris (Linnaeus) en el Nordeste del Brasil. Arq. Cién. Mar. 10 (1): 65-68.

- ALLEN, G.R. 1975: Anemonefishes. T.F.H. Publ., 2nd. ed., Jersey City, 351 pp.
- ALM, G. 1936: Havslaxöringen i Afvaan Stockholms Sportfiskeklubb. Arsbok, Universitet Bergen.
- ALM, G. 1952: Rep. Inst. Freshw. Drottning. 33, 17 p.
- ALONCLE, H. and F. DELAPORTE 1974: Données nouvelles sur le germon atlantique Thunnus alalunga Bonnaterre 1788 dans le nordest Atlantique 2ième Partie. Rev. Trav. Inst. Pêch. marit. 38 (1): 5-102.
- ALVERSON, F.G. and C.L. PETERSON 1962: Synopsis of biological data on bigeye tuna Parathunnus sibi (Temminck and Schlegel) 1844. FAO Fish. Rep., (6), vol. 2: 482-514.
- ANANIADIS, C.I. 1952: Quelques Considerations biométriques sur l'allache (Sardinella aurita C.V.). Prakt. Hellen. Hydrobiol. Inst., 5 (1): 45 p.
- ANDERSEN, K. 1938: An investigation into the alterations in the growth-rate of the haddock. I.C.E.S. Rapp. & Proc. Verb. 108 (1): 85-87.
- ANCELLIN, J. 1954: Observations sur la morue de Terre-Neuve et du Labrador. Cons. Int. Explor. Mer, Rapp. & P.-V. Réun., 136 (13): 72-76.
- ANDREU, B. and J. RODRIGUEZ-RODA 1953: Contribucion a la biologia de la caballa Scomber scombrus, del Levante Espanol (Mediterranea Occidental). Publ. Inst. Biol. Apl. Barcelona, 12: 129-185.

- ANDREU, B., J. RODRIGUEZ-RODA and M.G. LARRANETA 1950:  
Contribución al estudio de la talla, edad y crecimiento de la sardina (Sardina pilchardus Walb.) de las costas españolas de Levante (Noviembre 1949-Mayo 1950). Publ. Inst. Biol. Apl. Barcelona, (7): 159-189.
- ANON. 1954: Recherches sur les poissons / Thon rouge, Thunnus thynnus (Linné), Rev. Trav. Inst. Pêch. marit., 18 (2-4): 65-67.
- ANON. 1963: Report of the Federal Fisheries Services 1961-62 and 1962-63. Lagos, Nigeria, 24 pp.
- ANON. 1973: Valores geograficos de la cpue, distribution de talles y crecimiento de la albacora (Thunnus alalunga B.) durante la temporada 1972 en la Region Catabro-Galaica. Comm. Int. Conserv. Thonides de l'Atlantique, Rapp. Cons. Prem. Rech. & Statistique 49, II: 1-7.
- ANTHONY RAJA, B. 1970: Estimation of age and growth of the Indian oil sardine, Sardinella longiceps Valenciennes. Ind. J. Fish., 17 (1-2): 26-46,
- APPARAO, T. 1966: On some aspects of the biology of Lactarius lactarius (Schneider). Ind. J. Fish., 13 (2): 334-349.
- ARNÉ, P. 1938: Rapp. P. v. Réun. Comm. int. Expl. scient. Mer Méditerr. 11: 77-116.
- ARORA, H.L. 1951: An investigation of the California Sand dab Citharichthys sordidus (Girard). Calif. Fish Game, 37 (1): 3-42.
- ARRIGNON, J. 1966: L'Anchois (Engraulis encrasicolus L.) en Côte d'Oranie. Rev. Trav. Inst. Pêch. marit., 30 (4): 317-342.

- ASAMI, T. and M. HANAOKA 1957: On the variation of vertebral number of iwashi, with special reference to its possibility of being a clue for testing difference of races or of environmental conditions during their development. Rep. Nankai Reg. Fish. Res. Lab, 5: 59-73.
- ASLANOVA, N. E. 1954: [Black sea sprat] in Russian. Trud. vsesoinz. nauch. issled. morsk. ryb. Khoz. 28: 75-101.
- AUDOIN, J. 1962: La daurade de l'étang de Thau (Chrysophrys aurata (Linné)). Rev. Trav. Inst. Pêch. marit. 26 (1): 105-126.
- AUSTIN, H. M. 1971: Some aspects of the biology of the Rhomboid Mojarra Diapterus rhombeus in Puerto Rico. Bull. Mar. Sci. 21 (4): 886-903.
- BABEL, J. S. 1967: Reproduction, life history and ecology of the round stingray, Urolophus halleri Cooper. Calif. Fish Game, Fish Bull. 137: 104 pp.
- BABU RAO, M. 1965: Biological studies on the gizzard shad, Anadontostoma chacunda Hamilton (Fam. Clupeidae). J. mar. biol. Ass. Ind., 7 (1): 89-101.
- BACKIEL, T. and J. ZAWISZA 1968: Synopsis of biological data on the Bream Abramis brama (Linnaeus, 1158). FAO Fisheries Synopsis, No. 36.
- BACKUS, R. H. 1962: Age in a small sample of Blue Fish (Pomatomus saltatrix Linnaeus). Brevoria 159: 1-4.
- BAKER, A. N. 1972: Reproduction, early life history, age growth relationships of the New Zealand pilchard, Sardinops neopilchardus (Steindachner). Fish. Res. Bull., N.Z. (5): 1-64.

BALAN, V. 1967: Biology of the silver-belly, Leiognathus bindus (Val.) of the Calicut Coast. Ind. J. Fish. 10 (1): 118-134.

BANERJI, S. K. 1973: An assessment of the exploited pelagic fisheries of the Indian Seas: pp. 114-136. In: Proc. Symp. Living Res. Seas Around India, Spec. Publ., Cen. Mar. Fish. Res. Inst., Cochin, 748 pp.

BAPAT, S. V. 1970: The Bombay duck, Harpodon neherus (Ham.) Bull. Centr. Mar. Fish. Res. Inst., No. 21, 66 pp.

BAPAT, S. V., S. K. BANERJI and D. V. BAL 1952: Observations on the biology of Harpodon neherus (Hamilton). J. Zool. Soc. Ind., 3 (2): 341-356.

BARD, F. X. 1971: La croissance du Germon de l'Atlantique Nord-Est. Comm. Int. Conserv. Thonides de l'Atlantique, Rapp. Cons. Perm. Rech. & Statistique. No. 34, 10, p. 1-4.

BARRETT, I. and G. V. HOWARD 1961: Studies of the age, growth, sexual maturity and spawning of populations of anchoveta (Cetengraulis mysticetus) of the coast of the eastern tropical Pacific Ocean, in English and Spanish. Inter-Amer. Trop. Tuna Comm., Bull. 5 (2): 133-216.

BARNABÉ, G. 1973: Contribution à la connaissance de la croissance et de la sexualité du Loup (Dicentrarchus labrax) de la région de Sète. Ann. Inst. Océan. 49 (1): 49-75.

BARRACA, I. 1964: Quelques aspects de la biologie et de la pêche du Chinchard Trachurus trachurus (L.) de la côte portugaise. Notas Est. Inst. Biol. marit. 29: 1-45.

- BARRO, M. 1968: Premières estimations sur la croissance de Brachydenterus auritus (Val. 1834) en Côte d'Ivoire. O.R.S.T.O.M. Doc. Sci. Prov. Centre d'Abidjan (31): 18 pp.
- BAS, C. 1965: Ecology and rhythm of growth of Gadus poutassou. C. G. Pêches Medit 8, Techn. Pap. 29: 277-279.
- BAUDIN-LAURENCIN, F. 1968: Croissance et âge de l'albacore du Golfe de Guinée. Doc. Sci. Prov. Cent. Rech. Océanogr, Abidjan (21) 12 pp. miméo.
- BAXTER, J. et al. 1960: A study of the yellowtail Seriola dorsalis (Grill). Calif. Fish Game, Fish. Bull. 110: 96 pp.
- BAYAGBONA, E. 1969: Age determination and the Bertalanffy parameters of P. typus and P. senegalensis using the "burnt otolith technique". UNESCO Symp. Oceanogr. Fish. Res. Trop. Atlantic. Abidjan pp. 349-359.
- BAYLIFF, W. H. 1963: Observations on the life history and identity of intraspecific groups of the anchoveta, Cetengraulis mysticetus, in Montijo Bay and Chiriqui Province, Panama. in English and Spanish. Inter-Amer. Trop. Tuna Comm., Bull. 8 (3): 167-197.
- BAYLIFF, W. H. 1966: Population dynamics of the anchoveta, Cetengraulis mysticetus, in the Gulf of Panama, as determined by tagging experiments. Inter-Amer. Trop. Tuna Comm., Bull. 11 (4): 173-352.
- BAYLIFF, W. H. 1967: Growth, mortality and exploitation of the Engraulidae, with special reference to the anchoveta Cetengraulis mysticetus, and the Colorado, Anchoa naso, in the eastern Pacific Ocean. Inter-Amer. Trop. Tuna Comm., Bull. 12 (5): 365-432.

- BAYLIFF, W. H. 1969: Synopsis of biological data on the anchoveta Cetengraulis mysticetus Günter, 1866. FAO. Species Synopsis (43).
- BEARDSLEY, G. L. 1967: Age, growth and reproduction of the dolphin, Coryphæna hippurus in the Strait of Florida. Copeia 1967 (2): 441-451.
- BEARDSLEY, G. L. 1971: Contribution to the population dynamics of Atlantic albacore with comments on potential yields. Fish. Bull. U.S. 69 (4): 845-857.
- BECK, U. 1974: Bestandskundliche Untersuchungen an einigen Fischarten der Grundschieppnetzfisherei auf dem Schelf von Togo (Westafrika). Thesis, Hamburg University, 126 pp.
- BECKETT, J. 1974: Biology of swordfish Xiphias gladius L. in the Northwest Atlantic ocean. In: Shromura and Williams (eds.) Proc. Intern. Billfish Symp. II: 103-106, NOAA, Techn. Rep. NMFS SSRF-675.
- BEESE, G. und R. Kändler 1969: Beiträge zur Biologie der drei nordatlantischen Katfischarten Anarhichas lupus L., A. minor Olafs. und A. denticulatus Kr. Ber. dt. wiss. Kommn für Meeresforsch. 20 (1): 21-59.
- BELL, R. R. 1962: Age determination of the Pacific albacore of the California coast. Calif. Fish Game, 48 (1): 39-48.
- BELLOC, G. 1929: Étude monographique du merlu. Rev. Trav. Inst. Pêch. marit. 2 (3): 231-288.

BELVEZE, H. 1972: Observations complémentaires sur le stock sardinier de l'Atlantique marocain de 1968 à 1971 et estimation de quelques paramètres. Bull. Inst. Pêches Marit, Maroc, (20): 5-69.

BELYANINA, T. S. 1958: Synopsis of biological data on smelt Osmerus eperlanus (Linneaus), 1758. FAO Fisheries Synopsis, No. 78.

BENNETT, S. 1961: Further observations on the fishery and biology of "Chooday" (Sardinella spp.) of Madapam area. Ind. J. Fish. 8 (1): 152-168.

BEN-TUVIA, A. 1956: The biology of Sardinella aurita with a systematic account of the family Clupeidae of the Mediterranean coast of Israel. Fish. Bull., Haifa, (7): 20-4; (8): 14-5, 20-5.

BEN-TUVIA, A. 1960: The biology of the Cichlid fishes of Lake Tiberias and Huleh. Bull. Res. Council, Israel, 8B (4): 153-188.

BEN-TUVIA, A., G. W. KISSIL and D. POPPER 1973: Experiments in rearing rabbitfish (Siganus rivulatus) in sea water. Aquaculture 1: 359-364.

BEN YAMI, M., and T. GLASER 1974: The invasion of Saurida undosquamis (Richardson) into the Levant Basin - an example of biological effect of interoceanic canals. Fish. Bull. (U.S.), 172 (2): 359-373.

BERDEDUE, J. 1955: La pesqueria de la totoaba (Cynoscion macdonaldi Gilbert) en San Felipe, Baja California. Rev. Soc. Mex. Hist. Nat., 16 (1-4): 45-78.



- BERG, L. S., L. S. BOGDANOV, N. I. KOZHIN, and T. S. RASS (editors) 1949: [Commercial Fishes of the USSR. Description of the Fishes in Russian] Pshhepromizdat. 787 pp.
- BERRY, R., S. SAILA, and D. HORTON 1965: Growth studies of winter flounder, Pseudopleuronectes americanus (Waldbaum), in Rhode Island. Trans. Am. Fish. Soc., 94 (3): 259-362.
- BERTALANFFY, L. von 1951: Theoretische Biologie - Zweiter Band: Stoffwechsel, Wachstum. A. Francke AG Verlag, Bern, 418 pp.
- BERTIN, L. 1923: Esquisse du cycle vital de l'épinoche. Comptes Rendus Séances Soc. Biol., 89 p. 730.
- BEVERTON, R. J. H. 1963: Maturation, growth and mortality of clupeid and engraulid stocks in relation to fishing. Cons. Perm. Int. Explor. Mer, Rapp. P.-V. Réun., 154: 44-67.
- BEVERTON, R. J. H. 1965: Catch effort of some I.C.N.A.F. Fisheries. Res. Bull., I.C.N.A.F., 2: 59-72.
- BEVERTON, R. J. H. and S. J. HOLT 1959: A review of the lifespans and mortality rates of fish in nature and their relation to growth and other physiological characteristics. CIBA - Foundation Colloquia on Ageing, 5: 142-180.
- BLACKBURN, M. 1949: Age, rate of growth and general life history of the Australian pilchard in New South Wales waters. S.C.I.R.O. Bull., (242): 86 pp.

- BLACKBURN, M. 1950: Studies on age, growth and life history of the pilchard, Sardinops neopilchardus (Steindachner), in southern and western Australia. Aust. J. Mar. Freshw. Res., 1 (2): 221-258.
- BLACKER, R. 1971: Synopsis of biological data on haddock, Melanogrammus aeglefinus (Linnaeus) 1758. FAO Species Synopsis, No. 84.
- BOEREMA, L. K., G. SAETERSDAL, I. TSUKAYAMA, J. E. VALDIVIA, and B. ALEGRE 1965: Report on the effects of fishing on the Peruvian stock of anchovy. FAO Fish. tech. paper, 55: 44 pp.
- BOLIN, R. L. 1956: Age and growth of the myctophyd fish Lampanictus leucopsauus (Eigenmann & Eigenmann). Proc. intern. Congr. Zool., 14: 541-542.
- BONHAM, K. et al. 1949: Biological and vitamin A studies of dogfish landed in the state of Washington (Squalus suckleyi). St. of Wash., Dept. Fish. Biol. Rept., 49a: 83-114.
- BORTONE, S. A. 1971: Studies on the biology of the sand perch, Diplectum formosum (Perciformes, Serranidae). Tech. Ser. Fla. Dept. Nat. Res., No. 65: 1-27.
- BOTHA, L. 1971: Growth and otolith morphology of the cape Hake Merluccius capensis Cast. and M. paradoxus Franca. Div. Fish. Invest. Rep., (97). 32 pp.
- BOUGIS, P. 1952: La croissance des poisson Méditerranéens. Vie et Milieu, Suppl. 2: 118-146.

- BOUTIÈRE, H. 1958: Les Scorpenidés des eaux marocaines. Trav. Inst. Scientifique Cherifien. Ser. Zool., No. 15, 84 pp.
- BOWERS, A. B. 1954: Breeding and growth of Whiting (Gadus merlangus L.) in Isle of Man waters. J. Mar. Biol. Ass. U.K., 33: 97-122.
- BOWERS, A. B. 1959: Growth of the Witch (Glyptocephalus cynoglossus) (L.) in the Irish Sea. J. du Conseil., 25 (1): 168-176.
- BROCK, V. E. 1954: Some aspects of the biology of the aku, Katsuwonus pelamis, in the Hawaiian Islands. Pacific Sci. 8 (1): 94-104.
- BRUTON, M. and B. ALLANSON 1974: The growth of Tilapia mossambica Peters (Pisces, Cichlidae) in Lake Sibaya, South Africa. J. Fish. Biol., 6: 701-715.
- BURD, A. C. 1962: Growth and recruitment of Herring of the southern North Sea. Fisheries Investigations, Ser. II, 23 (5): 1-42.
- BURHANUDDIN et al. 1974: Some biological aspects of Lemuru, Sardinella sirm (Waldbaum), around Panggang Island. Oseanologi di Indonesia, 1974 (2): 17-24.
- CADENAT, J. 1935: Les Serranidés de la Côte Occidentale d'Afrique (du Cap Spartel au Cap Vert). Off. Sci. Tech. Pêch. Marit., Rev. Trav., 8: 377-422.
- CAMERON, J. 1958: Studies on the Ammodytidae of Isle of Man waters. Diss. Univ. of Liverpool.

- CARLES MARTÍN, C. A. 1967: Datos sobre la biología del banano Elops saurus Linnaeus (Teleostomi: Elopidae). Inst. Nac. de la Pesca, Cuba Centr. Invest. Pesq. Contr. (27), 53 pp.
- CARLES MARTÍN, C. A. 1975: Evaluación de la Pesquería de Bonito en la zona Occidental de Cuba, Rev. Invest. INP, 1 (1): 73-76.
- CARLISLE, J., J. SCHOTT, and N. ABRAMSON 1960: The barred surfperch (Amphistichus argenteus Agassiz) in Southern California. Calif. Fish Game, Fish. Bull., No. 109, 79 pp.
- CASSIE, R. M. 1956: Age and growth of the snapper Chrysophrys auratus Forster, in the Hauraki Gulf. Trans. Royal Soc., N. Z., 84 (2): 329-339.
- CECH, J. and D. WOHLSCHLAG 1975: Summer growth depression in the striped Mullet, Mugil cephalus L. Contr. Mar. Soc., 19: 91-100.
- CHABANNE, J. and R. PLANTE 1969: Les populations benthiques (endofaune, crevettes penaeides, poissons) d'une baie de la côte nord-ouest de Madagascar: écologie, biologie et pêche. Cah. O.R.S.T.O.M., Sér. Océanogr. 7 (1): 41-71.
- CHADWICK, E. 1976: A comparison of growth and abundance for tidal pool fishes in California and British Columbia. J. Fish. Biol., 8: 27-34.
- CHANG, HSIAO-WEI, 1951: Age and growth of Callionymus lyra L. J. Mar. Biol. Ass., 30 (2): 281-296.

CHOO, W. I. 1970: A summary report on yellow croaker in Korea. In: Marr, J. C. (ed.). The Kuroshio: A Symposium on the Japan Current, East West Center Press, Honolulu, 493-496.

CHOTIYARNWONG, A. 1971: Studies on Tilapia nilotica Linnaeus Tilapia mossambica Peters and their hybrids. Thesis, Kasetsart University (Bangkok), 76 pp.

CHYZAN, F. 1959: Salmon and sea trouts in Polish catches in the Baltic in the years 1945-1955. Prace Morskiego Instytutu Rybackiego w Gdyni Nr. 10/A.

CLARK, F. N. and J. B. PHILLIPS 1952: The northern anchovy (Engraulis mordax mordax) in the California fishery. Calif. Fish Game, 38 (2): 189-207.

CLELAND, K. 1947: Studies on the economic biology of the sand whiting (Sillago ciliata C&V). Proc. Linn. Soc. New South Wales, 72 (3-4): 215-228.

CLEMENS, H. B. 1961: The migration, age, and growth of Pacific albacore (Thunnus germon), 1951-1958. Fish. Bull., Sacramento (115): 128 pp.

CLEMENS, W. A. and L. C. CLEMENS 1921: Contribution to the biology of the Muttonfish, Zoarces anguilaris. Contr. Can. Biol. 1918-1920, 6: 69-83.

COLLIGNON, J. 1968: Les trigles des eaux marocaines 1<sup>ière</sup> note: Généralités - Trigla hirundo. Bull. Inst. Pêches Marit. Maroc, (16): 3-34.

CONNOLLY, C. J. 1920: Histories of food fishes. III. The angler. Bull. Biol. Board of Canada, No. 3, 17 pp.

COOPER, R. A. 1967: Age and growth of the tautog, Tautoga onitis (Linnaeus) from Rhode Island. Trans. Am. Fish. Soc., 96 (2): 134-142.

CRIDLAND, C. 1960: Laboratory experiments on the growth of Tilapia spp. I. The value of various foods. Hydrobiol., 15 (1-2): 135-160.

CRISP, D. and S. CARRICK 1975: Some observations on the growth and length-weight relationship of the South Georgia Cod, Notothenia rossii marmorata Fisher, during the first four years of life. J. Fish. Biol., 7: 407-409.

CROCKER, R. 1962: Growth and food of the grey snapper, Lutjanus griseus, in Everglades National Park. Trans. Am. Fish. Soc., 91 (4): 379-383.

DAAN, N. 1974: Growth of North Sea Cod, Gadus morhua. Netherland J. Sea Res. 8 (1): 27-48.

DA FRANCA, P. 1953: Bull. Soc. Portug. Sci. nat. 4, 198.

DAMAS, D., 1909: Contribution à la biologie des Gadides. I.C.E.S. Rapp. & Proc. Verb. 10 (3): 1-277.

DAMM, U. 1975: Untersuchungen zum Wachstum des Wittlings (Merlangius merlangus L.) der Nordsee. Thesis Kiel University 70 pp.

D'ANCONA, U. 1937: La croissance chez les animaux méditerranéens. Rapp. et P.-V. Comm. Int. Expl. Médit., X, 162-224.

DARDIGNAC, J. 1961: Les ombrines des côtes Atlantique du Maroc. Rev. Trav. Inst. Pêch. marit. 25 (3): 263-279.

- DARDIGNAC, J. 1962: La bonite du Maroc Atlantique (Sarda sarda Bloch). Rev. Trav. Inst. Pêch. marit. 26 (4): 399-405.
- DARK, T. 1975: Age and growth of pacific Hake, Merluccius productus. Fish. Bull., U.S., 73 (2): 336-355.
- DAVIES, D. H. 1958: The South African Pilchard (Sardinops ocellata). Preliminary report on the age composition of the commercial catches, 1950-57. Invest. Rep. Div. Fish., S. Afr., (33): 1-20.
- DAVIS, J. and D. LONDER 1969: Life history and ecology of Menidia extensa. Trans. Am. Fish. Soc., 98: 466-472.
- DAVIS, T. 1977: Age determination and growth of the fresh-water Catfish, Tandanus tandanus Mitchell, in the Gwydir River, Australia. Austr. J. Mar. Freshw. Res. 28: 119-137.
- DE, K. and P. DATTA 1974: A note on the analysis of size frequencies of Setipinna taty (Valenciennes) of the lower Sunderbans Region (Hoogly-Mattah Estuary). Ind. J. Fish. 21 (1): 285-289.
- DEASON, H. 1933: Preliminary report on the growth and maturity of the pike-perches (Stizostedion) of Lake Erie. Trans. Am. Fish. Soc. 63: 348-360.
- DEASON, H. and H. HILE 1947: Trans. Am. Fish. Soc. 74: 88.
- DESBROSSES, P. 1937: Note sur le Saint Pierre de Mauritanie, du Senegal et de Guinée, Zeus faber L. mauritanicus natio nova. Rev. Trav. Off. Pêch. marit. 10 (3): 379-414.

- DESBROSSES, P. 1939: Composition et fluctuations du stock de Saint Pierre Zeus faber L. à l'entrée occidentale de la Manche et dans le Golfe de Gascogne. I.C.E.S. Rapp. & Proc. Verb. III. App. 10: 84-87.
- DESBROSSES, P. 1948: Le merlan (Gadus merlangus L.) de la côte française de l'Atlantique, Rev. Trav. Off. Pêch. marit., 14: 71-104.
- DESHMUKH, V. M. 1973: Fishery and biology of Pomadasys hasta (Bloch). Ind. J. Fish. 20 (2): 497-522.
- DE SYLVA, D. 1957: Studies on the age and growth of the Atlantic sailfish, Istiophorus americanus (Cuvier), using length-frequency curves. Bull. Mar. Sci. Gulf Caribb. 7 (1): 1-20.
- DE SYLVA, D. 1963: Systematics and life history of the great barracuda Sphyraena barracuda Waldbaum. Studies in Trop. Ocean. Miami Univ. Inst. of Mar. Sci. (1): 1-179.
- DEVOLD, F. 1938: Fiskeridir Skr. Havundersøk 5 (6).
- DEW, C. BRAXTON 1976: A contribution to the life-history of the cunner, Tautogalabrus adspersus, in Fisher Island Sound, Connecticut. Chesapeake Sci. 17 (2): 101-113.
- DIEUZEIDE, R. and J. ROLAND 1957: Etude biométrique de Sardina pilchardus Walb. et de Sardinella aurita C. V., capturées dans la Baie de Castiglione. Bull. Sta. Aquic. Pêche Castiglione, n.s. 1956 (8): 111-216.



- DIXON, B. 1931: Morphometrical features of the sea trout of Polish Rivers. J. du Conseil 6 (1): 448-457.
- DOUTRE, M. 1960: Les merlus du Sénégal. Mise en évidence d'une nouvelle espèce. Rev. Trav. Inst. Pêch. marit. 24 (4): 513-536.
- DRAGANIK, B. and J. NETZEL 1966: An attempt to estimate the rate of growth of cod in the Southern Baltic on the basis of tagging experiments. I.C.E.S. C. M. 1966/D: 12 (mimeo).
- DRAGESUND, et al. 1973: Regulation of the fishery for capelin. Fiskets Gang 59 (3): 63-67.
- DRUZHININ, A. D. 1975: Some data on Sparid fish (Fam. Sparidae) in the Gulf of Aden Region. J. of Ichthyology 15 (4): 531-541.
- DWIPONGGO, A. 1972: The fishery for and preliminary study of "Lemuru" (oil sardine) at Muntjar, Bali Strait. M.S. Marine Fish. Res. Inst. (Jakarta) 29 pp. and Proc. IPFC 15 III: 221-240.
- EATON, R. C. and R. D. FARLEY 1974: Growth and reduction of dispensation of zebrafish, Brachydanio rerio, in the laboratory. Copeia (1): 204-209.
- EBERT, V. W. and R. SUMMERFELT 1969: Contribution to the life-history of the piute sculpin, Cottus beldingii Eigenmann & Eigenmann, in Lake Tahoe. Calif. Fish Game 55: 100-120.
- EGGLESTON, D. 1970: Biology of Nemipterus virgatus in the Northern part of the South China Sea, pp. 417-424. In: Marr, J. C. (ed.) The Kurishio: A Symposium on the Japan Current, East West Center Press, Honolulu.

- EGGLESTON, D. 1972: Pattern of biology in the Nemipteridae.  
J. Mar. Biol. Ass. India 14 (1): 357-364.
- EHRENBAUM, E. 1923: The Mackerel, Spawning-larval and post  
larval forms, age groups - food - enemies. I.C.E.S.  
Rapp. P.-V. Réun. 30: 1-39.
- EHRICH, S. 1976: Zur Taxonomie, Ökologie und Wachstum von  
Macroramphosus scolopax (Linnaeus, 1758) (Pisces,  
Syngnathiformes) aus dem Subtropischen Nordostatlantik.  
Ber. dt. wiss. Kommn Meeresforsch. N.F. 24 (4): 251-266.
- ELDER, R. D. 1976: Studies on age and growth, reproduction  
and population dynamics of Red Gurnard (Chelidonichthys  
kumu (Lesson and Garnot) in the Hauraki Gulf, New  
Zealand. Fish. Res. Bull., (12): 77 pp.
- ELLIS, R. W. and R. JONES 1956: The Whiting (Gadus merlangus  
L.) in the North Sea. Mar. Res. Scot. 2, 25 pp.
- ELVERTOWSKY, J. 1964: Observations sur les poissons du  
genre Sebastes (Compagne océanographique de la  
"Thalassa" en Juillet-Août 1962 dans l'Atlantique  
Nord-Ouest). Rev. Trav. Inst. Pêch. marit. 28. (4):  
427-435.
- EL ZARKA, S. and H. M. EL SEDFY 1970: The Biology of  
Mugil saliens (Risso) in Lake Quarun, U.A.R.  
Bull. Inst. Ocean. & Fish., Cairo, 1: 1-26.
- EMERY A. and E. D. McCracken 1966: Biology of the Atlantic  
argentine (Argentina silus Ascanius) on the scotian  
shelf. J. Fish. Res. Bd Can. 23 (8): 1145-1160.

- de ESPINOSA, V. 1972: The biology and fishery of the curvina, Cynoscion maracaiboenensis, of Lake Macaraibo. Ser. Recursos y Expl. Pesq. 2 (3): 1-40.
- ERMAN, F. 1959: Observations on the biology of the common grey Mullet (Mugil cephalus L.). Techn. Pap. No. 16 Proc. techn. Pap. gen. Fish. Coun. Med, 5: 157-169.
- EVERSON, I. 1970: The population dynamics and energy budget of Notothenia neglecta Nybelin at Signy Island, South Orkney Islands. Br. Antarct. Surv., Bull. 23: 25-50.
- FAGADE, S. O. 1974: Age determination in Tilapia melanotheron (Rüppel) in the Lagos Lagoon, Nigeria, with a discussion of the environmental and physiological basis of growth marking in the tropics: pp. 71-77 in: Bagenal (ed.) Ageing of Fish, Unwin Bros. Ltd., Old Woking, Surrey, England, 234 pp.
- FAGE, L. 1920: Engraulidae, Clupeidae. Report on the Danish Oceanographical Expeditions 1908 - 1910 to the Mediterranean and Adjacent Seas. No. 6, Vol. 2 (Biology), A. 9: 140 p.
- FAIRBRIDGE, W. 1951 a: The overfishing of the east Australian trawl fishery. Proc. I.P.F.C. 2, Sect. II & III: 73-79.
- FAIRBRIDGE, W. 1951 b: The New South Wales Flathead Neoplatycephalus macrodon (Ogilby). Austr. J. Mar. Freshw. Res. 2 (1): 117-178.
- FARRUGIO, H. 1975: Les muges (Poissons Téléostéens) de Tunisie. Repartition et Pêches. Contribution à leur étude systématique et biologique. Thesis, University of Languedoc, 201 pp.

- FEDDERN, H. A. 1965: The spawning, growth and general behavior of the bluehead wrasse Thalassoma bifasciatum (Pisces, Labridae). Bull. Marine Sciences (Miami) 15 (4): 896-941.
- FELIN, F. E. 1954: Population heterogeneity in the Pacific Pilchard. Fish. Bull. U.S., 54 (86): 201-225.
- FIGUERAS, A. 1955: Datos sobre la edad y crecimiento de la pescadilla (Merluccius merluccius L.) de Levante (sector de Castellon) determinados por medio de los otolitos. Invest. Pesq. I: 73-86.
- FIGUERAS, A. 1964: Edad y crecimiento del bacalao (Gadus morhua) en las pesquerias del Atlantico Nozoeste, en 1962. Publicaciones Technicas de la Junta de Estudio de Pesca. Direccion General de Pesca Maritima. (3): 249-261.
- FIGUERAS, A. 1967: Age and growth of hake (Merluccius merluccius L.) from the Western Mediterranean (Costa Brava, North of Spain). Doc. Techn. 16, Proc. gen. Fish. Counc. Medit. 8: 161-171.
- FITCH, J. 1951: Age composition of the Southern California catch of Pacific mackerel 1939-40 through 1950-51. Calif. Fish Game, Fish. Bull. (83) 73 pp.
- FITCH, J. 1956: Jack Mackerel. Calif. Coop. Oceanic Fish. Invest. Progr. Rep. 1955-1956, pp. 27-28.
- FITCH, J. 1956: Pacific mackerel. Progr. Rep. Calif. Oceanic Fish. Invest.: 29-32.

- FITZ, E. and F. DAIBER 1963: An introduction to the biology of Raja eglanteria Bosc 1802 and Raja erinacea Mitchell 1825 as they occur in Delaware Bay. Bull. Bingham Ocean. Coll. 18 (3): 69-97.
- FOERSTER, R. E. 1929: Contr. Biol, 8: 1 (as cited in: Beverton and Holt, 1959).
- FONDS, M. 1973: Sand gobies of the Dutch Wadden Sea (Pomatoschistus, Gobidae, Pisces). Netherland J. Sea Research 6 (4): 417-478.
- FONTELES-FILHO, A. A. 1970: Estudio sobre a biologia da pesca do parjo, Lutjanus purpureus Poey, no Nordeste Brasileiro, - Dado de 1969. Arq. Ciên. Mar. 10 (1): 73-78.
- FROST, W. and W. SMYLY 1952: J, Animal Ecol. 21: 62.
- FULTON, T. W. 1901: On the rate of growth of the cod, haddock, whiting and Norway pout. Scotland, 19th. Ann. Rep. Fish. Bd., Part III, 154-228.
- FULTON, T. W. 1904: The rate of growth of fishes. 22nd. Ann. Rep. Fish. Bd. Scotland, Part III, 141-240.
- FURNESTIN, J. 1935: La cardine (Lepistoma megastoma Donovan = Lepidorhombus whiff - jagonis Waldbaum). Résumé des connaissances acquises sur la biologie de ce poisson. Rev. Trav., Off. Pêches marit. 8: 203.
- FURNESTIN, J. 1943: Notes préliminaire sur l'anchois (Engraulis encrasicolus L.) du Golfe de Gascogne. Rev. Trav., Off. Pêches marit. 13 (1-4): 197-209.
- FURNESTIN, J. 1948: Observations sur le sprat (Clupea sprattus Linné) des côtes meridionales de France (Atlantique es Méditerranée). Rev. Trav. Inst. Pêch. marit. 14 (1-4): 39-69.

- FUSTER DE PLAZA, M. L. undated: Algunos datos sobre la biologia de la anchoita del sector Bonaerense, C.A.R.P.A.S./2/D, Tech. 12: 11 pp.
- GARNAUD, J. 1962: Nonographie de l'Apogon méditerranéen: Apogon limberbis (L.) 1758. Bull. Inst. Ocean. Monaco No. 1248, 83 pp.
- GARROD, D. J. 1959: The growth of Tilapia esculenta Graham in Lake Victoria. *Hydrobiologica* 12 (4): 268-298.
- GARROD, D. J. & R. GAMBELL 1965: The whiting of the Irish Sea and the Clyde. Fish. Invest. Ser. II 24 (3): 1-64.
- GELDENHUYS, N. D. 1973: Growth of the South African Maarsbanker Trachurus trachurus Linnaeus and age composition of the catches, 1951-1971. South Afr. Sea Fish. Branch Invest. Rep. No. 101, 24 pp.
- GELHAAR, C. 1972: Beiträge zur Biologie der Meeresforelle (Salmo trutta f. trutta L.) in Schleswig-Holstein unter besonderer Berücksichtigung der Farver Au und der Rantzau. Thesis, Kiel University, 109 pp.
- GEORGE, K. and S. K. BANERJI 1964: Age and growth studies on the Indian Mackerel Rastrelliger kanagurta (Cuvier) with special reference to length frequency data collected at Cochin. Ind. J. Fish. 11 (2): 621-638.
- GHÉNO, Y. 1975: Nouvelles études sur la détermination de l'âge et de la croissance de Sardinella aurita Val. Cah. O.R.S.T.O.M., sér. Océanogr. 13 (3): 251-262.
- GHÉNO, Y. and J. C. LE GUEN 1968: Détermination de l'âge et croissance de Sardinella eba (Val.) dans la région de Pointe-Noire. Cah. O.R.S.T.O.M., sér. Océanogr., 6 (2): 69-82.

- GHIRARDELLI, E. 1959: Contribution à la connaissance de la biologie du merlu (Merluccius merluccius L.) en Moyenne Adriatique. Proc. gen. Fish. Counc. Medit. 5 (69): 489-494.
- GIBSON, R. 1970: Observations on the biology of the giant goby, Gobius cobitis Pallas. J. Fish. Biol. 2: 281-288.
- GJØSAETER, J. 1973: Age, growth and mortality of the myctophid fish Benthosema glaciale (Reinhardt) from Western Norway. Sarsia 52: 1-14.
- GORMAN, T. 1963: Biological and economic aspects of the elephant fish Callorhynchus milii Bory in Pegasus Bay and the Canterbury Bight. N. Z. Fish. Techn. Rep. No. 8, 54 pp.
- GOTTLIEB, E. 1957: On the dynamics of the population of coal-fish Gadus virens L., in Norwegian waters, manuscript used by Beverton & Holt (1959).
- GRAHAM, M. 1934. Report on the North Sea Cod. Fish. Invest. Lond., Ser. 2. 13 (4).
- GRANADOS, R. R. 1958: Aspectos biológicos y económicos de la pesquería de Sardina (Sardinops caerulea Girard 1854) en aguas mexicanas del Pacífico. Sec. de Marina, Dir. Gen. de Pesca et Indust. Conex, 135 pp.
- GRANT, C. 1972: The biology of the soldier fish, Gymnapistes marmoratus (Pisces, Scorpaenidae). Austr. J. Mar. Freshw. Res. 23 (2): 151-163.

- GRAVIER, R. 1961: Les Bars (loups) du Maroc Atlantique Morone labrax (Linné) et Morone punctata (Bloch). Rev. Trav. Inst. Pêch. marit. 25 (3): 281-292.
- GUEGUIN, J. 1969: Croissances de la dorade Pagellus centrodontus Delaroche. Rev. Trav. Inst. Pêch. marit. 33 (3): 251-264.
- GUINARD, J. P. 1966: Recherches sur les Labridae (Poissons teleostéens) perciformes des côtes européennes. Systematique et biologie. Naturalia Monspeliensia Ser. Zoologique fasc. 5: 1-247.
- GUNDERMANN, N. and D. POPPER 1975: Some aspects of Recolonization of Coral Rocks in Eilat (Gulf of Aqaba) by fish populations after poisoning. Marine Biology 33: 109-117.
- HAMADA, R. 1971: Age and growth of the Ribbon-fish, Trichiurus lepturus Linné, based on the transverse section of the otolith. Bull. Sekai Fish. Res. Lab., (41): 11-29.
- HANABUCHI, N. 1967: Study on the Nibe Croaker Miichthys imbricatus (Matsubara) in the East China Sea and Yellow Sea II on the age and growth. Bull. Sekai, Reg. Fish. Res. Lab. (35): 51-60.
- HANATAKA, M. and K. SEKINO 1962: Ecological studies on the Japanese Sea Bass Lateolabrax japonicus. I. Growth. Bull. Jap. Soc. Sci. Fish. 28 (9): 857-861.
- HANNERZ, L. 1964: Regional and annual variations in the growth of whiting (Gadus merlangus Linné). Inst. Mar. Res. Lysekil Ser. Biol., Rept. No. 14: 1-66.



- HANSEN, D. 1970: Food, growth, migration, reproduction and abundance of Pinfish, Lagodon rhomboides and Atlantic croaker, Micropogon undulatus, near Pensacola, Florida. U.S. Fish. Comm. Bull. 68 (1): 135-146 (1968).
- HANSEN, P. 1949: Studies on the Biology of the Cod in Greenland Waters, Rapp. Cons. Explor. Mer. CXXIII.
- HANUMANTHARAO, L. 1974: Studies on the biology of Cirrhinus mrigala (Hamilton) of the River Godavary. Ind. J. Fish. 21 (2): 303-322.
- HANYU, I. 1956: On the age and growth of argentina (Argentina semifasciata Kishinouye). Bull. Jap. Soc. sci. Fish. 21 (9): 991-998.
- HART, I. 1948: Publ. Ontario Fish. Res. Lab. No. 34.
- HASHEM, M. T. 1972: The age, growth and maturity of Labeo niloticus Forsk. from the Nozha Hydrodrome in 1968-1970. Bull. Inst. Ocean. Fish. 2: 84-102.
- HASHEM, M. T. and K. HUSSEIN 1973: Some biological studies of the Nile perch (Lates niloticus (C. & V.) in the Nozha Hydrodrome. Bull. Inst. Ocean. Fish. 3: 321-393.
- HASS, G. 1937: Variabilitätsstudien an Gobius niger L., Gobius minutus Pallas und Cottus scorpius L. Kieler Meeresforsch. I, 279-321.
- HATANAKA, M. and K. SEKINO 1962: Ecological studies on the Japanese Sea Bass Lateolabrax japonicus II. Growth. Bull. Japn. Soc. sci. Fish. 28 (9): 857-861.

- HAYASHI, S. and K. KONDO 1957: Growth of the Japanese anchovy - IV. Age determination with the use of scales. Bull. Tokai Reg. Fish. Res. Lab., 17: 31-64.
- HAYASHI, Y. 1976: Studies on the growth of the red tilefish in the East China Sea. II. Estimation of age and growth from otoliths reading. Bull. Jap. Soc. sci. Fish. 42 (11): 1243-1250.
- HEALD, E. and R. GRIFFITHS 1967: The age determination, from scale readings, of the sardine, Sardinella anchovia, of the Gulf of Cariaco, Eastern Venezuela, Ser. Recursos y Expl. Pesq. 1 (10): 375-446.
- HENNEMUTH, R. C. 1961: Size and year class composition of catch, age and growth of yellowfin tuna in the eastern tropical Pacific Ocean for the years 1954-1958. Bull. Inter-Amer. trop. Tuna Comm. 5 (1): 1-112.
- HENRY, K. A. 1971: Atlantic Menhaden (Brevoortia tyrannus). Resource and Fishery - Analysis of Decline. N.O.A.A. Technical Report NMFS SSRF-642, 32 pp.
- HERALD, R. and M. RAKOWICZ 1951: Aquarium Journal 22: 234 p.
- HILE, R. 1936: Summary of investigations on the morphometry of the cisco, Leucichthys artedi (Le Sueur), in the Lakes of the Northern Highland, Wisconsin. Rap. Mich. Acad. Sci. Arts & Letters 21: 619-634.
- HILE, R. and F. JOBES 1941: Age, growth and production of the yellow perch, Perca flavescens (Mitchell), of Saginaw Bay. Trans. Am. Fish. Soc. 70: 102-122.

- HIRSCHHORN, G. 1974: The effect of different age ranges on estimated Bertalanffy growth parameters in three fishes and one mollusk of the northeastern Pacific Ocean. 192-199 in: Bagenal (ed.) Ageing of Fish, Unwin Bros. Ltd., Old Woking, Surrey, England, 234 pp.
- HNATEVIČ, B. 1960: Das Wachstum des Barsches (Perca fluviatilis L.) in dem Orava-Staubecken (Slowakei). Zeitschr. f. Fischerei und d. Hilfswiss. IX (1); 85-93.
- HOEK, P. 1912: Les Clupeidé (Le Hareng excepté) et leurs migrations I l'Anchois. Rapp. Proc. Verb. I.C.E.S. 20.
- HOHENDORF, K. 1966: Eine Diskussion der Bertalanffy Funktionen und ihre Anwendung zur Charakterisierung des Wachstums von Fischen. Kieler Meeresforsch. 22; 70-97.
- HOFFMEISTER, H. 1957: Beiträge zur Biologie des Wittlings (Gadus merlangus L.) und der Seequabbe (Onos cimbrius L.) Thesis, Kiel Univ. 111 pp.
- HOLDEN, M. T. 1972: The growth rates of Raja brachyura, R. clavata and R. montagui as determined from tagging data. J. du Conseil 34 (2):161-168.
- HOLDEN, M. T. 1974: Problems in the rational exploitation of elasmobranch populations and some suggested solutions, p. 117-137 in: Harden Jones, F.R. (ed.) 1974, Sea Fisheries Research. Elek Scientific Books, Ltd. London, 510 pp.
- HOLDEN, M. J. and P. MEADOW 1962: The structure of the spine of the spur dogfish (Squatus acanthias L.) and its use for age determination. J. Mar. Biol. Ass. U.K. 42: 179-197.

- HOLDEN, M. J. and T. WILLIAM 1974: The biology and population structure of bass, Dicentrarchus labrax, in English waters. J. Mar. Biol. Ass. U.K. 54: 91-107.
- HOLLAND, G. A. 1969: Age, growth and mortality of races in English sole (Parophrys vetulus) in Puget Sound, Washington. Pac. Mar. Fish. Comm. Bull. 7: 35-50.
- HONDA, K. 1950: On the scales of Amber-fishes, Seriola quinqueradiata (T. & S.). Bull, Jap. Soc. sci. Fish. 15 (2): 97-99.
- HONGSKUL, V. 1974: Population dynamics of Platu Rastrelliger neglectus (van Kampen) in the Gulf of Thailand. Proc. IPFC 15 III: 297-350.
- HORNELL, J. and M. R. NAIDU 1924: A contribution to the life history of Indian oil sardine with notes on plankton of the Malabar Coast. Madras Fish. Bull, 17 (5): 129-197.
- HORSTMANN, U. 1975: Some aspects of the mariculture of different siganids species in the Philippines. The Phillipine Scientist, 12: 5-20.
- HUBBS, C. L. 1921: The ecology and life history of Amphigonopterus aurora and other viviparous perches of California. Biol. Bulletin 40 (4): 181-209.
- HUBOLD, G. 1975: Das Wachstum des Buchan-Herings (Clupea harengus L.) in den Jahren 1955 bis 1973. Thesis, Kiel University 51 pp.
- HUGHES, S. E. 1974: Stock composition, growth, mortality and availability of pacific saury, Cololabis saira, of the northeastern Pacific Ocean. Fishery Bull. U.S., 72 (1): 121-131.

- HUREAU, J. C. 1970: Biologie comparée de quelques Poissons antarctique (Notothenidae). Bull. Inst. Oceanogr., Monaco, 68 (1391): 244 pp.
- HUREAU, J. C. and C. OZOUF 1977: Détermination de l'âge et croissance du coelacanth Latimera chalumnae Smith, 1939 (Poisson, Crossopterygien, Coelacanthidé). Cybium 3<sup>ième</sup> Série 2: 129-137.
- HYND, J. and C. LUCAS 1974: Population dynamics of the southern Bluefin tuna. Proc. I. P.F.C. 15 (III): 424-435.
- ICNAF Redbook 1973 - Part 1 p. 94 - Dartmouth, Canada.
- IDYLL, C. P. & D. SILVA 1963: Synopsis of biological data on the blackfin tuna Thunnus atlanticus (Lesson) 1830 (western Atlantic). Sp. Synopsis No. 25: 761-770. FAO Fish. Biol. Synopsis No. 68.
- IH-HSIU TUNG 1970: Studies on the fishery biology of the Grey Mullet, Mugil cephalus, in Taiwan, pp. 491-504. In: Marr, J. C. (ed.), The Kuroshio: A Symposium on the Japan Current, East West Center Press, Honolulu.
- IKENOUYE, H. and H. MASUZAWA 1968: An estimation of growth equation basing on the results of tagging experiments of the Japanese Alfonsin fish. Bull. Jap. Soc. sci. Fish. 34 (2): 97-102.
- ILES, T. D. and P. O. JOHNSON 1962: The correlation table analysis of a sprat (Clupea sprattus L.) Year-Class to separate two groups differing in growth characteristics. J. du Conseil 27 (1): 287-303.

ISAKOV, V. 1973: Growth and total mortality of mackerel from the New England area. ICNAF Fish. Res. Doc. 73/23 Ser. No. 2956.

ISHYAMA, R. 1951: Age determination of Raja hollandi Jordan & Richardson, chiefly in the waters of the East China Sea. Bull. Jap. Soc. sci. Fish 16 (12): 119-124.

IVERSEN, E. S. 1955: Size frequencies and growth of central and western Pacific bigeye tuna. Spec. sci. Rep. U.S. Fish Wildl. Serv.-Fish., (162): 1-40.

JAMES, P. S. B. R. 1967: Ribbon fishes of the family Trichiuridae of India. Mem. I., Mar. Biol. Ass. India, 228 pp.

JAPANESE FISHERIES AGENCY 1952: Biology and Fisheries of mackerel in Japan (a review) mimeo, in Japanese.

JENSEN, A. S. and P. M. HANSEN 1931: Investigations on the Greenland cod (Gadus callarias L.) with an introduction on the history of the Greenland cod fishery. I.C.E.S. Rapp. & Proc. Verb. 72 (1).

JHINGRAN, V. G. 1975: Fish and Fisheries of India. Hindustan Publishing Corp., Delhi 954 pp.

JILLETT, J. B. 1968: The biology of Acanthoclinus quadridactylus (Bloch & Schneider) (Teleostei-Blenniodea) I Age, growth, food. Austr. J. Mar. Freshw. Res. 19 (1): 1-8.

JOENOS, G. 1961: Über die Biologie und fischereiliche Bedeutung der Lengfische (Molva molva L., Molva byrkelange Wal) und des Lumb (Brosmius brosme Asc.). Ber. dt. Wiss. Kommn Meeresforsch. 16 (2): 129-160.

- JOHANSEN, F. 1925: Natural history of the cunner (Tautoglabrus adspersus Waldbaum). Can. Contr. Biol. No. 17: 425-467.
- JOHNSON, A. G. and H. HORTON 1972: Length-weight relationships, food habits and sex and age determination of the ratfish, Hydrolagus collei (Lay & Bennett). Fish. Bull, 70 (2): 421-429.
- JOHNSON, C. K. 1972: Biology and ecology of Callionymus belcheri (Pisces, Callionymidae). Copeia (3): 461-470.
- JOHNSON, C. R. 1970: Age of the snowy snailfish, Liparis pulchrettus, Jap. J. Ichthyol. 17: 121-123.
- JOHNSON, L. 1966: Consumption of food by the resident populations of Pike, Exos lucius, in Lake Windermere. J. Fish. Res. Bd Canada 23 (10): 1523-1535.
- JONES, A. 1974: Sexual maturity, fecundity and growth of the turbot, Scophthalmus maximus L. J. Mar. Biol. Assn. U.K. 54: 109-125.
- JONES, B. W. and J. JONSSON 1971: Coalfish tagging experiments at Iceland. Rit, Fiskideildar 5 (1): 27 pp.
- JONES, B. W. 1966: The cod and the cod fishery at Faroe. Fish. Invest., Lond. Ser. 2 24 (5): 1-32.
- JONES, J. W. and H. B. N. HINES 1950: The age and growth of Gastreosteus aculeatus, Pygosteus pungitius and Spinachia vulgaris as shown by their otoliths. J. Anim. Ecol., 19, 59-73.
- JONES, R. 1962: The growth of haddock in the North Sea and at Faroe. Mar. Res. Scotland 1962 (2): 1-19.

- JONES, R. 1976: Mesh Regulation in the Demersal Fisheries of the South China Sea Area. South China Sea Dev. & Coord. Progr. SCS/76/WP/34 Manila 75 pp.
- JONES, R. and J. R. G. HISLOP 1975: Changes in North Sea haddock and whiting. ICES Symposium on the changes in the North Sea fish stocks and their causes No. 25, 32 pp.
- JONSSON, J. 1957: Icelandic cod investigation in South west and South east Greenland waters 1957. Cons. Int. Explor. Mer. Ann. biol. 14: 115-118.
- JONSSON, J. 1965: Temperature and growth of cod in Icelandic waters. I.C.N.A.F. Spec. Publ. 6: 537-539.
- JOSEPH, D. 1962: Growth characteristics of two southern California surf fishes, the California Corbina and Sponfin Croaker, family Sciaenidae. Calif. Fish Game, Fish Bull. (119): 1-54.
- JOSEPH, J. 1963: Contribution to the biology of the engraulid Anchoa naso (Gilbert and Pierson, 1898) from Ecuadorian waters. Bull. Inter-Amer. Trop. Tuna Comm., 8 (1): 1-30.
- JOSEPH, J. and T. CALKINS 1969: Population dynamics of the skipjack tuna (Katsuwonus pelamis) in the eastern Pacific Ocean. Bull. Inter-Amer. Trop. Tuna Comm. 13 (1): 1-273.
- KAGANOWSKAYA, S. 1933: Method for ascertaining the age and composition of the catch of Squalus acanthias L. Bull. Far East Branch of the Acad. of Sci. of USSR Nos. 1, 2, 3 pp 5-6 and 139-141.



- KAGWADE, P. V. 1970: The Polynemid fishes of India.  
Bull. Centr. Mar. Fish. Inst. No. 18: 69 pp.
- KAISER, C. 1973: Age and growth of Horse Mackerel,  
Trachurus murphyi off Chile. Trans. Am. Soc. Fish.  
(3): 591-595.
- KAJIKAWA, H. 1973: Determination of age and growth in the  
marine Catfish Plotosus anguilaris by the use of  
otoliths. Sci. Rep. Shima Marineland 2: 5-22.
- KALININA, E. M. 1963: [Growth and feeding of Ctenilabrus  
and Symphodus] in Russian . Trud. Sevastopol Biol.  
Stn. 16: 323-326.
- KAKUDA, S. 1976: Personal communication.
- KÄNDLER, R. 1941: Untersuchungen über Fortpflanzung,  
Wachstum und Variabilität der Arten des Sandaals  
in Ost- und Nordsee, mit besonderer Berücksichtigung  
von Ammodytes tobianus L. Kieler Meeresforsch. 5  
(1): 45-139.
- KÄNDLER, R. 1944: Untersuchungen über den Ostseedorsch  
während der Forschungsfahrten mit dem R. F. D.  
"Poseidon" in den Jahren 1925-1938. Ber. dt. Wiss.  
Kommn Meeresforsch. 11 (2): 137-245.
- KARNELLA, C. and R. GIBBS 1977: The lanternfish Lobianchia  
dofleini: An example of the importance of life-  
history information in prediction of oceanic sound  
scattering, pp. 316-379 in: Andersen, N. and B.  
Zahuranec (eds.) Oceanic Sound Prediction, Plenum  
Press, New York.

- KELLY, G. and R. WOLF 1959: Age and growth of the Redfish Sebastes marinus in the Gulf of Maine. Fish. Bull., U.S. 156 Vol. 60:1-31.
- KENNEDY, W. A. 1954: Growth, maturity and mortality in the relatively unexploited lake trout, Cristivomer namaycush, of Great Slave Lake. J. Fish. Res. Bd Canada 11: 827-852.
- KESTEVEN, G. 1942: Studies in the biology of Australian Mullet. Bull. Counc. Scientific and Industr. Res. Melbourne, 157: 1-147.
- KETCHEN, K. S. 1964: Preliminary results of studies on growth and mortality of Pacific cod (Gadus macrocephalus) in Hecate Strait. J. Fish. Res. Bd Canada 21: 1051-1068.
- KETCHEN, K. S. 1975: Age and growth of dogfish Squalus acanthias in British Columbia Waters. J. Fish. Res. Bd Canada 32: 43-59.
- KETCHEN, K. S. and C. FORRESTER 1966: Population Dynamics of the petrale sole, Eopsetta jordani, in waters off Western Canada. Bull. Fish. Res. Bd Canada (115) 195 pp.
- KEYSLER, H. D. 1966: Beiträge zur Biologie und wirtschaftlichen Bedeutung von Argentina silus. Thesis, Kiel University, 100 pp.
- KIBESAKI, O. 1949: Ecological studies on Döderlinia berycoides in the South-Western part of Sea of Japan. I. Age and growth. Bull. Jap. Soc. sci. Fish., 15 (5): 221-225.

- KIKUCHI, T. 1966: An ecological study on animal communities of the Zostera marina belt in Tamioka Bay, Amakusa, Kyushu. Publ. Amakusa Mar. Biol. Lab. 1 (1) 106 pp.
- KIMURA, K. 1932: Growth curves of bluefin tuna and yellowfin tuna based on the catches near Shigedera, on the west coast of Prov. Izu. Bull. Jap. Soc. sci. Fish., 1 (1): 1-4.
- KIRILLOV, F. N. 1936: The sand eel of Novaya Zemla. Dokl. Akad. Nauk SSSR 4 (13) (5): 233-235.
- KLIMA, E. F. 1959: Aspects of the biology and the Fishery for Spanish Mackerel, Scomberomorus maculatus (Mitchill), of Southern Florida. Florida Board of Conservation, Tech. Bull., No. 27, 38 pp.
- KLIMA, E. F. and D. C. TABB 1959: A contribution to the biology of the spotted weakfish, Cynoscion nebulosus (Cuvier), from Northwest Florida, with a description of the fishers. Florida Board of Conservation, Tech. Bull. No. 30, 25 pp.
- KNUDSEN, J. 1950: Contributions to the biology of the whiting (Gadus merlangus L.) in the Danish waters. Rep. Danm. Biol. Stat. No. 52: 28-40.
- KNUDSEN, H. 1964: Studies on whiting (Merlangius merlangus L.) in the North Sea, Skagerrak and Kattegat I-II. Medd. Danm Fisk Havunders NS 4 (5): 95-136.
- KONDO, K. 1966: Growth of Japanese mackerel II. Age determination with use of scale. Bull. Tokai Reg. Fish. Res. Lab. 45: 31-60.

- KONDO, K. 1974: Ecological Monograph of the Japanese anchovy, Engraulis japonicus (Houttuyn) - the perspective, methodology and problems for the future. Proc. I.P.F.C. 15 (III): 195-211.
- KOSAKA, M. et al. 1963: Ecological studies on the Ribbon-fish, Trichiurus lepturus Linné, in Suruga Bay. J. Coll. Mar. Sci. Techn. Tokai Univ. (2): 131-146.
- KOSIOR, M. 1976: The South Baltic cod growth parameters and length-weight relationship. I.C.E.S. CM/1976/P: 20 7 pp.
- KOTO T. and T. KODAMA 1962: Some considerations on the growth of Marlin, using size frequencies in commercial catches I. Attempts to estimate the growth of sailfish. Rep. Nakai Reg. Fish. Res. Lab. (15): 97-108.
- KRISHNAMOORTHY, B. 1971: Biology of the threadfin bream, Nemipterus japonicus. Ind. J. Fish. 18 (1-2): 1-21.
- KRISHNAYA G. 1963: On the use of otoliths in the determination of age and growth in the Gangetic whiting Sillago panijus (H. & B.), with notes on its fishery in the Hoogly Estuary. Ind. J. Fish. 10 A (1): 391-412.
- KRÜGER, K. 1942: Erneutes Auftreten der Scharbzunge, Drepanopsetta plattessoides Fabr., in der Westlichen Ostsee. Kieler Meeresforsch. 4: 18-37.
- KÜHLMORGEN-HILLE, G. 1970: A contribution to the knowledge of the growth of Saurida undosquamis Richardson in the Gulf of Thailand, pp. 467-470. In: Marr, J. C. (ed.), The Kuroshio: A Symposium on the Japan Current, East West Center Press, Honolulu, 614 pp.

- KÜHLMORGEN-HILLE, G. 1977: Preliminary study on the life history of the flatfish Psettodes erumei from the Gulf of Thailand. pp.261-268. In: K. Tiews (ed.) Fisheries Resource and their Management in South East Asia. DSE/BFA Fisch./FAO, Berlin 1977. 522 pp.
- KUME, S. and J. JOSEPH 1966: Size composition, growth and sexual maturity of bigeye tuna, Thunnus obesus (Lowe), from the Japanese longline fishery in the Eastern Pacific Ocean. Inter-Amer. Trop. Tuna Comm., Bull., 11 (2); 47-49.
- KUNIYUKI, K. et al. 1975: The age determination and growth of immature Red Sea Bream, Chrisophrys major Temnick & Schlegel in the Central Waters of Hiroshima Prefecture. Bull. Nankei Reg. Fish. Res. Lab. 8 : 81-100.
- KUTTY, M. NARAYANAN 1961: Scales and otoliths of the "Koth" Otolithoides brunneus (Day) as age indicator. Ind. J. Fish. 8 (1): 145-151.
- KYLE, H. 1926: Beobachtungen über das Wachstum des Steinbutts (Rhombus maximus). Fischerbote 18: 209-212.
- LAM, T. J. 1974: Siganids: their biology and mariculture potential. Aquaculture 3: 325-354.
- LAMP, F. 1965: Beiträge zur Bestandskunde und Fortpflanzungsbiologie der Seeskorpiene, Myoxocephalus scorpius (Linnaeus 1758) und Taurulus bubalis (Euphrasen 1786) in der Kieler Förde. Thesis, Kiel University, 140 pp.
- LANE, E. 1967: A study of the atlantic midshipmen, Porochthys porisissimus in the vicinity of Port Aransas, Texas. Contr. Mar. Sci 12: 1-53.

- LARRAÑETA, G. 1965: Les constantes de la croissance de la sardine de Castellon. Proc. gen. Fish. Coun. Médit. 8: 273-276.
- LARRAÑETA, M. G. and J. LÓPEZ 1958: El crecimiento de la sardina (Sardinia pilchardus Walb.) de las costas de Castellón. Invest. Pesqu. 6: 53-82, (1957).
- LASKARIDIS, K. 1948: Study of the biology of the sardine (Clupea pilchardus Walb.) in Greek waters. Prakt. Hellen. hydrobiol. Inst. 2: 11-88.
- LASSÈRE, G. and P. J. LABOUR 1974: Étude comparée de la croissance de la daurade, Sparus auratus L. des régions d'Arcachon et de Sète (deuxième note). Vie et Milieu 24 (2)A: 357-364.
- LATIF, A. and S. SHENOUDA 1972: Biological studies on Rhonciscus striatus (Fam. Pomadasyidae) from the Gulf of Suez. Bull. Inst. Ocean. Fisheries (Egypt), 2: 103-134.
- LEBEAU, A. and J. C. CUEFF 1975: Biologie et Pêche du Capitaine, Lethrinus enigmaticus (Smith 1959) du Banc de Saya de Malha (Océan Indien). Rev. Trav. Inst. Pêch. marit. 39 (4): 415-442.
- LE DREW, B. and J. GREEN 1975: Biology of the radiated shanny Ulvaria subbifurcata Storer in Newfoundland (Pisces, Stichaeidae). J. Fish. Biol. 7: 485-495.
- LEE, J. Y. 1961: Le sardine du Golfe du Lion (Sardina pilchardus sardina Regan). Rev. Trav. Inst. Pêch. marit. 25 (4): 417-512.

- LEE, J. Y. 1963: Les Argentines du golfe du Lion, Argentina sphyraena L., Argentina leioglossa Val. Rev. Trav. Inst. Pêch. marit. 27 (1-4): 189-202.
- LEE, J. Y. and C. JUGE 1965: Observations morphologiques et biologiques sur les anchois (Engraulis encrasicolus) du Golfe du Lion. Commn Int. Explor. Sci. Mer. Medit. Rapp. P.-V. Réun. 18 (2): 221-224.
- LE FRANC, G. 1966: Note Préliminaire sur la morue du Sud de la Mer du Nord et de la région du Pas-de-Calais. Science et Pêche Bull. Inst. Pêches marit. (154): 1-8.
- LE FRANC, G. 1970: Biologie de la morue du Sud de la Mer du Nord et de la Manche Orientale. Rev. Trav. Inst. Pêch. marit. 34 (3): 277-296.
- LE GALL, J. 1939: Quelques resultats des recherches faites sue la biologie du maquereau de l'Atlantique. Rapp. Cons. Expl. Mer 111: 13-14.
- LE GALL, J. 1951: Corrélation taille/fréquence des germes en 1950. Ann. biol., Copenhague 7: 64-66.
- LE GALL, J. 1969: Croissance d'un poisson téléostéen, l'Agonus cataphractus (Linné). Rapports avec le cycle sexuel et le cycle de l'activité throidienne. Vie et Milieu 20 (1)A: 153-234.
- LEGAND, M. 1967: Cycles biologiques des poissons mesopelagiques dans l'Est de l'océan Indien. Première Note. Cah. ORSTOM, sér. Océanogr. 5 (2): 69-78.

- LE GUEN, J. C. 1971: Dynamique des populations de Pseudotolithus (Fonticulus) elongatus (Bowd. 1825) Poissons, Sciaenidae. Cah. O.R.S.T.O.M., sér. Océanogr. 9 (1): 3-84.
- LE GUEN, J. C., F. BAUDIN-LAURENCIN and C. CHAMPAGNAC 1969: Croissance de l'albacore (Thunnus albacores) dans les régions de Pointe-Noire et de Dakar. Cah. O.R.S.T.O.M., sér. Océanogr. 7 (1): 19-40.
- LE GUEN, J. C. and C. CHAMPAGNAC 1968: Croissance des albacores dans les régions de Pointe-Noire et de Dakar. Doc. Centre O.R.S.T.O.M. Pointe-Noire (431): 25 pp. miméo.
- LE GUEN, J. C. and G. T. SAGAKAWA 1973: Apparent growth of yellowfin tuna from the eastern Atlantic Ocean. Fish. Bull., U.S. 11: 175-187.
- LE ROUX, P. J. 1961: Growth of Tilapia mossambica Peters in some Transvaal Impoundments. Hydrobiol. 18 (1-2): 165-175.
- LESSENT, P. 1968: Essais d'hybridation dans le genre Tilapia à la station de recherches agricoles de Bouaké, Côte d'Ivoire. F.A.O. Fisheries Rep. 44, IV/E-5: 148-159. Vol. 4.
- LETACONNOUX, R. 1947: Notes sur le Gadiculus thori Johs Schmidt. Bull. Inst. Ocean. No. 913 7 pp.
- LETACONNOUX, R. 1954: Recherches sur les poissons: Morue. Rev. Trav. Inst. Pêch. marit. 18 (2-4): 79-81.
- LETACONNOUX, R. 1955: Rapport sur le merlu. I.C.E.A. Rapp. & Proc. Verb. 137: 40-43.



- LE TRONG PHAN and A. KOMPOWSKI 1972: The bronze bream-  
Pagellus acarne Risso from North West African Region.  
Acta Ichthyologica et Piscatoria 2 (1): 1-18.
- LE TRONG PHAN and A. KOMPOWSKI 1972: A study of Pagellus  
coupei Dieuzeide from the North West African Region.  
Acta Ichthyologica et Piscatoria 2 (1): 19-30.
- LI, KWANG-MING 1958: Age determination of the Hongkong  
Wave Sea Bream Taius tumifrons (Tanaka & Schlegel).  
Hongkong Univ. Fish. J. (2): 59-69.
- LI, KWANG-MING 1960: On the biology of the Hongkong golden  
thread, Nemipterus virgatus (Houttuyn). Hongkong  
Univ. Fish. J. 3: 1-18.
- LICHATOWICH, T. and D. POPPER 1975: Report on the growth  
of rabbitfish in fish ponds in Fiji. Aquaculture 5:  
211-212.
- LILLELUND, K. 1961: Untersuchungen über die Biologie und  
Populationsdynamik des Stintes (Osmerus eperlanus  
eperlanus L.) der Elbe. Arch. f. Fischereiwiss. 12,  
Beiheft.
- LIMBAUGH, C. 1961: Life history and ecological notes on the  
black croaker. Calif. Fish Game 41 (2): 163-174.
- LIN, F. H. 1956: Studies on black Croaker (Argyrosomus  
nibe J. & T.) in the Northern Trawling Ground of  
Taiwan. Rep. Inst. Fish. Biol. Ministry of Econ.  
Affaires & National, Taiwan Univ. 1 (1): 62-64.
- LONGHURST, A. R. 1963: The bionomics of the fisheries  
resources of the eastern tropical Atlantic. Fishery  
Public. Colon. Off. (London) (20), 65 pp.

- LONGHURST, A. R. 1964: A study of the Nigerian trawl fishery. Bull. I.F.A.N. Ser. A 26 (2): 337-402.
- LONGHURST, A. R. 1965: The biology of West-African Polynemid Fishes. J. du Conseil 30 (1): 58-74.
- LOPEZ, J. 1963: Age de la Sardine (Sardina pilchardus Wald.) de Barcelone. Doc. Tech. No. 29, Proc. gen. Fish. Coun. Médit. 7: 299-308.
- LUNDBECK, J. 1953: German Market investigation on cod mainly in the North/eastern area. J. Cons. int. Explor. Mer. 136, p. 33-39, (as cited in LeFranc 1970).
- LUTHER, G. 1973: The dorab fishery resources of India. PP 445-454 in: Proceedings of the Symposium on the Living Resources of the Seas around India, Spec. Publ., Cen. Mar. Fish. Res. Inst., Cochin, 748 pp.
- MACER, C. T. 1966: Sandeels (Ammodytidae) in the southwestern North Sea; their biology and fishery. Fish. Invest. Min. of Agri., Food & Fish. Ser. 2, 24 (6): 1-55.
- MACKAY, D. 1970: Populations of trouts and graylings in two Scottish rivers. J. Fish. Biol. 2 (1): 39-45.
- MAGNIN, E. 1965: Croissance de l'esturgeon Acipenser fulvescens, Raf. vivant dans le bassin hydrographique de la rivière Nottaway, tributaire de la baie James. Le Naturaliste Canadien 92 (8-9): 193-203.
- MAGNIN, E. and C. FRADETTE 1975: Croissance de la barbu (Ictalurus punctatus) du fleuve St. Laurent près de Quebec. J. Fish. Res. Bd Canada 32 (10): 1867-1870.

- MAJAROVA, A. A. and N. I. CHUGUNOVA 1954: [Biology, distribution and evaluation of the Stock of the Black Sea anchovy] in Russ. Trudy. Vses. Nauchno-Issled. Inst. Morsk. Ryb. Khoz. Okeanogr. 28: 5-33.
- MAJOR, R. and H. SHIPPEN 1970: Synopsis of biological data on Pacific Ocean Perch Sebastes alutus. F.A.O. Species Synopsis No. 79 and NOAA Circular No. 347, 38 pp.
- MARCILLE, J. and B. STEQUERT 1976: Croissance des jeunes albacores Thunnus albacares et patudos Thunnus obesus de la côte nord-ouest de Madagascar. Cah. O.R.S.T.O.M., sér. Océanogr. 14 (2): 153-162.
- MARR, J. C. 1943: Stanford Ichthyol. Bull. 2: 157.
- MARR, J. C. 1960: The causes of major variations in the catch of the Pacific Sardine (Sardinops caerulea (Girard)). F.A.O. Proceedings of the World Scientific Meeting on the Biology of Sardines and Related Species. Vol. III, 667-791.
- MARTIN, F. S. and G. G. GONZALES 1960: Observaciones y notas sobre la explotación de la sardina (Clupanodon pseudohispanicus Poey) en Venezuela. In: F.A.O. Proceedings of the World Scientific Meeting on the Biology of Sardines and Related Species. Vol. III, 855-876.
- MARTIN, W. R. 1953: Vital statistics of the cod and haddock Subarea 4. I.C.N.A.F. 3rd Ann. Meet. Doc. 39.
- MASUZAWA, H. 1967: Studies on the resources of Isaki, Parapristipoma trilineatum I. Age determination and growth. Bull. Jap. Soc. sci. Fish. 33 (9): 812-817.

- MATHER, F. J. III and H. A. SCHUCK 1960: Growth of bluefin tuna of the western North Atlantic. Fish. Bull. U.S. 61 (179): 39-52.
- MATHER, F. J. III et al. 1974: Result of sailfish tagging in the western North Atlantic Ocean. Proc. Intern. Billfish Symp. II: 194-210. N.O.A.A, Tech. Rep, NMFS SSRF-675.
- MATHEW, C. P. 1974: An account of some methods of overcoming errors in ageing tropical and subtropical fish populations when the hard tissue growth markings are unreliable and the data sparse; pp. 158-166 in: Bagenal (ed.) Ageing of Fish, Unwin Bros. Ltd., Old Woking, Surrey, England, 234 pp.
- MATHISEN, O. and S. OLSEN 1968: Yield isopleths of the halibut Hippoglossus hippoglossus in Northern Norway. Rep. Norw. Fish and Mar. Invest. 14: 129-159.
- MATSUI, I. and M. AMIO 1951: On the age composition and the circuli of the Black Croaker Nibea nibe (Jordan & Thompson). Bull. Jap. Soc. sci. Fish. 16 (12): 168-173.
- MATSUI, I. and T. TAKAI 1951: Ecological studies on the valuable fish in the East China Sea and Yellow Sea. II. Ecological studies on the black Croaker, Nibea nibe (Jordan & Thompson). Bull. Jap. Soc. sci. Fish. 16 (12): 125-143.
- MATSUURA, S. 1961: Age and growth of flatfish Ganzobriame, Pseudorhombus cinamoneus (Temn. & Schl.) Rec. Ocean. Works in Jap. Sp. Nr. 5: 103-110.
- MATTA, F. 1956: L'accrescimento del merluzzo del mediterraneo. Boll. Pesca Pisc. Idrobiol. XI: 31-36.

MATTA, F. 1959: Données préliminaires sur la biométrie de certaines espèces de poissons dans l'Archipel toscan. Proc. Tech. Pap. Gen. Fish. Counc. Médit. 5: 121-134.

MATTHEW, J. 1960: Synopsis on the biology of the South West African Pilchard (Sardinops ocellata Pappé). FAO Symp. II: 115-135.

MAURIN, C. 1954: Les merlus du Maroc et leur Pêche. Bull. Inst. Pêches marit. (Maroc) (2): 7-65.

MAY, A. W. 1959: Cod investigations in Subarea 2 - Labrador 1950 to 1958. I.C.N.A.F. Ann. Proc. 9: 103-105.

MAY, A. W., A. T. PINHORN, R. WELLS and A. M. FLEMING 1965: Cod Growth and Temperature in the Newfoundland Area. I.C.N.A.F. Spec. Publ. No. 6, Environmental Symposium.

MC CRACKEN, F. D. 1954: Seasonal movements of the winter flounder, Pseudopleuronectes americanus (Waldbaum), on the Atlantic Coast. Thesis, Univ. of Toronto, 141 pp.

MC ERLAN, A. 1963: A study of the age and growth of the gag Mycteroperca microlepis Goode and Bean (Pisces, Serranidae) on the West Coast of Florida. Florida Board of Conserv. Tech. Ser. 41: 1-35.

MEEK, A. 1916: The Migration of Fish. Edward Arnold, London, XVIII + 427 pp.

MENGI, T. 1963: Über das Wachstum des Steinbutts (Scophthalmus maximus L.) in der Nordsee. Ber. dt. Wiss. Kommn Meeresforsch. N.F. 7 (2): 119-132.

MENON, M. D. 1950: The bionomics of the poor cod (Gadus minutus L.) in the Plymouth area. J. Mar. Biol. Ass. U.K. 29 (1): 195-239.

- MERETT, N. 1971: Aspects of the biology of billfish (Istiophoridae) from the equatorial western Indian Ocean. J. Zool. Lond. 163: 35-395.
- MESSERSMITH, J. D. 1969: The Northern Anchovy (Engraulis mordax) and its fishery 1965-1968. Calif. Fish Game, Fish. Bull., 147: 102 pp.
- MESSTORFF, J. 1959: Untersuchungen über die Biologie des Wittlings Merlangius merlangus (L.) in der Nordsee. Ber. dt. Wiss. Kommn. Meeresforsch. 15 (4): 277-334.
- MILES, D. W. 1950: The life histories of the spotted sea trout, Cynoscion nebulosus, and the redfish, Sciaenops ocellatus. Annual Rep. for the Fiscal Year Sept. 1, 1949 to August 31, 1950 (Texas), 38 pp. (mimeo).
- MILLER, D. J., A. E. DAUGHTERTY, F. E. FELIN, and J. MACGREGOR 1955: Age and length composition of the northern anchovy catch off the coast of California in 1952-53 and 1953-54. Calif. Dept. Fish Game, Fish. Bull., 101: 37-66.
- MILLER, D. J. and R. S. WOLF 1958: Age and length composition of the northern anchovy catch off the coast of California in 1954-55, 1955-56, and 1956-57. Calif. Dept. Fish Game, Fish. Bull. 106: 27-72.
- MILLER, D. and J. GEIBER 1973: Summary of blue rockfish and lingcod life histories. Calif. Fish Game, Fish. Bull., No. 158, 137 pp.
- MILLER, P. J. 1961: Age, growth and reproduction of the rock goby, Gobius paganellus L. in the Isle of Man. J. Mar. Biol. Ass. U.K. 41: 737-769.

- MILLER, P. J. 1975: Age, structure and life-span in the Common Goby, Pomatoschistus microps. J. Zool. Lond. 175: 425-448.
- MILLER, R. R. 1960: Systematics and biology of the gizzard shad (Dorosoma cepedianum) and related fishes. Fish. Bull., U.S. (173) 60: 371-392.
- MINET, J. P. 1974: Données sur la biologie de la plie américaine Hippoglossoides platessoides des bancs méridionaux de Terre-Neuve et du plateau du Cape Breton. Rev. Trav. Inst. Pêch. marit. 38 (4): 343-434.
- MISU, H. and S. HAMASAKI 1971: Age and growth of Peruvian Hake, Merluccius gayi (Guichenot). Bull. Sehai. Reg. Fish. Res. Lab. (41): 93-105.
- MITANI, F. and E. IDA 1964: A study on the growth and age of the jack mackerel in the East China Sea. Bull. Jap. Soc. Sci. Fish. 30 (12): 958-977.
- MITANI, F. and T. SATO 1959: Studies on the growth and age of the yellow-tail Seriola quinqueradiata (T. & S.) found in Japan and adjacent region II Estimation of age and growth from the opercular bone. Bull. Jap. Soc. Sci. Fish. 24 (10): 803-808.
- MOFLETT, A. 1961: Movements and growth of spotted sea trouts, Cynoscion nebulosus, Cuvier, in West Florida. Fla. St. Bd Cons. Tech. Ser. (36): 1-35.
- MOLLOY, J. 1967: Irish Sprats and Sandeels. Irish Fish. Invest (B), (2): 5-18.

- MOORE, E. 1947: The Sand Flounder, Lophopsetta aquosa (Mitchill): A general study of the species with special emphasis on age determination by means of scales and otoliths. Bull. Bingham Ocean. Coll. 11 (3): 79 pp.
- MOORE, H. L. 1951: Estimation of age and growth of yellowfin tuna (Neothunnus macropterus) in Hawaiian waters by size frequencies. Fish. Bull., U.S. 52 (65): 131-149.
- MOORES, J. et al. 1975: Migrations and biological characteristics of Atlantic Mackerel (Scomber scombrus) occurring in Newfoundland waters. J. Fish. Res. Bd Canada 32 (8): 1347-1357.
- MOREAU, J. 1964: Contribution à l'étude du Lieu Jaune (Gadus pollachius L.). Rev. Trav. Inst. Pêch. marit, 28 (3): 237-255.
- MORGAN, A. and A. GERLACH 1950: Striped bass studies in Coos Bay, Oregon, in 1949 and 1950. Oregon Fish Comm./ Oregon Game Comm. Report to the 46th Legislature, 31 pp.
- MORSUWAN, P. 1970: On the biology of slender trevally, Caranx leptolepis, in the Gulf of Thailand. Marine Fish. Lab., Bangkok MS, 16 pp.
- MOSHER, K. H. 1953: Use of otoliths for determining the age of several fishes from the Bering Sea. J. du Conseil 19: 337-344.
- MOSS, S. 1972: Tooth replacement and body growth rate in the smooth Dogfish, Mustelus canis (Mitchill). Copeia (4): 808-871.



- MOTAIS, R. 1960: Quelques observations sur la biologie d'un poisson abyssal, Trachyrinchus trachyrinchus Risso, et sur les conditions de vie en mer profonde. Bull. Inst. Océanogr., Monaco, (1165): 79 pp.
- MOZZI, C. and A. DUO 1958: Croissance et age des sardines de la Haute Adriatique débarquées à Chioggia. Tech. Pap. gen. Fish. Coun. Médit. 5 (10): 1-15.
- MUKARAMI, S. and S. SHINDO 1949: Studies on the stocks of some economically important marine fishes caught around Amakusa IV. Some observations on the round herring (Temminck and Schlegel). Bull. Jap. Soc. sci. Fish. 15 (2): 122-130.
- MUKARAMI, S. and S. SHINDO 1949: Studies on the stocks of some economically important marine fishes caught around Amakusa V. Consideration on the age of Trachurus japonicus (Temn. & Schl.). Bull. Jap. Soc. sci. Fish. 15 (4): 155-157.
- MUKARAMI S. and S. SHINDO 1950: Studies on the stocks of some economically important marine fishes caught around Amakusa. IV. On the snapper, Pagrosomus major (T. & S.). Bull. Jap. Soc. sci. Fish. 15 (4): 158-160.
- MUKARAMI, S. and K. OKADO 1967: Studies on the Fishery Biology of the Sea Bream, Chrysophrys major Temm. & Schl. in the East China and Yellow Sea. Bull. Sekai Reg. Fish. Res. Lab. (35): 23-40.
- MUZINIČ, R. 1957: Sur la croissance de la jeune sardine (Sardina pilchardus Wald.) dans les eaux de Split. Biljeske, Split, (12).

NAIR, R. V. 1960: Synopsis on the biology of the Indian Sardines. Proceedings of the World Scientific Meeting on the Biology of Sardines and Related Species Vol. 2: 329-414.

NAKAI, Z. 1960: Changes in the population and catch of the Far East Sardine area. In: Proceedings of the World Scientific Meeting on the Biology of Sardines and Related Species, vol. 3: 807-853.

NAKASHIMA, E. 1916: Cynoscion macdonaldi Gilbert. Copeia (31): 85-86.

NALL, G. 1927: Fisheries, Scotland, Salmon Fish, II (as cited in Beverton & Holt 1959),

NAMMALWAR, P. 1974: A note on determination of age from skeletal structure of Pomadasys hasta (Bloch). Ind. J. Fish. 21 (1): 294-296.

NAMURA, H. 1966: Idade e crescimento da Pescada - Branea Cynoscion leiarchus (Cuvier), das águas cearenses. Arq. Est. Biol. Mar. Univ Fed. Ceara 6 (2): 135-137.

NATARAJAN, A. V. and V. G. JHINGRAN 1963: On the biology of Catla catla (Ham.) from the River Yamuna. Ind. J. Fish. 8 (1): 54-59.

NAVARRO F. de P. 1932: Nuevos estudios sobre la alacha (Sardinella aurita C. V.) de Baleares y de Canarias. Notas Inst. esp. Oceanogr. 58: 35 pp.

NEDELEC, C. 1958: Biologie et Pêches des Maqueraux. Rev. Trav. Inst. Pêch. marit. 22 (2): 121-134.

- NEWMAN, G. 1970: Stock assessment of the pilchard Sardinops ocellata at Walfis Bay, South West Africa. Div. Fish. Invest. Rep. (85): 1-17.
- NICHOLLS, G. 1973: Growth of the Australian "Salmon" Arripis trutta (Bloch & Schneider). Austr. J. Mar. Freshw. Sci. 24 (2): 159-176.
- NIKOLOV, D. K. 1960: [Biology of the bonito Sarda sarda (Bloch) from the Black Sea.] Trud. nauch.-izsled. Inst. Rib. Prom. Varna, 3: 91-115 (In Bulgarian).
- NIKOLSKY, G. W. 1957: Spezielle Fischkunde. VEB Deutscher Verlag der Wissenschaften: Berlin, 632 pp.
- NIVAL, S. 1966: Contribution à l'étude biologique de Symphodus rostratus Bl. (Pisces: Labridae) en Rade de Villfranche-sur Mer. Vie et Milieu 17 (3)A: 1273-1315.
- NOMURA, H. 1967: Dados biológicos sobre a serra Scomberomorus maculatus (Mitchill) das águas caerenses. Arq. Est. Biol. Mar. Univ. Fed. Caera 7 (1): 29-39.
- NOSE, Y., H. KAWATSU and Y. HIYAMA 1957: Age and growth of Pacific tunas by scale reading. In: Suisan Gaku Shusei, Tokyo Univ. Press, pp. 701-716.
- NÜMANN, W. 1955: Die Pelamiden des Schwarzen Meeres, des Bosporus, der Marmara und der Dardanellen (Sarda sarda). Hidrobiol., Istanbul, 3 (2-3): 75-127.
- O'CONNEL, C. P. 1953: The life history of the cabezon, Scorpaenichthys marmoratus, (Ayres). Calif. Fish Game, Bull. No. 93, 76 pp.

- ODATE, S. 1966: Studies on the fishes of the family Myctophidae in the Northeastern Sea of Japan. III. The determination of age and growth of Susuki-Hadaka, Myctophium affine (Lüdken). Bull. Tohoku Reg. Fish. Res. Lab. 26: 35-43. See also 19: 90-97.
- OHSIMA, Y. and N. NAKAMURA 1941: Notes on the life history of Triacanthus brevirostris Temminck and Schlegel. Bull. Jap. Soc. sci. Fish., 10 (4): 171-176.
- OLIVIER, R. 1928: La dorade (Pagellus centrodontus) (résumé pratique de nos connaissances sur ce poisson). Rev. Trav. Off. Pêches marit. 1 (4) p. 5.
- OLSEN, A. M. 1954: The biology, migration and growth rate of the school Shark Galeorhinus australis (McLeay), Charcharhinidae, in South Eastern Australian Waters. Austr. J. Mar. Freshw. Res. 5 (3): 353-410.
- OLSEN, S. 1954: South Georgian Cod. (Notothenia rossii marmorata Fischer). Norsk Havkangst-tid 7: 373-382.
- OLSEN, S. 1955: A contribution to the systematics and biology of Chaenichthid fishes from South Georgia. Nytt Magasin Zoologi 3: 79-93.
- OLSEN, Y. and D. MERRIMAN 1946: The biology and economic importance of the ocean pout Macrozoarces americanus (Bloch and Schneider). Bull. Bingham Ocean. Coll., 9 (4): 184 pp.
- ORCUTT, H. 1950: The life history of the Starry Flounder Plathyichthys stellatus (Pallas). Calif. Fish Game, Fish. Bull., (78) 64 pp.

- OSHIMA, Y. 1942: An ecological note on the fish Sparus swinhonis Günther. Bull. Jap. Soc. sci. Fish. 10 (6): 249-255.
- ØSTVEDT, O. J. 1964: On the life history of the spotted catfish (Anarhichas minor Olafsen), Fish. Dir. Skr. 13 (6): 54-72.
- OTSU, T. 1960: Albacore migration and growth in the North Pacific Ocean as estimated from tag recoveries. Pacific Sci., 14 (3): 257-266.
- PACHECO, A. 1962: Age and growth of Spot in lower Chesapeake Bay, with notes on Distribution and Abundance of Juveniles in the York River System. Chesapeake Sci. 3 (1): 18-28.
- PADOAN, P. 1963: Prime osservation sulle accinghe (Engraulis encrasicolus L.) catturate al largo delle foci del Po. I.C.E.S. Rapp. & Proc. Verb. 17 (2): 327-332.
- PANTULU, R. 1962: On the use of Pectoral spines for the determination of age and growth of Pangasius pangasius (H. & S.) J. du Conseil 27 (1): 192-216.
- PEARLMUTTER, A. and G. M. CLARKE 1949: Age and growth of immature rosefish (Sebastes marinus) in the Gulf of Maine and off Western Nova Scotia. Fish. Bull., U.S., 51 (45): 207-228.
- PEARSON, J. 1929: Natural history and conservation of redfish and other commercial Sciaenidae of Texas. Bull. U.S. Bur. of Fish. 44, doc 1046: 129-214.

- PHAM-THUOC and J. SZYPULA 1975: Age and growth rate of gilt Sardine (Sardinella aurita Cuv. et Val.) from North East African Waters. I.C.E.S. CM, 1975/J:5 Pel. Fish (Southern) Committee.
- PHETTONGKAM, M. and W. THASANANULKIT 1972: On the biology of Sciaena russelli (Cuvier et Valenciennes) in the inner Gulf of Thailand. 3rd. Symposium on Marine Fisheries, Mar. Fish. Lab. Bangkok, M.S. 18 pp.
- PHILLIPS, J. B. 1948: Growth of the sardine Sardinops caerulea 1941-42 through 1945-47. Fish. Bull., Sacramento 71: 33 pp.
- PILLAY, S. R. and H. ROSA, Jr. 1963: Synopsis of Biological Data on Hilsa ilisha (Ham.) 1822. FAO Fisheries Biology Synopsis No. 25.
- PILLAY, T. V. R. 1958: Biology of the hilsa, Hilsa ilisha of the River Hoogly. Ind. J. Fish. 5 (2): 201-257.
- PITT, T. 1975: Changes in abundance and certain biological characteristics of Grand Bank American Plaice, Hippoglossoides platessoides. J. Fish. Res. Bd Can. 32 (8): 1383-1398.
- POINSARD, F. 1973: Croissance des Pseudotolithus typus Blk dans la région de Pointe Noire. Doc. Centre O.R.S.T.O.M. de Pointe Noire.
- POINSARD, F. and J. TROADEC 1966: Détermination de l'âge par la lecture des otolithes chez deux espèces de Sciaenides ouest africain (Pseudotolithus senegalensis C. V. et Pseudotolithus typus Blkr.). J. du Conseil 30 (3): 291-307.

- POPIEL, J. 1958: The growth rate of the Southern Baltic Herring. I.C.E.S. C.M. 1958/Herring Comm. (48) 8 pp.
- PORCHE, P. 1976: Le sprat (Clupea sprattus L.) du golfe de Gascogne. Observations biologiques et écologiques sur les échantillons prélevés en 1975. I.C.E.S. C.M. 1976/ H:9.
- POSTEL, E. 1955 a: Contribution à l'étude de la biologie de quelques Scombridae de l'Atlantique tropico-oriental. Ann. Stn. océanogr. Salammbô 10: 168 p.
- POSTEL, E. 1955 b: Résumé des connaissances acquises sur les Clupéidés de l'Ouest Africain. Rapp. Cons. Explor. Mer, 137: 14-17.
- POSTEL, E. 1956: Données biométriques sur quelques Scombridés tunisiens. Bull. Sta. océanogr. Salammbô, (53): 50-61 (SS/1a).
- POULSEN, E. 1935: Recent investigations into the stock of salmon and sea trout in the River Gudenaa. Rep. Danish biol. Stn. 40.
- PRABHU, M. 1955: Some aspects of the biology of ribbon fish, Trichiurus haumela (Forsk.) Ind. J. Fish. 2 (1): 132-163.
- PRIOL, E. P. 1944: Observations sur les germons et les thons rouges capturés par les pêcheurs bretons. Rev. Trav. Off. Pêches Marit. 13 (1-4): 384-439.
- PRUTER, A. 1954: Age and growth of the Oregon sablefish, Anoploma fimbria. Pacific Marine Fisheries Comm. Bull. 3: 121-128.

- QASIM, S. Z. 1957: The biology of Blennius pholis L. (Teleostei). Proc. Zool. Soc. Lond. 128: 161 p.
- QUIGNARD, J. P. 1973: Recherche sur la biologie d'un sélacien du Golfe de Tunis, Torpedo torpedo Linné 1978 (Croissance relative, croissance absolue, coefficient de condition). Ann. Inst. Michel Pacha 6: 72-110.
- RADHAKRISHNAN, N. 1957: A contribution to the biology of Indian Sand Whiting - Sillago sihama (Forsk.) Ind. J. Fish. 4 (2): 254-283.
- RADHAKRISHNAN, N. 1964: Notes on some aspects of the biology of the fringe scale sardine, Sardinella fimbriata (C. and V.). Ind. J. Fish. 11 A (1): 127-134.
- RAFAIL, S. Z. 1972: Investigation on Sciaenidae and Moronidae catch and on the total catch by beach seine on U.A.R. Mediterranean Coast. Coun. Gen. Peches Medit. Studies & Reviews (48): 1-26.
- RAFAIL, S. Z. 1972a: Red Sea fisheries by light and purse-seine near Al-Ghardaqa. Bull. Inst. Ocean. Fish (Cairo) 2: 25-49.
- RAFAIL, S. Z. 1972 b: A statistical study of length-weight relationship of eight Egyptian fishes. Bull. Inst. Ocean. Fish. (Cairo) 2: 136-156.
- RAFAIL, S. Z., A. ALKOHOLY and M. MALIDI 1973: Biological studies on Sundanese Inland fishes. III. Hydrocyon forskalii Cuvier. Bull. Inst. Ocean. Fish. (Cairo), 3: 276-293.



- RAITT, D. 1966: The biology and commercial potential of the blue whiting (Micromesistius poutassou) in the North East Atlantic. I.C.E.S. Symp. Ecology of Pelagic fishes in Arctic waters, 1966.
- RAITT, D. 1968: The population dynamics of the Norway pout in the North Sea, Mar. Res. (Scotland) (5) 24 pp.
- RAITT, D. and V. SAGUA 1969: A preliminary investigation on the biology of Brachydeuterus auritus (Val. 1831), in Nigerian waters, pp. 397-401, in: UNESCO Symp. Trop. Atlantic, Abidjan, 430 p.
- RAJAN, S. 1964: The biology and fishery of Pseudosciaena (Hamilton) from the Chilka Lake. Ind. J. Fish. 11 (2): 639-663.
- RANDALL, J. E. 1961: A contribution to the biology of the convict surgeonfish of the Hawaiian Islands: Acanthurus triostegus sandvicensis. Pacific Science 15 (2): 215-272.
- RANDALL, J. E. 1962: Tagging Reef Fishes in the Virgin Islands. Proc. Gulf & Caribb. Fish Inst., 14: 201-241.
- RANNOU, M. 1976: Age et croissance d'un poisson bathyal: Nezumia glerorhynchus (Macrouridae, Gadiformes) de la Mer d'Alboran. Cah. Biol. Mar. 17 (4): 413-421.

- RAO, V. S. 1961: Studies on the age determination of "Ghol" Pseudosciaenia diacanthus (Lacépède) by means of scales and otoliths. Ind. J. Fish. 8 (1): 121-126.
- RAO, V. S. 1966: Age and growth of "Ghol", Pseudosciaenia diacanthus (Lacépède), in Bombay and Saurashtra Waters. Ind. J. Fish. 13 (1-2): 251-291.
- RAWSON, D. S. 1956: The life history of the yellow wall-eye, Stizostedion vitreum, in Lac La Ronge, Saskatchewan. Trans. Am. Fish. Soc. 86: 15-37.
- REAY, P. 1970: Synopsis of biological data on North Atlantic sandeels of the genus Ammodytes. FAO Species Synopsis (82) pag. var.
- REAY, P. 1973: Some aspects of the biology of the sandeel, Ammodytes tobianus L. in Langstone Harbour, Hampshire. J. Mar. Biol. Ass. U.K. 53: 325-346.
- REINTJES, J. 1969: Synopsis of biological data on the atlantic Menhaden Brevoortia tyrannus. FAO Fish. Synopsis No. 42 and U.S. Fish. Wildlife Serv. Circ. 320, 30 pp.
- REYNOLDS, L. F. and R. MOORE 1973: Barramundi (Lates calarifer) research in Papuan waters, 21-28 in: Annual Report, Dept. of Agriculture, Stock and Fisheries, Port Moresby, 51 pp.
- RICHARD, C. E. 1967: Age, growth and fecundity of the cobia, Rachycentron canadum, from Chesapeake Bay and adjacent Mid-Atlantic Waters. Trans. Am. Fish. Soc. 96 (3): 343-350.

- RICHARD, C. E. 1972: Age, growth and distribution of the black drum (Pogonias cromis). Trans. Am. Fish. Soc. 102 (3): 584-590.
- RICHARD, S., D. MERRIMAN and L. CALHOUN 1963: The biology of the Little Skate Raja erinacea ~~Mitchill~~. Bull. Bingham Ocean. Coll. 18 (3): 5-67.
- RICHARDSON, I. D. et al. 1960: Report on sardine investigations in Brazil. FAO Proceeding of the World Scientific Meeting on the Biology of Sardines and Related Species Vol. III, 667-791.
- RICKER, W. E. 1958: Handbook of computations for biological statistics of fish populations. Bull. Fish. Res. Bd Can., (119): 300 p.
- RIVAJEC, L. 1973: Biology and Dynamics of Pagellus coupei (Dieuz. 1960), Pagrus ehrenbergi (Val. 1830) and Dentex canariensis (Poll. 1954) in Ghana Waters. Doc. Sci. Centr. Rech. Ocean. Agidjan O.R.S.T.O.M. 14: 42-97.
- ROBB, A. P. 1973: Age, growth and stock assessment of the Megrin Lepidorhombus whiffiagonis (Wald.) in Northern Scottish Waters. Diss. Aberdeen 1973, 75 pp.
- ROBINS, J. P. 1963: Synopsis of the biological data on bluefin tuna Thunnus thynnus maccoyii (Castelnau) 1872. FAO Fish. Rep. (6), 2: 562-587.
- RODRIGUEZ-PINO, Z. 1962: Estudios estadísticos y biológicos sobre la biajaiba (Lutjanus synagris). Centro de Investigaciones Pesqueras. Nota Sobre Investigaciones No. 4, 91 pp. (Cuba).

- RODRIGUEZ-RODA, J. 1972: Research on Bluefin Tuna (Thynnus thynnus L.) in Spain during the years 1970-1971: pp. 193-194 in: Report Int. Comm. Cons. Atl. Tuna, 1970-71, III. Madrid.
- RODRIGUEZ-RODA, J. and M. G. LARRAÑETA, 1955: El crecimiento de la sardina (Sardina pilchardus Wald.) de las costas de Alicante, Invest. pesqu., (2): 9-20.
- ROJO, A. and J. SILVOSA 1969: Estudio biológica de la merluza (Merluccius merluccius hubbsi) del sector patagonico. FAO Argentino-Proyecto de Desarrollo Pesquero Publicacion No. 18, 40 pp.
- ROLLEFSEN, G. 1934: The cod otolith as a guide to race, sexual development and mortality. I.C.E.S. Rapp. & Proc. Verb. 88 (2).
- RONQUILLO, I. A. 1960: Synopsis of biological data on Philippine Sardines (Sardinella perforata, S. fimbriata, S. sirm, S. longiceps). In: Proceedings of the World Scientific Meeting on the Biology of Sardines and Related Species, Species Synopsis 13.
- ROSEN, N. 1918: Om Havslaxöringen i. Övre Norrland. Medd. Kungl. Lantbruksstyrelsen (212).
- ROSSIGNOL, M. 1955: Premières observations sur la biologie des sardinelles dans la région de Pointe Noire (Sardinella eba Val., S. aurita Val.). Rapp. Cons. Explor. Mer, 137: 17-21.
- ROTHSCHILD, B. 1967: Estimates of the growth of the skip-jack tuna (Katsuwonus pelamis) in the Hawaiian Islands. Proc. Indo-Pacif. Fish. Comm. 12 (II): 100-111.

- ROUNSEFELL, G. A. 1930: Contribution to the biology of the pacific herring Clupea pallasii and the conditions of the fishery in Alaska. U.S. Bur. of Fish. Doc. No. 1080: 227-320.
- RUSH, W. 1952: Observations of age and growth in the Menhaden (Brevoortia tyrannus) as determined by scale examination. Copeia 1952 (3): 208-209.
- RUTKOWICZ, St. 1963: The results of investigations on the stock of cod (Gadus morhua L.) in the southeastern area of the Baltic Sea in the years 1946-1960. Prac. Morsk. Inst. Ryb. 12A: 165-211.
- SAEGER, J. 1974: Der Befischungszustand der Flunderpopulation in der Kieler Bucht. Thesis, Kiel University, 115 pp.
- SAEMUNDSSON, B. 1923: On the age and growth of the cod (Gadus callarias L.) in Iceland waters. Medd. Fisk. Havundersøg., Ser. Fisk. 7 (3): 1-35.
- SAEMUNDSSON, B. 1925: On the Age and Growth of the Haddock (Gadus aeglefinus L.) and the Whiting (Gadus merlangus L.) in Icelandic waters. Medd. Komm Havundersøg., Ser. Fisk. 8 (1): 1-26.
- SAEMUNDSSON, B. 1926: Fiskarnir (Pisces Islandiae) in Iceland. Reykjavik, 583 pp., here: 182-189.
- SAEMUNDSSON, B. 1929: On the age and growth of the coal-fish (Gadus virens L.), the Norway Cod (Gadus esmarki Nilsson) and the Poutassun (Gadus poutassou Risso) in Icelandic Waters). Medd. Komm. Havundersøg., Ser. Fisk. Bd. 8 (7): 1-37.

SAETERSDAL, G. and L. VILLEGAS 1968: Estudio del tamaño, crecimiento y madurez de la merluza (Merluccius gayi) en aguas chilenas. Publ. Inst. Fom. Pesq. (34). 52 pp.

SAGE, M. et al. 1972: Growth and seasonal distribution of the elasmobranch Dasyatis sabina. Contr. Mar. Sci. (Texas) 16: 71-74.

SANCHEZ, R. J. E. 1966: General aspects of the biology and ecology of the anchovy (Engraulis ringens). Proc. Gulf & Carib. Fish. Inst. 18: 84-93.

DOS SANTOS, E. P. 1963: Growth of "Goete": Quantitative aspects. Bol. Inst. Ocean (São Paulo) 13 (1): 185-190.

SARHAGE, D. 1970: Ein Beitrag zur Biologie des Stückers (Trachurus trachurus L.) in der Nordsee. Bericht dt. wiss. Kommn Meeresforsch. N. F. 21 (1-4): 122-169.

SAROJINI, K. K. 1957: Biology and Fisheries of the Grey Mullet of Bengal. I. Biology of Mugil parsia Hamilton with notes on its fishery in Bengal. Ind. J. Fish. 4 (2): 160-207.

SAROJINI, K. K. 1958: Biology and fisheries of grey mullets of Bengal. II. Biology of Mugil cunnesius Valenciennes. Ind. J. Fish 5 (1): 56-74.

SARRATIMSKY, P. I. 1973: The age of the rock grenadier in the North West Atlantic and a possible influence of fisheries on its population number. Tranl. Serv. Res. Board Canada 2491, 26 pp.

SCHAEFER, M. B. 1961: Rep. Inter-American Tropical Tuna Comm., 1960: 40-183.

SCHMIDT, U. 1959: German investigations on commercial saith landings in 1958/59. Appendix in: Stat. Newslett. I.C.E.S. (8) Dem. Sp. 1957-58, pp. 37-42.

SCHROEDER, W. 1930: Migrations and other phases in the life history of the cod off Southern New England. Bull. U.S. Bur. Fisheries 46: 1-136. (Doc. 1081).

SCHWARZ, F. J. 1965: Age, growth and egg complement of the stickleback Apeltes quadracus at Solomons, Maryland. Chesapeake Sci. 6 (2): 116-118.

SCHWARZ, F. J. 1971: Biology of Microgobius thalassinus Pisces: Gobiidae, a sponge inhabiting goby of Chesapeake Bay, with range extension of two goby associates. Chesapeake Sci. 12 (3): 156-166.

SCHWARZ, F. J. and B. DUTCHER 1963: Age, growth and food of the oyster toadfish neas Solomons, Maryland. Trans. Am. Fish. Soc. 92: 170-173.

SCOFIELD, E. C. 1931: The striped bass of California (Roccus lineatus). Div. Fish & Games of Calif. Fish Bull. 29, 82 pp.

SCOTT, J. S. 1968: Morphometrics, distribution, growth and maturity of offshore sand launce (Ammodytes dubius) on the Nova Scotia Banks. J. Fish. Res. Bd. Can. 25 (9): 1775-1785.

SEKARAN, K. V. 1955: Observations on the choodal fishery of Mandapam area. Ind. J. Fish. 2 (1): 113-131.

- SEKERAN, DENEVASAN & JOHN (1940) cited without further reference) in Table I, p. 1963 by: Banerji S. K. and T. S. Frishnan 1973. Acceleration of assessment of fish populations and comparative studies of similar taxonomic groups, p. 158 - 175 in: Proc. Symp. Living Res. around India. Spec. Publ. Centr. Mar. Fish. Res. Inst. Cochin, India, 748 pp.
- SELLA, M. 1930: Distribution and migrations of the tuna (Thunnus thynnus L.) studied by the method of hooks, and other observations. Int. Rev. Hydrobiol., 24: 446-466.
- SEMAKULA, S. and P. LARKIN 1968: Age, growth and yield of the white sturgeon (Acipenser transmontanus) of the Fraser River, British Columbia. J. Fish. Res. Bd. Can. 25 (12): 2589-2602.
- SERCHUK, F. and C. COLE 1974: Age and growth of the cunner, Tautogalabrus adspersus (Waldbaum), Pisces (Labridae), in the Weweeantic River Estuary, Massachusetts. Chesapeake Sci. 15 (4): 205-213.
- SERVENTY, D. L. 1956: Additional observations on the biology of the northern bluefin tuna, Kishinoella tonggol (Bleeker) in Australia: Aust. J. Mar. Freshw. Res., 7 (1): 44-63.
- SESHAPPA, G. 1958: Occurrence of growth cheeks in the scales of Indian Mackerel, Rastrelliger kanagurta (Cuvier). Curr. Sci. 27: 262-263.
- SHINDO, S. 1972: Note on the study on the stock of Lizard fish, Saurida tumbil in the East China Sea. Proc. IPFC 13 (III): 298-305.



- SHIOKAWA, T. 1962: Growth and maturity of the common Rockfish, Sebasticus marmoratus Cuvier et Valenciennes. Rec. Ocean. Works in Jap. (Spec. No. 6).
- SHOMURA, T. 1962: Studies on habits of coastal fish in the Amakusa Islands. Part II. Growth and maturity of the purple rockfish Sebastes pachycephalus Temminck & Schlegel. Rec. Ocean. Works in Jap. (Sp. No. 6): 103-111.
- SHOMURA, R. S. and B. A. KEALA 1963: Growth and sexual dimorphism in growth of bigeye tuna (Thunnus obesus). A preliminary report. FAO Fish. Rep. (6) 2: 1409-1417.
- SIBLOT-BOUTEFLIKA 1976: Contribution à l'étude des Scorpenidés de la Région d'Alger. Thesis, Univ. of Marseille.
- SILLIMAN, R. P. 1943: Studies on the Pacific pilchard or Sardine (Sardinops caerulea). 5. A method of computing mortalities and replacement. Spec. sci. Rep. U.S. Fish wildl. Serv., (24): 10 pp.
- SIMPSON, J. G. 1971: The present status of exploitation and investigation of the clupeoid resources of Venezuela. Fish. Rep. FAO No. 71-2: 263-278. Caribbean Symposium.
- SIMPSON, J. G. and R. GRIFFITHS 1973: Age, growth and maturity of the "rabo amarillo", Cetengraulis edentulus, in Eastern Venezuela, based on length frequency studies. Offic. Nacional de Pesca, Venezuela Ser. Recurs. y Expl. Pesqu. 2 (6).
- SIMPSON, J. G. and R. BUZETA 1967: El crecimiento y la edad de la anchoveta (Engraulis ringens, Jenyns) en Chile basado en estudios de frecuencia de longitud. Inst. Fomento Pesq., Boletín Cien., 3: 53 pp.

- SIVASUBRAMANIAM, K. 1973: Co-occurrence and the relative abundance of narrow and broad corseletted frigate mackerel, Auxis thazard (Lacépède) and Auxis rochei (Risso) around Ceylon: 537-547. In: Proc. Symp. Living Res. Seas around India, Spec. Publ. Cen. Mar. Fish. Res. Inst. Cochin, 748 pp.
- SKILLMAN, R. and M. YOUNG 1976: Von Bertalanffy growth curves for striped marlin, Tetrapterus audax and blue marlin, Makaira nigricans in the North Central Pacific. (US) Fish. Bull. 74 (3): 553-566.
- SMIDT, E. 1969: The Greenland halibut, Rheinhardtius hippoglossoides (Waldb.) Biology and exploitation in West-Greenland. Meddelser fra Danmarks Fiskeri-og Havunderøgelser 6 (4): 79-148.
- SMYLY, W. J. P. 1957: Proc. Zool. Soc. London 128, 431.
- VAN SOMEREN, V. D. and P. J. WHITEHEAD 1960: The culture of Tilapia nigra (Günther) in ponds. IV. The seasonal growth of male T. nigra. E. African Agric. & For. J. 26 (2): 79-86.
- SOMJAIWONG, D., CHULLASORN and SUPONGPAN 1972: Assessment on growth and mortality of Indo-Pacific Mackerel (Rastrelliger neglectus van Kampen) from tagging data in the Gulf of Thailand. M.S. 1st Seminar of Mar. Res. Lab., Bangkok.
- SOUTHWARD, G. and D. CHAPMAN 1965: Utilization of Pacific halibut stocks: study of Bertalanffy's growth equation. Rep. Int. Pac. Halibut Comm. 39: 33 pp.

- SPRINGER, V. and A. MCERLAN 1961: Spawning seasons and growth of the code goby, Gobiosoma robustum (Pisces: Gobiidae) in the Tampa Bay area.. Tulane Studies in Zoology 9: 77-85.
- STANEK, E. 1962: The stock of the cod in the southern part of the Bornholm Basin in the years 1955-56. Prac. Morsk. Inst. Ryb. (11)A.
- STAPLES, D. 1972: Growth of the red gurnard (Teleostei, Triglidae) from Pegasus Bay, Canterbury, New Zealand. N. Zealand J. Mar. Freshw. Res. 6 (3): 365-374.
- ST. PIERRE, R. and J. DAVIS 1972: Age, growth and mortality of the white perch, Morone americana, in the James and York River, Virginia. Chesapeake Science 13 (4): 272-281.
- STEINITZ, H. 1959: Dr. H. Lissner's study of the biology of Acanthobrama terrae-sanctae in Lake Tiberia. Sea Fishery Res. Station (Israel) Bulletin (24), 64 pp.
- STEVEN, G. 1952: Contribution to the biology of the mackerel, Scomber scombrus. III. Age and growth. J. Mar. Biol. Ass. U.K. 30: 549-568.
- STEVEN, G. and P. CORBIN 1939: Mackerel investigation at Plymouth. Preliminary report. Rapp. Cons. Explor. Mer. 11: 15-18.
- STEVENS, J. D. 1975: Vertebral rings as a means of age determination in the blue shark (Prionace glauca L.). J. Mar. Biol. Ass. U.K. 55: 657-665.

- STOBO, W. and J. HUNT 1974: Mackerel biology and history of the Fishery in Subarea 4, I.C.N.A.F. Fish. Res. Doc. 74/9 (3155).
- STOYANOV, S. 1961: [Condition of the Anchovy stock of the Black Sea fished near the Bulgarian Coast during the period 1952-1960]. In Bulgarian, Inst. Centr. Rech. Sci. Pisc. Pech. Varna, Bull., 1:1-57.
- STRAWN, K. 1958: Life history of the pygmy sea-horse, Hippocampus zosterae Jordan & Gilbert, at Cedar Key, Florida. Copeia 1958 (1): 16-22.
- STRUHSAKER, P. and J. UCHIYAMA 1976: Age and growth of the nehu Stolephorus purpureus (Pisces, Engraulidae), from the Hawaii Islands as indicated by daily growth increment of sagittae. Fishery Bull, 74 (1): 9-17.
- SUCONDHARMAN, P., C. TANTISAWETRAT and U. SRIRUANGCHEEP 1970: Estimation of age and growth of Chub mackerel Rastrelliger neglectus (van Kampen) in the Western Gulf of Thailand, pp. 471-480. In: Marr, J. C. (ed.). The Kuroshio: A Symposium on the Japanese Current, East West Center Press, Honolulu, 614 pp.
- SUDJASTANI, T. 1974: The species of Rastrelliger in the Java Sea, their taxonomy, morphometry and population dynamics. Thesis, Univ. British Columbia, 147 pp.
- SUNADA, J. 1974: Age and growth of the pacific saury, Cololabis saira. Calif. Fish Game 60 (2): 64-73.
- SUND, O. 1911: Undersøkelser over brislingen i norske farvand vaesentlig paa grundlag av "Michael Sars" togt i 1908. Årsberetn. Norg. Fisk. 1910, Hft. 3: 357-474.

SUNDARAJ, B. 1960: Age and growth of the Spot, Leiostomus xanthurus Lacépède. Tulane Studies in Zoology 8 (2): 41-62.

SVÄRDSON, G. 1943: Studien über den Zusammenhang zwischen Geschlechtsreife und Wachstum bei Lebistes. Mitt. Anst. f. Binnenfischerei bei Drottingholm, Stockholm (21) 48 pp.

SVETOVIDOV, A. N. 1964: [Handbook of the Fauna of the USSR, Fishes of the Black Sea] in Russian. Izdatel'stvo Nauka, Moscow: 550 pp.

SWEDMARK, M. 1957: Variations de la croissance et de la taille dans différentes populations du Téléostéen Gobius minutus. (Communication faite au colloque de Biologie Marine à la Station Biologique de Roscoff, 1956). L'Année Biologique 33 (3-4): 163-170.

TALBOT, F. H. and M. J. PENRITH 1963: Synopsis of biological data on species of the genus Thunnus (sensu lato) (South Africa). FAO Fish. Rep., (6)2: 608-646.

TAMPI, P. 1960: Utilization of saline mud flat for fish culture: an experiment in marine fish farming. Ind. J. Fish. 7 (1): 137-146.

TANAKA, S. 1956: A method of analysing the polymodal frequency distribution and its application to the length distribution of Porgy, Taius tumifrons (T & S.). Bull. Tokai Reg. Fish. Res. Lab. (14): 1-12.

TEMPLEMAN, W. 1948: The life history of the capelin (Mallotus villosus Müller) in Newfoundland Waters. Bull. Nfld. Gov. Lab. (17): 1-151.

- THAM AH KOW 1967: A contribution to the study of the growth of members of the genus Stolephorus Lacépède in Singapore Strait. I.P.F.C, Proc. 12 (2); 1-25.
- THOMAS, J. 1968: Management of the white seabass (Cynoscion nobilis) in California waters. Calif. Fish Game Bull. 142, 34 pp.
- THORPE, J. 1974: Estimation of the number of brown trout, Salmo trutta (L.), in the Loch Leven, Kinross, Scotland. J. Fish. Biol. 6: 135-152.
- THUROW, F. 1973: Growth Parameters of Baltic Salmon (Salmo salar L.). Ber. dt. wiss. Kommn Meeresforsch. 22 (4): 445-451.
- TIEWS, K. 1971: Further results of studies on the spawning stock of cod in the middle Baltic Sea. I.C.E.S. C.M, 1971/ special meeting on cod and herring in the Baltic, cod No. 3, pp. 1-33.
- TIEWS, K. and P. CACES-BORJA 1965: On the Availability of Fish of the Family Leiognathidae Lacépède in Manila Bay and San Miguel Bay and their Accessibility to Controversial Fishing Gears, Phil. Jour. Fish. 7 (1): 59-86.
- TIEWS, K., A. MINES and I. A. RONQUILLO 1972: On the Biology of Saurida tumbil (Bloch 1801), Family Synodontidae in Philippine Waters. Proc. I.P.F.C. 13 (III): 100-120.
- TIEWS, K., I. A. RONQUILLO and L. M. SANTOS 1970: On the biology of Anchovies (Stolephorus Lacépède) in Philippine waters. Proc. I.P.F.C. 13 (II): 20-48.

- TIEWS, K., I. A. RONQUILLO and P. CAECES-BORJA 1970:  
On the biology of Roundscads (Decapterus Bleeker)  
in the Philippine Waters. Proc. I.P.F.C. 13 (11):  
82-106.
- TOKAI LABORATORY 1960: (:Tokai Regional Fisheries Research  
Laboratory, Tokyo). Synopsis on the biology of Sar-  
dinops melanostica. Proceedings of the World Scien-  
tific Meeting on the Biology of Sardines and Re-  
lated Species. Vol. 2: 213-244.
- TOOR, H. S. 1964: Lethrinus lentjan, Lacépède, from  
Indian Waters. III. age and growth. Ind. J. Fish.  
11 (A)(2): 597-620.
- TRAVIN, V. I. 1951: A new species of sea perch in the  
Barents Sea (Sebastes mentella travin sp. nov.).  
Akad. Nauk S.S.S.R. Doklady 77 (4):741-744.
- TROADEC, J. 1969: Observations sur la biologie et la  
dynamique des Pseudotolithus senegalensis dans la  
Région de Pointe Noire. PP. 365-380. UNESCO Symp.  
Oceanogr. Fish. Res. Trop. Atlantic Abidjan.
- TROUT, G. C. 1954: Otolith growth of Barents Sea Cod.  
I.C.E.S. Rapp. & Proc. Verb. 136: 89-102.
- TSUDA, R. T., P. G. BRYAN, W. J. FITZGERALD and W. J.  
TOBIAS 1974: Juvenile - adult rearing of Siganus (Pisces:  
Siganidae) in Guam. South Pacific Commission Seventh  
Technical Meeting on Fisheries, Nuku'alofa, Tonga,  
July 15-19, 1974 (mimeo), 6 pp.
- TUGGAC, M. 1957: On the biology of the Scomber colias  
Gmelin. Tech. Pap. (21). Gen. Fish. Counc. Medit. (4):  
145-159.

- TÜRGAN, G. 1958: The age determination of bonitos and pelamids. *Balik Balıkcılık*, 6 (3): 18-20 (In Turkish).
- TURNER, C., E. EBERT and R. GIVEN 1969: Man made reef ecology. *Calif. Fish Game, Fish. Bull.* 146.
- UCHIYAMA, H. and J. W. STRUHSACKER 1975: Age and growth of Skipjack tuna, Katsuwonus pelamis, Yellowfin tuna, Thunnus albacares and Albacore, Thunnus alalunga, as indicated by daily growth increment of sagitta. *Doc. ICCAT, SCRS/75/57*.
- URSIN, E. 1967: A mathematical model of some aspects of fish growth, respiration, and mortality. *J. Fish. Res. Bd Canada* 24 (11): 2355-2453.
- VALOZZER, G. 1962: Sobre a biologia da Corvina da Costa Sul do Brasil. *Bd. Inst. Ocean. (Sao Paulo)* 12 (1): 53-102.
- VAN DER ELST, R. 1976: Game fish of the east coast of South Africa. I. The biology of the elf, Pomatomus saltatrix (Linnaeus), in the coastal waters of Natal. *Ocean. Res. Inst. Invest. Rep.* (4-4): 1-59.
- VAN MULLEM, P. J. and VAN DER VLUGT 1964: On the age, growth and migration of the anadromous stickleback Gasteosteus aculeatus L. investigated in mixed populations. *Arch. Neerl. Zool.* 16 (1) 1964: 111-116.
- VENKATAMARAN, G. 1956: Studies on some aspects of the biology of the common anchovy, Thrissoclos mystax (Bloch & Schneider). *Ind. J. Fish.* 3 (2): 311-333.
- VERGHESE, T. 1961 a: Observations on the biology of Raconda russellina (Gray). *Ind. J. Fish.* 8 (1): 96-106.



- VERGHESE, T. 1961 b) Some observations on the biology of Coila borneesis (Blkr.). Ind. J. Fish. 8 (2): 312-325.
- VESCHEZEROV, 1944: [Materials on the biology and fisheries of marine perches in the Barents Sea] in Russian. Knipovitch Polar Scientific Institute of Sea Fish. and Ocean. (8): 236-270.
- VIDALIS, E. 1949: Contribution to the biology of the Anchovy (Engraulis encrasicolus L.) in Greek waters. Hellen. Hydrobiol. Inst., Prakt., 3 (2): 41-70.
- VLADYKOV, V. D. and D. H. WALLACE 1952: Studies of the striped bass, Roccus saxatilis (Waldbaum), with special reference to the Chesapeake Bay Region during 1936-1938. Bull. Bingham Ocean. Col. 14 (1): 132-177.
- VOOREN, C. M. 1972: The population dynamics of the New Zealand Tarakihi, Cheilodactylus macropterus (Bloch & Schneider), and changes due to fishing: an exploration. Oceanography of the South Pacific 1972: 493-503.
- WALFORD, L. A. 1932: The California Barracuda Sphyraena argentea. Calif. Fish Game, Fish. Bull. No. 37, 122 pp.
- WANATABE, H. 1958: Growth of the anchovy in the Japan Sea. Japan Sea Reg. Fish. Res. Lab., Ann. Rep. 4: 147-152.

- WEBER, W. 1970: Untersuchungen an den Beständen des Herings (Clupea harengus L.) der Westlichen Ostsee. Thesis, Kiel University, 85 pp.
- WEBER, W. and A. A. JOTHY undated: Observations on the fish Nemipterus spp. (Family: Nemipteridae) in the coastal waters of East Malaysia, 1972. Manuscript, with raw data pertaining to paper, same authors and title in Arch. FischWiss. 28 (2/3): 109-122.
- WELSH, W. and G. BREDER 1924: Contribution to the life history of Sciaenidae of the U.S. East Coast, U.S. Bur. Fish., Bull 39 (945): 141-201.
- WESTMAN, J. R. and P. W. GILBERT 1941: Notes on age determination and growth of the Atlantic bluefin tuna, Thunnus thynnus (Linnaeus). Copeia (2): 70-72.
- WESTMAN, J. R. and W. C. NEVILLE 1942: The tuna fishery of Long Island. New York, Board of Supervisors, Nassau County, 30 p.
- VON WESTERNHAGEN, H. V. and H. ROSENTHAL 1975: Rearing and spawning siganids (Pisces: Teleostei) in a closed seawater system. Helgoländer wiss. Meeresunters. 27: 1-18.
- VON WESTERNHAGEN, H. V. and H. ROSENTHAL 1976: Some aspects of the suitability of various Philippine Siganid species (Siganidae) for mariculture. Aquaculture 9: 297-311.
- WHEELER, J. and F. OMMANNEY 1953: Report on the Mauritius-Seychelles Fisheries Survey 1948-1949, Part 4. Fish. Publ., Lond. 1 (3): 120-140.

- WOHLSCHLAG, D. 1961: Growth of an Antarctic fish at freezing temperatures. *Copeia* 1961, 17-18.
- WOHLSCHLAG, D. 1962: Antarctic fish growth and metabolic differences related to sex. *Ecology* 43 (4): 589-597.
- WOJCIECHOWSKI, J. 1971: Some observations on the biology of Trichiurus lepturus L. from Cape Blane to Cape Verde Area. I.C.E.S. C.M. 1971/3:6.
- WONGSOMNUK, S., S. CHANTARAT and P. PONGSUWANA 1971: Experiments on Seabass (Lates calcarifer Bloch) culturing in brackish water ponds. 2nd. Symp. Mar. Fish. Marine Fish. Lab., Bangkok, MF/71-16, 12 pp.
- WÖRNER, F. 1975: Untersuchungen an drei Myctophidenarten, Benthoosema glaciale (Rheinhardt, 1837), Ceratoscopelus maderensis (Lowe, 1839) und Myctophum (M.) punctatum Rafinesque, 1810 aus dem Nordwestafrikanischen Auftriebsgebieten, im Frühjahr 1972. Thesis, Kiel University, 136 pp.
- WORTHMANN, H. 1975: Das Macrobenthos und Fischbesiedlung in verschiedenen Flachwassergebieten der Kieler Bucht (Westl. Ostsee). Thesis, Kiel University, 141 pp.
- YABUTA, Y. and M. YUKINAWA 1957: Age and growth of yellowfin tuna (Neothunnus macropterus) in Japanese waters by size frequencies. Rep. Nankai Reg. Fish. Res. Lab., (5): 127-133.
- YABUTA, Y., M. YUKINAWA and Y. WARASHINA, 1960: Growth and age of yellowfin tuna. 2. Age determination (scale method). Rep. Nankai Fish. Res. Lab., (12): 63-74.

- YAMADA and KAZIHARA 1955: Invest. Rep. Tsushima Warm Current. Proc. 3rd. Symp.
- YAMAGUTI, N. and E. P. DOS SANTOS 1966: Crescimento da pescada-goete (Macrodon ancylodon): Aspecto quantitativo. Bol. Inst. Ocean. (Sao Paulo) 15 (1): 75-78.
- YANG, R. T. 1970: Studies of age and growth (of) Atlantic albacore and a critical review on the structure, China Fisheries Monthly, No. 213, 9: 3-16.
- YASUDA, H. 1950: Growth of Japanese principal fish. III. Taius tumifrons (T. & S.), Bull. Jap. Soc. sci. Fish. 16 (6): 39-43.
- YASUDA, H. and A. KOIKE 1950: Growth of principal Japanese fish. II. Lateolabrax japonicus (C. & V.), Bull. Jap. Soc. sci. Fish. 16 (6): 256-258.
- YASUDA, H. and M. KOSAKA 1950: On the growth of Japanese principal fish. V. Nibea argentea (Houttuyn). Bull. Jap. Soc. sci. Fish. 15 (12): 859-861.
- YASUDA, H. and M. KOSAKA 1951: On the growth of the Japanese Principal Fish. IV. Branchiostegus japonicus (Houttuyn) and its allied forms. Bull. Jap. Soc. sci. Fish. 15 (12): 855-858.
- YASUDA, H. et al. 1962:  
cited (in Japanese) in Masuzawa (1967).
- YOKOTA, T. 1951: Studies on the stocks of sharks and rays.  
2. Age composition of the ray Dasyatis akajei (Müller & Henle) as observed in the catch by trawlers landed at Tororo, Miyazaki Prefecture, 1949 to May 1950. Bull. Jap. Soc. sci. Fish. 16: 188-189.

YOKOTA, T. and T. ASAMI 1956: Progress report on the results of coordinated research on the iwashi resources, 1953. Nankai Reg. Fish. Res. Lab.; 117 pp.

YOKOTA, T. and I. FURUKAWA 1952: Studies on the stocks of the clupeoid fishes in Hyuganada. III. Note on the variation of the number of vertebrae and monthly growth rate of the Japanese anchovy, Engraulis japonicus Temminck et Schlegel. Bull. Jap. Soc. sci. Fish. 17 (7-8): 60-64.

YOUNG, P. 1963: The Kelp Bass (Paralabrax clathrus) and its fishery 1947-1958. Calif. Fish Game, Fish. Bull. (122) 67 pp.

YUKINAWA, M. 1970: Age and growth of southern bluefin tuna Thunnus maccoyi by use of scale. Bull. Far East Fish. Res. Lab. (3): 229-257.

YUNOKAWA, Y. 1951: On the age and growth of Chelidonichthys kumu (Lesson and Garnot). Rec. Ocean. Works in Japan, (Spec. No. 5): 111-116.

ZAVODNIK, D. 1969: Studies on the life-history of Adriatic Sprat III on the Otoliths of the North Adriatic Sprat. Cons. gen. Pêches Medit. Studies and reviews, (40): 19-26.

ZEI, M. 1951: Monograph of the Adriatic species of Maenidae. Acad. Sc. Art. Slovenica, p. 3.

ZUPANOVIĆ, S. 1961: Contribution à la connaissance de la biologie des poissons de l'Adriatique. Acta Adriatica 9 (4): 1-84.

ZUPANOVIĆ, S. 1968: Study of the hake (Merluccius merluccius L.) biology and population dynamics in the central Adriatic. Conc. gen. Pêches Medit. Studies & Rev. (32), 24 pp.

ZUSSER, S. G. 1954: [The biology and production of Sarda sarda in the Black Sea.] Trud. vsesoiuz. nauch.-issled. morsk. ryb. Khoz., 27: 160-74 (In Russian).

2.5. Addenda to bibliography II

ALBERTI and ZUNIGA 1966: cited in Misu & Hamasaki 1971.

AWERINZEW, S. and N. SSYTSCH-AWERINZEWA 1935: Zur Biologie einiger Vertreter der Salmoniden, Familie aus dem Unterlaufe des Flusses Lena. Int. Rev. d. ges. Hydrobiol. u. Hydrogr. 32 (1/3): 49-102.

BAGENAL, T. B. 1970: Notes on the biology of the shelly Coregonus lavaretus L. in Haweswater and Hullswaters. J. Fish. Biol. 2 (1): 137-154.

BAS, C. 1963: Fluctuations de la pêche de Merlangus pou-tassou et quelques considerations sur son contrôle. Proc. tech. Pap. gen. Fish. Counc. Méditer. 7: 417-420.

EISELE 1943: Über das Wachstum der Coregonen im Voralpengebiet, insbesondere über das Verhältnis von Schuppen- und Längenwachstum. Z. Fisch. und d. Hilfswiss. 61 (1): 23-45.

FULTON, T. W. 1903: The distribution, growth and food of the angler (Lophius piscatorius). 21st. Ann. Rep. Fish. Bd Scotland, Part III: 186-217.

HABIB, G. 1976: Measuring growth in irregularly shaped otolith by means of a planimeter. Mauri Ora 4: 133-137 & Fish. Res. Publ. No 278, New Zealand.

HALLIDAY, R. 1970: Growth and vertical distribution of the glacier lanternfish, Bentho-sema glaciale, in the North-western Atlantic. J. Fish. Res. Bd Can. 27: 105-116.

HILE & DEASON 1934: Trans. Am. Fish. Soc. 64: 231.

- HOLT, S. J. 1960: A preliminary comparative study of the growth, maturity and mortality of sardine. Proc. World Scient. Meet. Biology Sardines and Related Species. FAO Rome, Vol. II, Subject Syn. (2): 553-561.
- JÄRVI, T. 1947; Über den Kleinmaränenbestand (Coregonus albula L.) in dem See Vesijärvi (Südfinnland). Acta Zool. Fennica 48: 43 pp.
- KOTO, T. 1963: Some consideration on the growth of marlins, using size frequencies in commercial catches. III attempts to estimate the growth of striped marlin, Tetrapterus audax (Philippi) in the Western North Pacific Ocean. Rep. Nankai Fish. Res. Lab. 17: 63-85.
- LASSERE, G. 1976: Thesis, Université du Languedoc, France.
- MATTHEW 1960: Synopsis on the biology of the South West African Pilchard (Sardinops ocellata Pappé). FAO Species Synopsis No 8 FB/60/S 8, 20 pp. in: Proceeding of the World Scient. Meet. Biol. Sardines and Related Species, Vol. II Rome (1960) p. 115-135.
- MORROW, J. 1951: The biology of the longhorn sculpin, Myoxocephalus octodecimspinosus Mitchill, with a discussion of the Southern New England "trash" fishery. Bull. Bingham Ocean. Coll. 13 (2) 89 pp.
- MÜLLER, W. 1960: Beiträge zur Biologie der Quappe (Lota lota L.) nach Untersuchungen in den Gewässern zwischen Elbe und Oder. Zeitschr. f. Fisch. u. d. Hilfswiss. IX (1): 1-72.
- NARAYANAN KUTTY 1961: Scales and otoliths of the "Koth", otolithoides brunneus (Day) as age indicator. Ind. J. Fish 8 (1): 145-151.



- NÜMAN, W. 1959: Das Wachstum der Blaufelchen und seine Berechnung bei Erfassung aller Individuen der einzelnen Jahrgänge in Abhängigkeit von der Stärke des Jahrganges, den jährlich variierenden Aussenfaktoren, der Eutrophierung und den regionalen Unterschieden des Sees. Arch. Fischwiss. 10 (1/2): 5-20.
- ROBERTSON, J. A. 1938: Fish Invest. London, Ser. 2, 16 (2).
- SURBECK, K. 1916: Beobachtungen und Untersuchungen an Sandfelchen des Bodensees. Allg. Fischerei Zeitung (11/12): 1-4.
- TABB 1961: A contribution to the biology of the spotted sea trout, Cynoscion nebulosus, in East Central Florida. Fla St. Bd. Cons. Techn. Ser (35): 1-23.
- THIENEMANN, A. 1935: Der Schnäpel (Coregonus lavaretus balticus) in Vorpommern. Dohrniana 14: 85-91.
- VAN OOSTEN 1939: The age, growth, sexual maturity and sex ratio of the common whitefish Coregonus clupeaformis Mitchill) of Lake Huron. Pap. Mich. Acad. Sci. Art & Letters 24 (II): 195-221.
- VINCENT-CUAZ, L. 1960: cited in: Vilela, H. and F. Frade 1963. Synopsis of biological data on yellowfin tuna Neothunnus albacora (Lowe) 1839 (Eastern Atlantic) FAO Fisheries Biology Synopsis No 76, 30 pp.